

## CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm  
**Comet<sup>®</sup> 400**

Herbicide

**ACTIVE CONSTITUENT: 400 g/L FLUROXYPYR present  
as the methylheptyl ester**  
**SOLVENTS: 316 g/L LIQUID HYDROCARBON  
100 g/L N-METHYL PYRROLIDONE**

GROUP  HERBICIDE

For the control of a wide range of Broadleaf Weeds in Fallow, Lucerne, Maize, Millets, Pastures, Poppies, Sorghum, Sugar Cane, Sweetcorn, Winter Cereals.  
Also for the control of Woody Weeds in Agricultural Non-Crop Areas, Commercial and Industrial Areas, Forests, Pastures and Rights-of-way, as specified in the Directions for Use.

**IMPORTANT: READ THE ATTACHED BOOKLET  
BEFORE USING THIS PRODUCT.**

Contents: 1 Litre  
5 Litres  
10 Litres  
20 Litres  
60 Litres  
110 Litres  
200 Litres  
500 Litres  
1000 Litres

Nufarm Australia Limited  
ACN 004 377 780  
103-105 Pipe Road  
Laverton North Victoria 3026  
Tel: (03) 9282 1000  
Fax: (03) 9282 1001



### **STORAGE AND DISPOSAL**

#### **1L, 5L, 10L, 20L, 200L**

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

#### **Refillable containers (500L, 1000L):**

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

#### **Envirodrum 60L, 110L Mini Bulk Returnable Container:**

Store the original sealed Envirodrum in a cool well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Nufarm Australia Limited.

### **SMALL SPILL MANAGEMENT**

Wear protective equipment (See SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal (See STORAGE AND DISPOSAL section). If necessary, wash the spill area with an alkali detergent and water and absorb as above the wash liquid for disposal.

### **SAFETY DIRECTIONS**

Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves, a face shield or goggles. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

### **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, DO NOT induce vomiting. Give a glass of water. If in eyes wash out immediately with water.

### **MATERIAL SAFETY DATA SHEET**

For further information refer to the Material Safety Data Sheet (MSDS), which can be obtained from your supplier or from the Nufarm website - [www.nufarm.com.au](http://www.nufarm.com.au)

### **CONDITIONS OF SALE**

"Any provisions or rights under the Trade Practices Act 1974 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute."

APVMA Approval No.: 61667/1L/0708

APVMA Approval No.: 61667/5L/0708

APVMA Approval No.: 61667/10L/0708

APVMA Approval No.: 61667/20L/0708

APVMA Approval No.: 61667/60L/0708

APVMA Approval No.: 61667/110L/0708

APVMA Approval No.: 61667/200L/0708

APVMA Approval No.: 61667/500L/0708

APVMA Approval No.: 61667/1000L/0708

### **In case of emergency: Phone 1800 033 498**

Ask for shift supervisor. Toll free 24 hours.

Batch Number:

Date of Manufacture:

Drummuster logo

Barcode

## CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm  
**Comet<sup>®</sup> 400**

Herbicide

**ACTIVE CONSTITUENT: 400 g/L FLUROXYPYR present  
as the methylheptyl ester**

**SOLVENTS: 316 g/L LIQUID HYDROCARBON  
100 g/L N-METHYL PYRROLIDONE**

GROUP  HERBICIDE

For the control of a wide range of Broadleaf Weeds in Fallow, Lucerne, Maize, Millets, Pastures, Poppies, Sorghum, Sugar Cane, Sweetcorn, Winter Cereals.

Also for the control of Woody Weeds in Agricultural Non-Crop Areas, Commercial and Industrial Areas, Forests, Pastures and Rights-of-way, as specified in the Directions for Use.

**IMPORTANT: READ THIS BOOKLET BEFORE USING THIS PRODUCT**

Nufarm Australia Limited  
ACN 004 377 780  
103-105 Pipe Road  
Laverton North Victoria 3026  
Tel: (03) 9282 1000  
Fax: (03) 9282 1001



APVMA Approval No.: 61667/0708

®Comet is a registered trademark of Nufarm Australia Limited.

**DIRECTIONS FOR USE:**

**RESTRAINTS:** DO NOT apply to plants which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (waterlogged or drought affected), poor nutrition, presence of disease, or previous herbicide treatment as reduced levels of control may result.

Thorough coverage of both foliage and stems, to the point of runoff, is essential for high volume applications (see GENERAL INSTRUCTIONS; application methods WOODY WEED SITUATIONS section). DO NOT spray if rain is likely within one hour.

Table 1	Woody Weeds in Agricultural Non-Crop Areas and Rights-of-way, Commercial and Industrial Areas, Forests and Pastures
Table 2	Established Grass Pastures (Ground and Aerial)
Table 3	Sorghum, Maize, Millets and Sweet corn
Table 4	Winter Cereals (Wheat, Barley, Oats and Triticale)
Table 5	Summer Fallow
Table 6	Winter Fallow
Table 7	Sugar cane
Table 8	Lucerne (established only)
Table 9	Poppies

**Table 1: Woody Weeds in Agricultural non-crop areas and Rights-of-way, Commercial and Industrial areas, Forests and Pastures**

Legumes present at the time of spraying will be severely damaged.

HIGH VOLUME APPLICATION: Dilute product with water See General Instructions - Application Method for application details				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /100L of water	CRITICAL COMMENTS
Bathurst burr Noogoora burr	Seedlings and young plants up to 40cm high	NSW, NT, Qld, WA only	38mL	
Black bindweed (Climbing buckwheat)	Seedlings and young plants before flowering	NSW, Qld only	150mL	
<i>Mimosa pigra</i>	Apply from mid to late summer	NT, WA only	250mL	Add Bonza Spraying Oil (see GENERAL INSTRUCTIONS; oils and surfactants)
Common sensitive plant	Seedlings and young plants up to flowering	Qld, WA only		
Bellyache bush		Qld, NSW, WA only		
Blackberry nightshade Bokhara clover		NSW, Qld only		
Caltrop (yellow vine) ( <i>Tribulus terrestris</i> ) ( <i>T. micrococcus</i> )		Seedlings and young plants up to 30cm diameter.		
Cobblers pegs	Up to 15cm high			
Cockspur thorn	Up to 3m high			
Creeping lantana	At flowering			
Crofton weed Mistflower	Seedlings and young plants up to flowering			
Docks ( <i>Rumex spp.</i> )	Seedlings and rosettes up to 30cm high			
Hexham scent	Seedlings and young plants up to flowering		Boom spray: Comet 400 at 300mL + 400mL of Amicide®625	
Honey locust	Seedlings and young plants up to 2m high		500mL	Apply to actively growing plants from October to April. Some regrowth may occur particularly when treating old woody plants with sparse canopies.
Small flowered mallow (Marshmallow) ( <i>Malva parviflora</i> )	Seedlings and young plants up to flowering			
Yellowflower Devil's claw	Seedlings and young plants up to flowering			
Lantana	Seedlings and regrowth 0.5 to 1.2m high Plants and regrowth 1.2 to 2m high			
Blue heliotrope	Flowering			
Limebush	Infestations up to 1.5m high only		250mL	
Madeira vine	Apply at time of active growth			
Milkweed ( <i>Euphorbia heterophylla</i> )	3 leaf to flowering	Qld only	500mL	Repeat applications will be necessary to control subsequent germinations.
Common sowthistle	Seedlings and young plants up to bolting	NSW, Qld only	250mL	Add Chemwet 1000 (See GENERAL INSTRUCTIONS; Oils and surfactants).
Mother-of-millions ( <i>Kalanchoe spp.</i> )	Seedling and young plants before flowering		300mL	

**Table 1: Woody Weeds in Agricultural non-crop areas and Rights-of-way, Commercial and Industrial areas, Forests and Pastures**

HIGH VOLUME APPLICATION: Dilute product with water See General Instructions - Application Method for application details				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /100L of water	CRITICAL COMMENTS
<b>Prickly acacia</b>	Seedlings and young plants up to 2m high	Qld only	375mL	Add Bonza Spraying Oil (see GENERAL INSTRUCTIONS; Oils and surfactants). Consult Tropical Weeds Research Centre, Charters Towers, for specific advice on application.
<b>Sida spp.</b>	Seedlings and young plants up to flowering	NSW, NT, Qld, WA only	500mL	
<b>Broadleaf Pepper tree (<i>Schinus terebinthifolius</i>)</b>	Mature leaves, fruiting	Qld only	250mL	Winter application only. Contact Alan Fletcher Research Station for more information.
<b>Flannel weed (<i>Sida cordifolia</i>)</b>				
<b>Snakeweed, (Dark and light blue)</b>	Seedlings and young plants before flowering		375mL	Add Bonza Spraying Oil (See GENERAL INSTRUCTIONS; Oils and surfactants).
<b>Stinking Passion Flower</b>	Established plants and regrowth	Qld, NT, WA only	225mL	Use 70mL/15L for a knapsack.
<b>Wandering jew (<i>Tradescantia albiflora</i>)</b>	Young plants up to and including flowering	All States	750mL	Some regrowth will usually occur and will require retreatment.
<b>Wattles (including <i>Acacia aulacocarpa</i>, <i>A. decora</i>, <i>A. harpophylla</i>, <i>A. leiocalyx</i>, <i>A. salicina</i>)</b>	Seedling plants or regrowth 0.5 to 1.2m high	NSW, Qld only	250mL	Apply to actively growing plants when soil moisture is plentiful. Some regrowth may occur particularly when treating old woody plants with sparse canopies and under dry conditions.
	Plants or regrowth 1.2 to 2.0m high only		500mL	

**Table 1: Woody Weeds in Agricultural non-crop areas and Rights-of-way, Commercial and Industrial areas, Forests and Pastures**

<b>BASAL BARK AND CUT STUMP APPLICATION: Dilute product with diesel</b> See General Instructions - Application Method for application details				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /100L of diesel</b>	<b>CRITICAL COMMENTS</b>
<b>Celtis</b> <i>(Celtis sinensis)</i>	<b>Basal Bark only:</b> Young plants up to 2m high and 20cm basal diameter	Qld only	1.8L	Treat stems from ground level to where multi-stemmed trunks branch.
<b>Chinee apple</b>	Up to 15cm basal diameter		1.5L	With basal bark, treat circumference of stem to height of 45cm from the ground. Contact The Land Protection Branch, Department of Natural Resources and Mines, Qld for further information on Chinee Apple.
<b>Cockspur thorn</b>	Basal Bark only: Up to 5cm basal diameter		1L	
<b>Mimosa bush</b> <i>(Acacia farnesiana)</i>	Up to 5cm basal diameter	Qld, WA only	1.5L	
<b>Prickly acacia</b>	Up to 10cm basal diameter	Qld only	750mL	
<b>Honey locust</b>	Plants up to 10cm basal diameter	Qld, NSW only	750mL	With basal bark, treat circumference of stem to height of 45cm from the ground. <b>For cut stump application</b> use a rate of 5L/100 diesel for all plant sizes. Contact The Land Protection Branch, Department of Natural Resources and Mines, Qld for further information on Honey Locust.
	Plants 10 to 20cm basal diameter		1.5L	
	Plants >20cm basal diameter		2.5L	
<b>Sisal hemp</b> <i>(Agave spp.)</i>	All growth stages	Qld only	1.5L	Treat as an overall spray. Contact The Land Protection Branch, Department of Natural Resources and Mines, Qld for advice to control large infestations.
			5mL undiluted product per plant	Lever out centre of plant with crowbar and immediately treat the exposed cut area.
<b>BROADCAST AND AERIAL APPLICATION: Dilute product with water</b> See General Instructions - Application Method for application details				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /ha</b>	<b>CRITICAL COMMENTS</b>
<b>Mimosa pigra</b>	Actively growing plants	NT, WA only	1.5L	Aerial application: Add Bonza Spraying Oil at the rate of 1L/100L spray mix. Apply to actively growing plants from mid to late summer. Contact The Department of Primary Industries and Fisheries, NT for further information.
<b>LOW VOLUME, HIGH CONCENTRATE APPLICATION: Using a drench gun or gas-powered gun</b> See General Instructions - Application Method for application details				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /10L of water</b>	<b>CRITICAL COMMENTS</b>
<b>Limebush</b>	Isolated bushes up to 1.2m high only	NSW, Qld only	500mL	Apply a 50mL dose per 5m <sup>2</sup> of bush surface area.
<b>Tree violet</b> <i>(Hymenanchera dentata)</i>	Apply from late flowering to green fruit up to 1.2m high	NSW only		Apply a 50mL dose per cubic metre of bush.

Table 2: Established Grass Pastures

WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha	CRITICAL COMMENTS
Blue billygoat weed Common sensitive plant Giant sensitive plant Spinyhead sida	Apply before flowering	Qld, WA only	750mL	Add Bonza Spraying Oil at 1L/ha
St John's wort	Apply from bud to full bloom (usually late Nov to early Jan)	ACT, NSW and Vic only	1.5L	Some regrowth will occur. Treat regrowth the following season for best results. Use at least 200L water/ha.
Silverleaf nightshade	From onset of flowering to early berry-set. (usually spring to mid-summer)	NSW only	375mL or 190mL + 1.2-1.6 Amicide 625	Add Bonza Spraying Oil at 1L/ha. To ensure maximum effect, delay application until the majority of shoots have emerged. Follow-up treatment of regrowth is critical for best control.

Table 3: Sorghum, Maize, Millets and Sweet Corn (NSW and QLD only)

CROP	CROP GROWTH STAGE	WEEDS CONTROLLED	WEED GROWTH STAGE	RATE /ha	CRITICAL COMMENTS
Sorghum	Apply when secondary roots are present, from 4 fully expanded leaves (15cm tall) up to boot (also see CRITICAL COMMENTS)	Annual ground cherry Wild gooseberry ( <i>Physalis</i> spp.)	2 to 8 leaf Up to 15cm tall	250mL	<p><b>Sorghum:</b> From 8 leaf to boot stage, use dropper nozzles to prevent herbicide coming in contact with the crop's leaves and the growing point (meristem).</p> <p><b>Maize and sweet corn:</b> From 6 leaf to just before tasselling, use dropper nozzles to prevent the herbicide coming in contact with the crop's leaves and the growing point (meristem).</p> <p><b>Millets:</b> DO NOT use mixes with Nu-trazine products.</p> <p><sup>1</sup> This treatment may be slightly damaging to the crop. To minimise crop damage apply using dropper nozzles at all crop stages.</p>
		Apple-of-Peru	Seedling plants up to 15cm tall	375mL	
		Bathurst burr Noogoora burr	2 to 8 leaf Up to 20cm tall	250mL	
Maize and Sweet corn	Apply when secondary roots are present, from 3 fully expanded leaves (10cm tall) up to just before tasselling (See CRITICAL COMMENTS)		20 to 50cm tall	375mL	
		Pigweed ( <i>Portulaca oleracea</i> )	Up to 10cm diameter	250mL	
			10 to 30cm diameter	375mL	
Millets	Spray when secondary roots have developed, usually early to mid-tillering, and not later than before heads start to form at the base of tillers. (See CRITICAL COMMENTS)	Sesbania pea	2 to 6 leaf Up to 10cm tall	750mL	
		Silverleaf nightshade (NSW only) <sup>1</sup>	Full flower to early berry	375mL + LI 700 at 300mL/100L	
		Starburr ( <i>Acanthospermum hispidum</i> ) (Qld only)	Up to 12 leaf and before flowering	750mL or 375mL + 1.6L Nu-trazine 600	
		Thornapples ( <i>Datura</i> spp.)	2 to 8 leaf Up to 15cm tall	375mL	
		Volunteer sunflower	2 to 5 leaf Up to 20cm tall	500mL	

Table 3 (Continued): Sorghum, Maize, Millets and Sweet Corn (NSW and QLD only)

CROP	CROP GROWTH STAGE	WEEDS CONTROLLED	WEED GROWTH STAGE	RATE /ha	CRITICAL COMMENTS
<b>Comet 400 in tank-mixes with Nu-triazine: Sorghum, Maize and Sweet Corn</b>					
Sorghum, Maize and Sweet corn (continued)	See Above	Amaranthus spp. including: Boggabri weed, Dwarf amaranth, Green amaranth Redshank Anoda weed Bladder ketmia Black pigweed ( <i>Trianthema portulacastrum</i> ) Caltrop (yellow vine) including: <i>Tribulus terrestris</i> , <i>T.micrococcus</i> and <i>T.maximus</i> Cowvine (peach vine) ( <i>Ipomoea lonchophyllia</i> ) Hairy wandering jew ( <i>Commelina benghalensis</i> ) Mintweed	Seedling plants up to 15cm tall or rosettes up to 15cm diameter	<b>Low Rate</b> 250mL + 1.25L of Nu-triazine 600 or 675g of Nu-triazine 900DF <b>OR</b> <b>High Rate</b> 375mL plus 1.6L of Nu-triazine 600 or 1.1kg of Nu-triazine 900DF	<b>Use the low rate</b> (250mL + 1.25L) when weeds are small (5-7cm tall/diameter). <b>Use the high rate</b> (375mL + 1.6L) when the weeds are larger (7-15cm tall/diameter). Comet 400 is generally more compatible with <b>Liquid Nu-triazine products</b> (see GENERAL INSTRUCTIONS; compatibility section). <b>Add a surfactant</b> (see GENERAL INSTRUCTIONS; Oils and surfactants). DO NOT add an oil to mixtures of Comet 400 and Nu-triazine 600/Nu-triazine 900DF.
		<i>Euphorbia davidii</i>	Cotyledons to 4 nodes up to 15cm	500mL + 1.6L of Nu-triazine 600 or 1.1kg of Nu-triazine 900DF	
		Volunteer peanuts	Up to 15cm diameter	500mL + 3.75L of Nu-triazine 600 or 2.5kg of Nu-triazine 900DF	
<b>Sweet Corn: Tasmania only</b>					
Sweet corn only	3 to 5 leaf	Blackberry nightshade Volunteer potatoes	3 to 5 leaf	500mL	

Table 4: Winter Cereals (Wheat, Barley, Oats and Triticale)

CROP GROWTH STAGE	WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha	CRITICAL COMMENTS
Apply from 3 leaf to flag (Zadoks 13 to 39)	Bedstraw ( <i>Galium tricornutum</i> )	1 to 3 whorl	Vic, SA, WA only	500mL	². Add either LI 700 or Activator Surfactant (see GENERAL INSTRUCTIONS; Oils and Surfactants).
	Cleavers ( <i>Galium aparine</i> )		NSW, Vic only		
	Black bindweed (Climbing buckwheat)	2 to 4 leaf	NSW, Qld only	250mL²	Useful suppression only.  <b>Mixtures:</b> Mixing partners with Comet 400 may reduce crop selectivity. Apply at crop growth stages according to the mixing partner's recommendation.
		2 to 6 leaf			
	Common sowthistle ( <i>Sonchus oleraceus</i> )	2 to 5 leaf		500mL	
	Deadnettle	2 to 6 leaf		750mL or 250mL + 5g Associate/Brushkiller²	
	Spiny emex (Doublegee, Three cornered jack)	2 to 4 leaf	NSW, SA, Qld, WA only	500mL	Plants 15 to 30cm tall will only be suppressed.
	Prickly lettuce	2 to 5 leaf	NSW, Qld, Tas, Vic, WA only		
	Volunteer lupins	2 to 8 leaf	NSW, Vic, WA only	750mL	
	Volunteer potato	10 to 15cm tall	WA and Tas only		
	Wireweed	2 to 3 leaf	NSW, Qld, SA, Tas, Vic, WA only	250mL + 5g Associate/Brushkiller²	
			NSW and Qld only		
Bittercress ( <i>Coronopus didymus</i> ) Mustards Shepherd's purse Turnip weed Wild radish Wild turnip	Up to 8 leaf and up to 15cm diameter	Qld, NSW, Vic, SA, Tas, WA only	250mL to 750mL + Associate/Brushkiller² or Eclipse² or Nufarm LVE MCPA or Nufarm Agritone 750	The Comet 400 rate depends on what other weeds are present as listed above. See Mixtures comment above. Associate²/Brushkiller™ @ 5g/ha (This mix does not control wild radish). Eclipse* @ 5-7g/ha (use the 5g rate on turnip weed only). LVE MCPA @ 700mL/ha Agritone 750 @ 660mL/ha.	

Table 5: Summer Fallow

WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha	CRITICAL COMMENTS
Annual ground cherry Wild gooseberry ( <i>Physalis</i> spp.)	2 to 8 leaf, up to 15cm tall	NSW, Qld only	375mL <sup>3</sup>	<sup>2</sup> Add Bonza Spraying Oil (see GENERAL INSTRUCTIONS; Oils and surfactants section)
Bathurst burr Noogoora burr	2 to 8 leaf, up to 20cm tall	NSW, Qld, Vic, WA only		
Bellvine	Pre-flowering	NSW, Qld only	250mL + 1.2L Nufarm Glyphosate CT or Roundup <sup>®</sup> CT or 1L Credit <sup>®</sup> plus 1L Bonus <sup>®</sup> or 1L Roundup PowerMAX <sup>™</sup>	When mixing with Nufarm Glyphosate CT, Roundup CT, Credit plus Bonus or Roundup PowerMAX to control both grass and broadleaf weeds, refer to the label for use rates and adjuvants recommended for the grasses (see GENERAL INSTRUCTIONS; Compatibility Section). DO NOT add an adjuvant to mixtures containing Credit and Bonus.
Bladder ketmia	4 to 8 leaf, up to 10cm tall			
Cowvine (Peach vine) <i>Ipomoea lonchophylla</i>	2 to 10 leaf up to 10cm diameter			
Caltrop (Yellow vine) including: <i>Tribulus terrestris</i> , <i>T. maximus</i> and <i>T. micrococcus</i>	Up to 15cm diameter			
Pigweed ( <i>Portulaca oleracea</i> )	Up to 10cm diameter			
	Up to 60cm diameter	375mL + 1L Nufarm Glyphosate CT or Roundup <sup>®</sup> CT or 830mL Credit plus 830mL Bonus <sup>®</sup> or 830mL Roundup PowerMAX <sup>™</sup>		
<i>Polymeria pusilla</i>	2 to 10 leaf up to 20cm diameter		500mL <sup>2</sup> or 250mL + 1.2L Nufarm Glyphosate CT or Roundup <sup>®</sup> CT or 1L Credit <sup>®</sup> plus 1L Bonus <sup>®</sup> or 1L Roundup PowerMAX <sup>™</sup>	<sup>3</sup> Delay treatment until the maximum number of shoots have emerged, but before the onset of fruiting (late summer) DO NOT treat plants showing symptoms from previous treatment. Use the high rate when longer term weed control (6-10 months) is required and delay planting crops during this period. The low rate will require follow-up treatments.
Rhynchosia	Seedlings to early flowering		500mL <sup>2</sup> or 190mL + 800mL Nufarm Glyphosate CT or Roundup <sup>®</sup> CT or 670mL Credit <sup>®</sup> plus 670mL Bonus <sup>®</sup> or 670mL Roundup PowerMAX <sup>™</sup>	
Smallflower mallow or Marshmallow ( <i>Malva parviflora</i> )	Up to 8 leaf up to 20cm		500mL <sup>2</sup>	
Thornapples ( <i>Datura</i> spp.)	2 to 8 leaf up to 15cm diameter	NSW, Qld, WA only	375mL <sup>2</sup> or 250mL + 1.2L Nufarm Glyphosate CT or Roundup <sup>®</sup> CT or 1L Credit <sup>®</sup> plus 1L Bonus <sup>®</sup> or 1L Roundup PowerMAX <sup>™</sup>	
Sesbania pea	2 to 6 leaf up to 10cm tall	NSW, Qld only	750mL <sup>2</sup> or 250mL + 1.2L Nufarm Glyphosate CT or Roundup <sup>®</sup> CT or 1L Credit <sup>®</sup> plus 1L Bonus <sup>®</sup> or 1L Roundup PowerMAX <sup>™</sup>	

Continue over

Table 5: Summer Fallow

WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha	CRITICAL COMMENTS
<b>Perennial Ground Cherry</b> <i>(Physalis virginiana)</i> <sup>1</sup> :	Bud to early flowering up to 20cm tall	NSW, Qld only	750mL or 1.5L	Refer to critical comments on previous page
<b>Silverleaf nightshade</b>	Full flower to early berry-set (usually Dec - Feb)	NSW only	375mL or 190mL + 1.2L-1.6L Amicide® 625	Add LI 700 at the rate of 300mL/100L spray mixture  To ensure maximum effect, delay application until the majority of shoots have emerged. Follow-up treatment will be required to control regrowth and is critical for optimal control. If wanting to prevent seed set repeat applications may be needed in the same season, although this does not lead to better long term control.
<b>Volunteer peanuts</b>	Up to 15cm diameter	Qld only	500mL + 3.75L Nu-trazine 600	Add Chemwet 1000 (see GENERAL INSTRUCTIONS; Oils and surfactants) Important: See GENERAL INSTRUCTIONS; Compatibility section.
<b>Volunteer sunflowers</b>	2 to 5 leaf up to 20cm	NSW, Qld only	500mL	Add LI 700 (see GENERAL INSTRUCTIONS; Oils and surfactants section).

Table 6: Winter Fallow

WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE /ha	CRITICAL COMMENTS
<b>Bedstraw</b> <i>(Galium tricornutum)</i>	Up to 5 whorl	Vic, SA, WA only	500mL <sup>2</sup>	<sup>2</sup> Add LI 700 (see GENERAL INSTRUCTIONS; Oils and surfactants section) <sup>3</sup> Add LI 700 or Chemwet 1000 (see GENERAL INSTRUCTIONS; oils and surfactants section)
<b>Cleavers</b> <i>(Galium aparine)</i>		NSW, Vic only		
<b>Black bindweed</b> (Climbing buckwheat)	2 to 8 leaf up to 10cm diameter	NSW, Qld only	375mL <sup>2</sup>	When mixing with Nufarm Glyphosate CT or Roundup CT to control both grass and broadleaf weeds, refer to the label for use rates and adjuvants recommended for the grasses (see GENERAL INSTRUCTIONS; Compatibility Section)  DO NOT add an adjuvant to mixtures containing Credit® and Bonus®.
<b>Common sowthistle</b> <i>(Sonchus oleraceus)</i>	2 to 5 leaf up to 10cm diameter		500mL <sup>2</sup> or 250mL + 600mL Nufarm Glyphosate CT or Roundup <sup>^</sup> CT or 500mL Credit® plus 500mL Bonus® or 500mL Roundup <sup>^</sup> PowerMAX™	
<b>Prickly lettuce</b>				
<b>Spiny emex</b> (Doublegee, Three cornered jack)	2 to 8 leaf		750mL <sup>2</sup> or 250mL <sup>3</sup> + 5g Associate/Brushkiller	
<b>Wireweed</b>	2 to 3 leaf up to 10cm tall		750mL <sup>2</sup> or 250mL <sup>3</sup> + 5g Associate/Brushkiller or 500mL <sup>3</sup> + 600mL Nufarm Glyphosate CT or Roundup <sup>^</sup> CT or 500mL Roundup <sup>^</sup> PowerMAX™	

**Table 7: Sugar cane (Qld, NSW, NT and WA only)**

CROP GROWTH STAGE	WEED CONTROLLED	WEED GROWTH STAGE	RATE /ha	CRITICAL COMMENTS
From early tillering to maturity	Balsam pear Blackberry nightshade Blue billygoat weed Centro Cowpea Giant sensitive plant Lablab bean Noogoora burr Phasey bean Pinkburr Prickly African cucumber Spinyhead sida Stinking passion flower (seedlings only)	Apply from 2 to 3 leaf until flowering	Ground: 650mL  Aerial: 750mL	For optimal weed control, delay application until just before the “close-in” stage <b>Aerial application:</b> Apply in not less than 60L/ha water and add LI 700 at 300mL/100L spray mixture. <b>Ground application:</b> Apply in 100-400L/ha water and add LI 700 at 300mL/100L of spray mixture.
	Bellvine Morning glory Red or pink convolvulus Star-of-Bethlehem		As above + 800mL Amicide 625	
	Stinking passion flower	Established or ratoon plants with at least 1.0m of regrowth	<b>High volume:</b> 225mL/100L water  <b>Knapsack:</b> 35mL/15L water	Thoroughly wet plants to the point of run-off.
	Milkweed ( <i>Euphorbia heterophylla</i> )	Seedlings and young plants up to flowering	1.5L or 1.15L + 3.3L Nu-trazine 600	Better control will be achieved with the Nu-trazine mixture. Delay application until just before the cane reaches the “close-in” stage. This will improve control and minimise the number of seedlings that germinate.

**Table 8: Lucerne (NSW Only)**

CROP GROWTH STAGE	WEED CONTROLLED	WEED GROWTH STAGE	RATE /ha	CRITICAL COMMENTS
Established crops at least eighteen months old	Annual ground cherry Bathurst burr Noogoora burr Wild gooseberry	2 to 8 leaf up to 15cm high	250mL	To minimise crop injury and to maximise weed control, cut, slash or heavily graze the lucerne before application. Wherever possible, irrigate before application to stimulate weed growth.  DO NOT treat crops growing on sandy or stony soils. DO NOT treat crops after the summer growing season (after end of March). To broaden the spectrum of weeds controlled, Comet 400 can be mixed with Buttress®.
	Pigweed	Up to 10cm diameter		

**Table 9: Poppies (TAS only)**

CROP GROWTH STAGE	WEED CONTROLLED	WEED GROWTH STAGE	RATE /ha	CRITICAL COMMENTS
4 to 6 leaf	Cleavers Fumitory	2 to 6 leaf	500mL	
	Shepherd’s purse Wireweed		500mL+5L Asulox*	
8-10 leaf	Common sowthistle Prickly lettuce	2 to 5 leaf	500mL	DO NOT apply Comet 400 to poppies later than the 8 to 10 leaf growth stage as a reduction of alkaloid content could occur.
	Black nightshade	cotyledon to 4 leaf	750mL	
	Fumitory	6 to 10 leaf		This rate will provide season long control of volunteer potato, but will not control all daughter tubers and will only suppress potatoes over 15cm tall.
	Volunteer potato	From tuber initiation to flower bud		

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIODS:**

**Grazing:**

**DO NOT GRAZE FAILED CROPS AND TREATED PASTURES OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.**

**Harvest:**

**Poppies - DO NOT SPRAY POPPIES LATER THAN 10 WEEKS BEFORE HARVEST**

**Other Crops - Not required when used as directed**

**MINIMUM RECROPPING PERIODS**

**Plant-back periods for crops following the application of Comet 400 for rates up to 750mL/ha.**

RATE /HA	190mL	375mL	750mL
CROP	DAYS		
Barley	7	7	7
Wheat	7	7	7
Chickpea	7	7	7
Cotton	14	14	28
Soybean	7	7	14
Sunflower	7	7	7
Maize	7	7	7
Sorghum	7	7	7

**Note: Before using Comet 400 in tank mixes with other herbicides, check the plant-back information on all product labels. The time between spraying and planting will be determined by the most residual product, ie. the product with the longest plant-back period.**

**GENERAL INSTRUCTIONS:**

**MIXING**

Comet 400 may be mixed with water or diesel. Mix only sufficient chemical for each days use and avoid storing.

**Mixing in Water:** Half fill the spray tank with water and add the required quantity of Comet 400 and complete filling. Agitate continuously to ensure thorough mixing before and during application.

**Mixing in Diesel:** Half fill the spray tank with diesel and add the required quantity of Comet 400. Add the remainder of the diesel and agitate or shake to mix contents.

**Tank mixtures:** Wettable powder or dry flowable formulations (eg water dispersible granules) should be added to the spray tank first, followed by suspension concentrates (Flowables), water soluble salts and then emulsifiable concentrate formulations (Comet 400). Add spraying oils and surfactants (wetters) last.

**OILS AND SURFACTANTS**

**Oils:**

Where specified use Bonza Spraying Oil at the rate of 500mL/100L of spray mix. When using less than 100L/ha spray volume, ensure a minimum of 250mL/ha of Bonza Spraying Oil is used, unless 1L/100L or 1L/ha is specified.

**Surfactants (wetters):**

LI 700 to be used at 300mL/100L.

Spraymate Activator to be used at 125mL/100L.

Use a 100% concentrate non-ionic surfactant such as Chemwet 1000 at 100mL/100L of spray mix where required.

**COMPATIBILITY**

Comet 400 is compatible with the herbicides listed. Follow any regional restrictions, and all directions and restrictions on the label, of any product mixed with Comet 400.

Nu-triazine 900DF	Topik* 240 EC (see below)
Nu-triazine 600	
Associate®/Brushkiller™	Tordon* 75-D
	Trooper
Broadstrike*	Touchdown*
Eclipse*	Amicide® 625
Nugrass®	Buttress®
Invader® 600	
Archer™	Credit®
Agritone 750	Bonus®
	Nufarm LVE MCPA
Nufarm Glyphosate CT	Roundup^CT
	Roundup PowerMAX™

**NU-TRAZINE**

AVOID USING HARD WATER WHEREVER POSSIBLE. Where hard water cannot be avoided, the addition of a water conditioning agent to the spray tank, at 100g/100L water, before adding any herbicide may improve compatibility.

AGITATION IS VERY IMPORTANT WHEN MIXING COMET 400 AND NU-TRAZINE.

Comet 400 plus Nu-triazine tank mixes **must be**

**agitated vigorously and continuously during mixing and application. After mixing DO NOT allow to stand without agitation. Ensure that the time from mixing to the end of application is not more than 2 hours.**

If settling out occurs re-suspension is difficult, even with vigorous agitation. Agitation using only the pump's by-pass is usually inadequate, particularly with larger tanks (more than 2000L). Additional mechanical agitation will be necessary in large tanks, computer sprayers and mixing tanks.

When additional surfactant is required, add Chemwet 1000 at 100mL/100L of spray mix. DO NOT use a spraying oil when tank mixing Comet 400 and Nu-triazine.

Guidelines for tank-mixing Comet 400 and Nu-triazine formulations:

Tank Mix	Rate /ha	Water Hardness		Min. Water Volume (L/ha)		Comments
		Soft	Hard (1026 ppm ca)	Ground	Aerial	
Comet 400 + Nu-triazine 900 DF	375mL + 1.1	✓	✓	100	Do not use	Sediment may be difficult to resuspend and may block nozzles
Comet 400 + Nu-triazine 600	375mL + 1.6	✓	✓	100	Do not use	Sediment may be difficult to resuspend and may block nozzles

**Topik 240 EC**

Always use Bonza with Comet 400 + Topik 240 EC tank-mixes at 500mL/100L of spray mix with a minimum of 250mL/ha.

DO NOT mix Comet 400 with Topik 240 EC if the grass weeds are not actively growing. Always use the maximum label rate of Topik 240 EC for the appropriate grass growth stage.

DO NOT use Comet 400 at more than 0.75L/ha in tank mixes with Topik 240 EC.

**Nufarm Glyphosate CT/Roundup CT/Credit plus Bonus/Roundup PowerMAX**

When mixing Comet 400 with one of these products to control both grass and broadleaf weeds, refer to that label for use rate and adjuvants recommended for grasses. DO NOT use Nufarm Glyphosate CT or Roundup CT at less than 1.2L/ha in tank mixes with Comet 400 or Credit plus Bonus at less than 1L/ha of each or Roundup PowerMAX at less than 1L/ha, when banyard grass, buttongrass, crowsfoot grass, native millet and liverseed grass are the target species.

**APPLICATION METHODS and WATER RATES**

**BROADCAST APPLICATION IN CROPPING, PASTURE AND FALLOW SITUATIONS.**

**A. Ground application (Boom)**

Apply Comet 400 with an accurately calibrated boom sprayer, in at least 50L/ha water (100-400L/ha for sugar cane).

Flat fan nozzles are recommended using pressures in the range 200 to 300kPa.

Set the boom at a height to ensure a double overlap of the nozzle patterns.

**B. Ground direct application (Dropper nozzles)**

To minimise crop effects, dropper nozzles should be used in sorghum when the crop is beyond the 8 leaf growth stage and in maize and sweet corn when the crop is beyond the 6 leaf growth stage.

Adjust the nozzles to direct the spray into the base of the crop and away from the leaves and the growing point. See manufacturers directions for setting up and calibration of dropper nozzles.

**C. Aerial Application**

Apply in a minimum volume of at least 35L/ha water (60L/ha in sugar cane)

Use equipment calibrated to produce droplets with an average diameter (Volume Mean Diameter: VMD) of 250-350 microns.

DO NOT apply when the temperature is above 30°C, when there is no wind or when the wind is blowing toward susceptible crops.

DO NOT use human flaggers unless they are protected by engineering controls such as enclosed cabs.

**WOODY WEED SITUATIONS**

Weeds must be actively growing to attain optimal effect. Delay the treatment of regrowth following bulldozing, slashing, burning, ploughing or a previous chemical treatment until it has at least 1 metre of new, vigorous growth.

**A. High Volume Application**

**Hand Gun**

Apply the recommended mix to obtain full coverage of leaves and stems using a number 6-8 tip at 700 to 1500 kPa. To obtain good coverage, a spray volume of 1500 to 4000L/ha (15 to 40L/100m<sup>2</sup>) is required per infested hectare.

Ensure thorough coverage to the point of runoff.

**Knapsack**

Knapsack sprayers may be used on smaller infestations where penetration and coverage of the canopy is easier to achieve. Use the same use rate and spray techniques as for handgun application.

**B. Low Volume, High Concentrate Application  
Drench Gun or Grass-Powered Gun.**

Apply the recommended mixture uniformly across the foliage by applying 50mL shots to cover 4 to 5m<sup>2</sup> of surface area of plant. This is approximately equivalent to 20 droplets per cm<sup>2</sup> of the leaf surface. Use a marking agent as recommended by the equipment manufacturer to check spray coverage.

**C. Basal Bark and Cut Stump Application  
Basal Bark**

DO NOT apply to wet stems as this can repel the diesel mixture.

Spray or paint the recommended mixture around the

base of each stem from ground level to a height of at least 30cm from the ground, wetting the bark to the point of runoff.

Apply with a paint brush or a pressure sprayer with an appropriate lance and solid cone nozzle. If using spray equipment use low pressures (≤ 200 kPa) sufficient to form a cone of spray.

Old rough bark will require more spray than smooth or young thin bark.

**Cut Stump**

Apply the recommended mixture liberally to the freshly cut stump immediately after cutting.

Apply by spraying or painting the cut surface and sides of the stump.

Best results are obtained when the stems are cut less than 15cm above the ground.

**CLEANING SPRAY EQUIPMENT**

**Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto wasteland away from desirable plants and water courses.**

**Cleaning equipment after using water-based sprays:**

**Rinsing:** After using Comet 400 Herbicide, empty the tank completely and drain the whole system.

Thoroughly wash inside the spray unit using a pressure hose. Drain, and clean any filters in the tank, pump, lines, hoses and nozzles.

After cleaning the tank as above, quarter fill with clean water and circulate through the pump, lines and nozzles. Drain and repeat the rinsing procedure twice.

**Decontamination (before spraying cotton and other sensitive crops; see PROTECTION OF CROPS):**

Wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent such as Spraymate Tank and Equipment Cleaner at 500mL/100L of water and circulate throughout the system for at least 15 minutes.

Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.

**Cleaning equipment after using diesel-based sprays:**

On completion of spraying, use a degreaser such as Caltex Kwik-D-Grease\* to remove traces of diesel from the sprayer. Rinse tank and spray through nozzles with water to remove degrease.

Then, quarter fill the tank with clean water and add an alkali detergent such as Spraymate Tank and Equipment Cleaner at 50mL/10L of water. Shake sprayer to circulate the washing solution throughout the sprayer, then spray the solution through the nozzles. Rinse well with clean water to remove the detergent.

To clean brushes and containers, spray liberally with degreaser. Hose off with clean water and repeat using detergents as above.

DO NOT use this equipment for any other purpose.

## RESISTANT WEEDS WARNING

**GROUP I HERBICIDE**

Comet 400 Herbicide is a member of the pyridine group of herbicides. Comet 400 has a disrupters of plant cell growth mode of action. For weed resistance management Comet 400 is a Group I Herbicide. Some naturally occurring weed biotypes resistant to Comet 400 and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Comet 400 or other Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from the failure of this product to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department or Agriculture, or Nufarm Australia Limited representative.

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS.

Susceptible crops include but are not limited to clovers, cotton, fruit, hops, lupins, ornamentals, peas, pine trees, potatoes, navy beans, safflower, shade trees, soybeans, sunflower, tobacco, tomatoes, vegetables, and vines.

Comet 400 can be damaging to susceptible crops during growing and dormant periods.

Grasses are normally unaffected by Comet 400 and establish quickly after treatment. Transitory damage can occur on some species particularly those that spread by stolons such as couch grass (*Cynodon dactylon*), Kikuyu grass and carpet grass (*Axonopus* sp.).

DO NOT allow spray to drift onto susceptible crops, shade trees and *Pinus* spp.

DO NOT use under weather conditions or from spraying equipment which could be expected to cause spray to drift onto nearby susceptible plants.

## PROTECTION OF LIVESTOCK

DO NOT graze or cut treated crops for stock food except as specified under withholding periods.

Poisonous plants may become more palatable after spraying.

DO NOT allow stock to re-enter paddocks containing treated poisonous plants, until the plants have died down.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

Alongside waterways, treat only noxious weeds and poisonous plants.

## STORAGE AND DISPOSAL

**1L, 5L, 10L, 20L, 200L**

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

## Refillable containers (500L, 1000L):

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

## Envirodrum 60L, 110L Mini Bulk Returnable Container:

Store the original sealed Envirodrum in a cool well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Nufarm Australia Limited.

## SMALL SPILL MANAGEMENT

Wear protective equipment (See SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal (See STORAGE AND DISPOSAL section). If necessary, wash the spill area with an alkali detergent and water and absorb as above the wash liquid for disposal.

## SAFETY DIRECTIONS

Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves, a face shield or goggles. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, DO NOT induce vomiting. Give a glass of water. If in eyes wash out immediately with water.

## MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS), which can be obtained from your supplier or from the Nufarm website - [www.nufarm.com.au](http://www.nufarm.com.au)

**CONDITIONS OF SALE**

“Any provisions or rights under the Trade Practices Act 1974 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded.

Nufarm Australia Limited and/or its affiliates (“Nufarm”) shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute.”

APVMA Approval No.: 61667/0708

**In case of emergency: Phone 1800 033 498**

Ask for shift supervisor. Toll free 24 hours.

™ Trademark pending

® Amicide, Associate, Comet, Nugrass, Invader, Buttress, Credit, Bonus are trademarks of Nufarm Australia Limited.

^ Roundup is a registered trademark of Monsanto Technology LLC used under licence by Nufarm Australia Limited.

\* Other registered trademarks