

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Antracol[®]

FUNGICIDE SPRAY

Active Constituent: 700 g/kg PROPINEB

GROUP Y FUNGICIDE

For the control of blue mould of tobacco, target spot (early blight) and late blight (Irish blight) of potatoes and tomatoes, grey leaf spot of tomatoes, downy mildew of cucurbits and onions, black spot of citrus and certain other diseases

GENERAL INSTRUCTIONS

Fungicide Resistance Warning

Antracol is a member of the multi-site group of fungicides. For fungicide resistance management Antracol is a Group Y fungicide. Some naturally occurring individual fungi resistant to the product and other Group Y fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product and other Group Y fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that result from failure of this product to control resistant fungi.

Export of treated produce

Growers should note that MRLs or import tolerances do not exist in all markets for edible produce treated with Antracol. If you are growing edible produce for export, please check with Bayer CropScience Pty Ltd for the latest information on MRLs and import tolerances before using Antracol.

Mixing

To the required quantity of Antracol add a small amount of water and stir to make a paste. Add further water and mix until a cream results. Pour this into water in spray vat while stirring or with agitators in motion.

Compatibility

Ensure that water is of suitable quality for mixing with agricultural pesticides. Poor quality water can cause physical incompatibility between some products which may result in crop damage in some circumstances.

Antracol is compatible with most commonly used insecticides and fungicides including Bayfidan[®] 250 EC, Diptorex[®] 500 SL, Folimat[®] 800, Lebaycid[®], Morestan[®], Nitofol[®]. Antracol is not compatible with lime sulphur and other alkaline materials. When mixing sprays, except when adding oil, add insecticides last. Do not mix concentrates together but add each to the spray tank separately. The compatibilities given are based on world wide experience and are considered correct. However, as changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays, we accept no responsibility for the behaviour of such mixtures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well ventilated area out of direct sunlight. Single rinse before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Repeated exposure may cause allergic disorders. Interacts with alcohol - avoid alcohol on days of use. Avoid contact with eyes and skin. Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (telephone 13 11 26). Avoid giving alcohol.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from www.bayercropscience.com.au.

EXCLUSION OF LIABILITY

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

Antracol® is a registered trademark of Bayer

NRA Approval Number 30613/1202

FOR 24 HOUR SPECIALIST ADVICE
IN EMERGENCY ONLY
PHONE 1800 033 111

DIRECTIONS FOR USE

CROP	DISEASE/PEST	STATE	RATE	WHP (DAYS)	CRITICAL COMMENTS
Celery	Septoria leaf spot	Vic, Tas, WA only	200 g/100 L or 2 kg/ha	7	Apply when conditions favour disease and repeat at 7-10 day intervals.
Citrus	Black spot	NSW, Vic, SA, WA only	200 g/100 L plus white oil 250 mL/100 L	7	Apply 6 weeks after copper spray at petal fall and 12 weeks after petal fall. Coastal lowlands and highlands (NSW): apply 8 and 14 weeks after the petal fall copper spray.
	Citrus rust mite	NSW, Vic, SA, WA only			
	Black spot Citrus rust mite	Qld only	200 g/100 L plus white oil 600 mL/100 L		
Cucurbits eg: cucumbers, rockmelons	Downy mildew	All States	200 g/100 L or 2 kg/ha	3	Apply at weekly intervals while conditions favour disease. Spray both sides of leaves thoroughly.
Lettuce	Downy mildew	Vic, Tas, WA only	200 g/100 L or 2 kg/ha	3	Apply when conditions favour disease and repeat at 7-10 day intervals.
Onions	Downy mildew	All States	200 g/100 L or 2 kg/ha	14	Apply at intervals of 7-10 days from early in the life of the crop. If conditions favour disease development consider using a systemic fungicide. Add wetting agent.
Potatoes	Target spot (Early blight) Irish blight (Late blight)	All States	200 g/100 L or 2 kg/ha	1	Apply when conditions favour disease and repeat at 7-10 day intervals.
Tobacco	Blue mould	Vic, WA only	200 g/100 L or 2 kg/ha	7	Apply at weekly intervals commencing 7 days after transplanting and terminating 7 days before first pick. Apply 350 L spray mix per ha after transplanting, increasing to 1350 L on a fully grown crop.
		Qld, NSW, WA only			Apply at weekly intervals terminating 7 days before first pick.
	Frogeye leaf spot Brown spot	Qld, WA only			
Tomatoes	Target spot (Early blight) Irish blight (Late blight) Grey leaf spot (Stemphylium)	All States	200 g/100 L or 2 kg/ha	3	Apply when conditions favour disease and repeat at 7-10 day intervals. In places where diseases are severe commence at seed bed stage.
Watermelon Rockmelon	Anthracnose Gummy stem blight	Qld, WA only	200 g/100 L or 2 kg/ha	3	Apply every 7-10 days depending on weather and disease incidence. First application may be made when vines begin to run.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

Potatoes: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION

Cucurbits, Lettuce, Tomatoes: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION

Celery, Citrus, Tobacco: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION

Onions: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION