

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

MACSPRED

EUCMIX™GR GRANULAR HERBICIDE

ACTIVE CONSTITUENTS: 44 g/kg TERBACIL
2 g/kg SULFOMETURON METHYL

GROUP	B C	HERBICIDE
-------	-----	-----------

For the control of certain annual and perennial weeds in *E. globulus*, *E. regnans* and *E. nitens* plantations.

NET 1, 5, 20 KG

MACSPRED PTY LTD
A.C.N. 011 029 495
Lot 22 Kennedy Drive
BALLARAT VIC 3350
TEL: 03 53358522

™ Macspred Trademark

GENERAL INSTRUCTIONS

- This product is a granular formulation that does not need to be mixed in water. It is an effective residual herbicide.
- For best residual control, apply during periods of active plant growth. Application when vegetation is dormant, semi-dormant or under moisture stress may not be as effective.
- In temperate winter rainfall areas where temperatures are generally below 25°C during the period of active growth, overall effect may be slow to appear. For best control of perennials, apply during the months of spring in these areas. In WA and SA apply only during April-August when adequate rainfall is expected.
- Sufficient moisture is required after treatment to promote root absorption by the weed species.
- Results obtained may vary with rainfall, soil types and other conditions.
- DO NOT apply Eucmix™ GR Granular Herbicide before (within 48 hours) or during periods of intense rainfall, when wind speed exceeds 30 kph (for ground application) or 20 kph (for aerial application) or under conditions which will cause granule movement to non-target areas.
- DO NOT broadcast apply to uncultivated or non-mounded or bare ground on slopes exceeding 30%.
- DO NOT apply within 10m of dirt tracks or roads.

RESISTANT WEEDS WARNING

Macspred Eucmix™ GR Granular Herbicide is a member of the uracils and sulfonylurea group of herbicides. Eucmix™ has the inhibitors of acetolactate synthase and inhibitors of Photosynthesis at Photosystem II mode of action. For weed resistance management Eucmix™ is a Group B C herbicide. Some naturally occurring weed biotypes resistant to Eucmix™ and other inhibitors of acetolactate synthase and inhibitors of Photosynthesis at Photosystem II herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Eucmix™ or other inhibitors of acetolactate synthase or inhibitors of Photosynthesis at Photosystem II.

Since the occurrence of resistant weeds is difficult to detect prior to use, Macspred Pty Ltd accepts no responsibility for any losses that may result from the failure of Eucmix™ to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

- Do not apply the product or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend. As guidance, this is twice their height or more depending on the species and other conditions.
- Do not apply the product where a surface flow of water could wash the chemical into areas where the roots of desirable trees or plants may extend.
- DO NOT apply to foliage or use on eroded areas, where sub-soil or tree roots are exposed, as injury to trees may result.
- DO NOT apply under weather conditions, or from equipment, that may cause granules to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

- DO NOT use in irrigation channels or in drains including roadside ditches and table drains.
- DO NOT empty equipment in the above situations.

- DO NOT use on leached, sandy soils, low in organic matter.
- Dangerous to aquatic life. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.
- DO NOT apply to water-logged or saturated soil.

PROTECTION OF LIVESTOCK

- Keep stock away from treated areas.

STORAGE AND DISPOSAL AND PROTECTION OF OTHERS

- During storage keep from contact with fertilisers, other pesticides and seeds.
- DO NOT re-use container.
- Store in the closed original container in a well ventilated area, as cool as possible but out of direct sunlight.
- Shake empty in application equipment. Destroy empty bags by puncturing or shredding them. Dispose of empty bags at a local authority landfill readily available in your area, or bury the bags at a depth of 500mm or more in a licensed/approved disposal site. In some States, wastes can only be buried at a licensed landfill. Do not burn empty containers or product.

SAFETY DIRECTIONS

Will damage eyes, nose, throat and skin. Avoid contact with eyes and skin. Do not inhale dust. When handling granules, wear elbow-length PVC gloves, goggles and a disposable dust mask. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves and goggles.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (TEL: 131126). For further information please refer to the Eucmix™ GR Material Safety Data Sheet.

NOTICE TO BUYER

To the extent permitted by law, all conditions and warranties and statutory or other rights or action which buyer or any other user may have against Macspred or Seller are hereby excluded. Macspred hereby gives Notice to Buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Macspred or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. Macspred's liability shall, in all circumstances, be limited to replacement of the product or a refund of the purchase price paid therefor.

NRA Approval Number: 49420/0401

Batch No:

Date of Manufacture:

Pat. Applied For

DIRECTIONS FOR USE:

Restrains - DO NOT use on leached sandy soils, low in organic matter.

METHOD OF APPLICATION - GROUND AND AIR (FIXED WING AND HELICOPTER)

CROP AND SITUATION	WEEDS CONTROLLED	STATES	RATE kg/ha	CRITICAL COMMENTS
<p><i>Eucalyptus globulus</i>, <i>E. regnans</i>, <i>E. nitens</i> plantations</p> <p><u>Post Planting</u> Tree Release Application (a) Pasture Sites (b) Native Woodland, Heath and Second Rotation Sites - Long term control</p>	<p>ANNUALS Capeweed (<i>Arctotheca calendula</i>), Annual Ryegrass (<i>Lolium rigidum</i>), White Clover (<i>Trifolium repens</i>)</p> <p>HARD TO KILL PERENNIALS Fog grass (<i>Holcus lanatus</i>)</p>	<p>ALL STATES</p>	<p>30</p>	<p>Allow 12 months after transplanting before application. Do not apply to weeds larger than 10cm in height or diameter.</p>

DIRECTIONS FOR USE

METHOD OF APPLICATION - WEED-A-METRE

CROP AND SITUATION	WEEDS CONTROLLED	STATES	RATE	CRITICAL COMMENTS
<p><i>Eucalyptus globulus</i>, <i>E. regnans</i>, <i>E. nitens</i> plantations</p> <p><u>Post Planting</u> Tree Release Application</p> <p>(a) Pasture Sites (b) Native Woodland, Heath and Second Rotation Sites - Long term control</p>	<p>ANNUALS Capeweed (<i>Arctotheca calendula</i>), Annual ryegrass (<i>Lolium rigidum</i>), White clover (<i>Trifolium repens</i>)</p> <p>HARD TO KILL PERENNIALS Fog grass (<i>Holcus lanatus</i>)</p>	ALL STATES	3 g/m ²	<p>Allow 12 MONTHS after transplanting before application.</p> <p>Use 3g trigger to apply 3 g/m²</p> <p>Do not apply to weeds larger than 10cm in height or diameter.</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.