



POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



TRIADIMENOL 250 EC

Fungicide

ACTIVE CONSTITUENT: 250 g/L TRIADIMENOL
Solvent: 600 g/L N-methyl-2-pyrrolidone

GROUP	C	FUNGICIDE
--------------	----------	------------------

Cabbage; Cauliflower & Broccoli:

Systemic control of Ring Spot

Cucurbits; Grapevines; Papaws

For control of powdery mildew

Sugar Cane

Control of Pineapple disease

20 Litre

*This information is adapted from the Triadimenol 250 EC Fungicide label.
Please consult the label on the pack before use.*

DO NOT DESTROY CONTAINER WHILE PRODUCT STILL REMAINS

4 FARMERS PTY. LTD.
A.C.N. 067 443 485
643 MURRAY STREET,
WEST PERTH, WESTERN AUSTRALIA 6005
Ph (08) 9356 3445



GENERAL INSTRUCTIONS

- 4 Farmers TRIADIMENOL 250 EC Fungicide is a broad spectrum, systemic, foliar fungicide. As it is quickly taken up by plants and rapidly translocated with systemic activity, 4 Farmers TRIADIMENOL 250 EC Fungicide is less affected by rainfall or irrigation than contact type fungicides.

FUNGICIDE RESISTANCE WARNING

GROUP	C	FUNGICIDE
--------------	----------	------------------

Triadimenol is a member of the D.M.I group of fungicides. For fungicide resistance management, 4Farmers Triadimenol 250 is a **Group C** fungicide.

Some naturally occurring individual fungi resistant to 4Farmers Triadimenol 250 and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by 4Farmers Triadimenol 250 and other **Group C** fungicides, thus resulting in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, 4Farmers Pty Ltd accepts no liability for any losses that may result from the failure of 4Farmers Triadimenol 250 to control resistant fungi.

MIXING

- Half fill the spray tank with clean water, then add the required quantity of 4 Farmers TRIADIMENOL 250 EC Fungicide with the agitator going. Fill to top with water and agitate thoroughly.
- DO NOT tank mix with trace elements or insecticides without first checking compatibility.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

- Harmful to fish.
- DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

- Keep out of reach of children.
- Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to tank. Do not dispose of chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.
- If not recycling, puncture top, bottom and sides of empty container then crush and bury in an approved local authority landfill. If no landfill available, bury under at least 500 mm of soil in a disposal pit set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers should not be burnt.

SAFETY DIRECTIONS:

Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing the spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow-length butyl rubber gloves and goggles. If product in eyes, washes it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.



FIRST AID:

- If poisoning occurs, contact a doctor or Poisons Information Centre on 131126 (24 hrs).

MSDS:

For further information see Material Safety Data Sheet which can be obtained from the supplier.

CONDITIONS OF SALE:

The use of 4 Farmers TRIADIMENOL 250 EC Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by 4Farmers Pty. Ltd. regarding its suitability, fitness or efficacy for any purpose for which it is used by the buyer, whether in accordance with the directions or not and 4 Farmers Pty. Ltd. accepts no responsibility for any consequences whatsoever resulting from the use of this product.

NRA APPROVAL NO.: 52067/1100

Batch No.:

Date of Manufacture:



DIRECTIONS FOR USE

Crop	Disease/ Pest	State(s)	Rate	Critical Comments
Cabbage Cauliflower Broccoli	Ring Spot	ALL STATES	200 or 400 mL/ha Plus Agridex [®] 200 mL/ha	Apply every 10 days. Start the program immediately following the first two consecutive days after transplanting. Use the higher rate under heavy disease conditions favour development or the disease.
Cucurbits	Powdery mildew		40 mL/100 L or 400 mL/ha	Apply a non-DMI fungicide such as Morestan [®] on a schedule commencing as soon as possible after emergence until fruit set. Then either tank mix or alternate Triadimenol with a non-DMI fungicide such as Morestan [®] on a 5 - 10 day schedule. Use Short Spray intervals when disease pressure is high or plants are growing rapidly. Good coverage is essential for good results.
Grapevines			10 mL/100 L Or 100 mL/ha	Apply thoroughly in the following spray program: 1. When shoots are 10cm long. 2. Pre flowering. 3. At flowering. 4. After fruit set. 5. Before bunches close. Do not apply in volumes less than 250 L/ha. For spray volumes between 250 and 1000 L/ha, use Triadimenol at 100 mL/ha. For volumes greater than 1000 L/ha, use the volume rate of 10 mL/100L.
Papaws			QLD NSW WA Only	20 mL/100 L Or 200 mL/ha
Sugar cane	Pineapple disease	QLD NSW ONLY	20 mL/100 L	Apply to setts by dipping or <i>spraying</i> . Ensure thorough wetting of cut ends.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

Cucurbits:	DO NOT HARVEST FOR 1 DAY AFTER APPLICATION
Cabbage, Cauliflower, Broccoli, Grapevines, Papaws	DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION
Cabbage, Cauliflower, Broccoli:	DO NOT GRAZE ANY TREATED AREA, OR CUT FOR STOCK FOOD
Sugar cane:	NOT REQUIRED WHEN USED AS DIRECTED