

KENSO
agcare

Ken-gran

750 WG

Selective Herbicide

ACTIVE CONSTITUENT: 750 g/kg TRIASULFURON

GROUP B HERBICIDE

For pre-plant control of Annual Ryegrass, Paradoxa Grass and certain broadleaf weeds in Wheat, and for post-emergent control of Wild Radish in Wheat, Oats and Barley as per Directions for Use table.

CONTENTS: 3.5 kg

IMPORTANT : READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT



KENSO
agcare

Kenso Corporation (M) Sdn Bhd 15a/243 Bradman Street, Acacia Ridge 4110 Ph: (07) 3711 5900

NRA Approval No: 56839/0203

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT store near food, feedstuffs, fertilizers or seed. Keep out of reach of children.

Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Ph.: 131126).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

WARRANTY AND EXCLUSION OF LIABILITY

Kenso Corporation (M) Sdn Bhd has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Kenso Corporation (M) Sdn Bhd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

In a Transport Emergency Dial 000 Police or Fire Brigade
--

NRA Approval No. : 56839/0203

Batch No. :

Date of Manufacture :



Ken-Gran 750 WG

Selective Herbicide

Active Constituent : 750 g/kg TRIASULFURON

GROUP	B	HERBICIDE
--------------	----------	------------------

For pre-plant control of Annual Ryegrass, Paradoxa Grass and certain broadleaf weeds in Wheat, and for post-emergent control of Wild Radish in Wheat, Oats and Barley as per Directions for Use table.

READ THIS LEAFLET BEFORE USE

NRA Approval No : 56839/0203



Kenso Corporation (M) Sdn Bhd
15a/243, Bradman Street, Acacia Ridge, 4110
Phone 07 3711 5900

**DIRECTIONS FOR USE
WESTERN AUSTRALIA ONLY**

Pre-emergent Application

Restraints: DO NOT apply to crops undersown with legumes.
DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season.

Crop	Weeds Controlled	Rate/ha	Critical Comments
Wheat	Burr Medic, Common Cotula, Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Denseflower Fumitory, Hedge Mustard, Indian Hedge Mustard, Matricaria, Paterson's Curse, Rough Poppy, Smallflower Fumitory, Wards Weed, Wild Turnip, Yellow Burrweed (Amsinckia) and suppression of Crassula	30 g	Apply to bare moist soil prior to sowing or at sowing and incorporate by the sowing operation using low profile 10cm combine points. Application should not be made to ridged or excessively cloddy soil. For best results apply to moist soil when follow up rain is likely to occur within 7 to 10 days. In conservation tillage situations where weeds and grasses have emerged apply as a tank mixture with the recommended rate of knockdown herbicide prior to sowing.
	Annual Ryegrass, Capeweed, Doublegee or Three-cornered Jack (Spiny Emex), Wireweed, suppression of Wild Radish, 60 to 80% suppression of Soursob	35 g	When used on Kulin Wheat on very acid soils (pH 5.5 in water) or under poor fertility conditions increased stem breakage may occur. Early season crop retardation may occur where the product is used on soils with a pH greater than 8, and which are prone to zinc deficiency.
	Annual Ryegrass, Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Hedge Mustard, Indian Hedge Mustard, Prickly Lettuce (Whipthistle), Smallflower Fumitory, Wild Turnip, Wireweed (Hogweed), Ward's Weed	10 g to 15 g plus 1 L of a 400 g/L trifluralin	This mixture to be only used on alkaline soils where pH is greater than 8 (1:5 Soil : Water suspension method). For best results apply mix to bare moist soil that has a minimum of trash and incorporate to a depth of 5cm just prior to sowing. Incorporation should be made within 4 hours of application. Heavy rainfall (greater than 50 mm) within 7 days of application may affect efficacy especially at the 10 g/ha rate. Late germination of some weeds eg Annual Ryegrass will not be controlled in seasons of above average rainfall. Use the higher rate where heavy density of weeds is expected.

WESTERN AUSTRALIA ONLY**Post-emergent Application**

Restrictions: DO NOT apply to crops undersown with legumes.
 DO NOT spray when very dry conditions prevail.
 DO NOT spray under dry frosty conditions
 DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season.
 If RAIN FALLS within 6 hours of application, the effect could be diminished.

Crop	Weeds Controlled	Rate/ha	Critical Comments
Barley, Oats, Wheat	Wild Radish	10 g to 15 g plus crop oil at 1 L per 100 L of spray mixture or surfactant at recommended label rates	Early Post-Emergent Application: Spray prior to the crop reaching mid-tillering (Zadoks - 23) and when Wild Radish is in the 2 to 6 leaf stage. Use rates towards the lower end of the range when weeds are small and growing conditions ideal. Spray only when weeds are actively growing. Late Post-Emergent Application: Spray during early flowering. DO NOT apply to the crop during crop anthesis or flowering (Zadoks 60-69). DO NOT spray on weeds under stress. WARNING: Application to Oats may cause some yellowing.
Barley, Oats, Triticale, Wheat From 3 leaf to early tillering stage	Turnip Weed	6.5 g + 300 mL Terbutryn 500 SC	Spray when weeds are in the 2 to 6 leaf stage (up to 6 node/leaflet for Field Peas, 10 leaflet for Vetch), except 2 to 4 leaf for Doublegee. Use rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist. Spray only when weeds are actively growing. Spray only after good rain and when top soil is moist. Best results are obtained when good soil moisture has been present since planting.
	Denseflower Fumitory, Hedge Mustard, Smallflower Fumitory	6.5 g + 300 mL Terbutryn 500 SC to 10 g + 440 mL Terbutryn 500 SC	
	Ball Mustard, Field Pea (volunteer)	10 g + 440 mL Terbutryn 500 SC	
	Deadnettle	10 g + 440 mL Terbutryn 500 SC to 13 g + 600 mL Terbutryn 500 SC	
	Australian Crassula, Doublegee or Three-cornered Jack (Spiny Emex), Volunteer Lupins	13 g + 600 mL Terbutryn 500 SC	

DIRECTIONS FOR USE SOUTH AUSTRALIA ONLY

Pre-emergent Application

Restraints: DO NOT apply to crops undersown with legumes
DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season.

Crop	Weeds Controlled	Rate/ha	Critical Comments
Wheat	Ball Mustard, Burr Medic, Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Denseflower Fumitory, Hedge Mustard, Indian Hedge Mustard, Prickly Lettuce, Rough Poppy, Smallflower Fumitory, Stemless Thistle, Vetch, Wards Weed, Wild Turnip, Yellow Burrweed (Amsinckia).	30 g	Apply to bare moist soil prior to sowing or at sowing and incorporate by the sowing operation using low profile 10cm combine points. Application should not be made to ridged or excessively cloddy soil. For best results apply to moist soil when follow up rain is likely to occur within 7 to 10 days. In conservation tillage situations where weeds and grasses have emerged apply as a tank mixture with the recommended rate of knockdown herbicide prior to sowing.
	Annual Ryegrass, Capeweed, Three-cornered Jack or Doublegee (Spiny Emex), Volunteer Chickpeas, Faba Beans and Field Peas, Wireweed, suppression of Skeleton Weed, Wild Radish, 60 to 80% control of Soursob and suppression of remaining plants.	35 g	KEN-GRAN 750 WG will provide good control of volunteer grain Legumes, however a small proportion of plants may survive and require an overspray to eliminate the potential for grain contamination. For Skeleton Weed a significant degree of control will be achieved on soil types of a predominantly sandy clay loam mixture with a pH greater than 8. Best control is observed where Skeleton Weed germinates in the very early stages of the crop. Surviving plants will be stunted.
	Annual Ryegrass, Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Hedge Mustard, Indian Hedge Mustard, Prickly Lettuce (Whipthistle), Smallflower Fumitory, Wild Turnip, Wireweed (Hogweed), Ward's Weed	10 g to 15 g plus 1 L of a 400 g/L trifluralin	This mixture to be only used on alkaline soils where pH is greater than 8 (1:5 Soil : Water suspension method). For best results apply mix to bare moist soil that has a minimum of trash and Incorporate to a depth of 5 cm just prior to sowing. Incorporation should be made within 4 hours of application. Heavy rainfall (greater than 50 mm) within 7 days of application may affect efficacy especially at the 10 g/ha rate. Late germination of some weeds eg Annual Ryegrass will not be controlled in seasons of above average rainfall. Use the higher rate where heavy density of weeds is expected.

SOUTH AUSTRALIA ONLY**Post-emergent Application**

Restraints: DO NOT apply to crops undersown with legumes.
 DO NOT spray when very dry conditions prevail.
 DO NOT spray under dry frosty conditions.
 DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season. If RAIN FALLS within 6 hours of application, the effect could be diminished.

Crop	Weeds Controlled	Rate/ha	Critical Comments
Barley, Oats, Wheat Late Post emergent application	Wild Radish	15 g plus crop oil at 1 L per 100 L of spray mixture or surfactant at recommended label rates	Late Post-Emergent Application: Spray during early flowering. DO NOT apply to the crop during crop anthesis or flowering (Zadoks 60-69). DO NOT spray on weeds under stress. WARNING: Application to oats may cause some yellowing.
Barley, Oats, Triticale, Wheat From 3 leaf to early tillering stage	Faba Beans (volunteer), Long Fruited Turnip, Long Headed Poppy, Paterson's Curse, Turnip Weed	6.5 g + 300 mL Terbutryn 500 SC	Spray when weeds are in the 2 to 6 leaf stage (up to 6 node/leaflet for Field Peas, 10 leaflet for Vetch), except 2 to 4 leaf for Doublegee. Use rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist. Spray only when weeds are actively growing. Spray only after good rain and when top soil is moist. Best results are obtained when good soil moisture has been present since planting.
	Hedge Mustard, Indian Hedge Mustard, Wild Turnip	6.5 g + 300 mL Terbutryn 500 SC to 10 g + 440 mL Terbutryn 500 SC	
	Ball Mustard, Crassula, Medic, Prickly Lettuce (Whipthistle)	10 g + 440 mL Terbutryn 500 SC	
	Field Pea (volunteer), Wild Radish	10 g + 440 mL Terbutryn 500 SC to 13 g + 600 mL Terbutryn 500 SC	
	Lupins (volunteer), Three-cornered Jack or Doublegee (Spiny Emex), Vetch, Wireweed (less than 3 leaves) suppression	13 g + 600 mL Terbutryn 500 SC	

DIRECTIONS FOR USE NEW SOUTH WALES ONLY

Pre-emergent Application

Restrictions: DO NOT apply to crops undersown with legumes.
DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season.

Crop	Weeds Controlled	Rate/ha	Critical Comments
Wheat	Black Bindweed (Climbing Buckwheat), Burr Medic, Common Cotula, Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Densflower Fumitory, Hedge Mustard, Indian Hedge Mustard, Lesser Swinecress, Paterson's Curse, Prickly Lettuce, Rough Poppy, Shepherd's Purse, Smallflower Fumitory, Sow Thistle, Stagger Weed, Turnip Weed, Wild Turnip, Yellow Burrweed (Amsinckia) and suppression of Variegated Thistle.	30 g	Apply to bare moist soil prior to sowing and incorporate by the sowing operation using low profile 10 cm combine points. Application should not be made to ridged or excessively cloddy soil. For best results apply to moist soil when follow up rain is likely to occur within 7 to 10 days. In conservation tillage situations where weeds and grasses have emerged apply as a tank mixture with the recommended rate of knockdown herbicide prior to sowing.
	Annual Ryegrass, Capeweed, Doublegee or Three-cornered Jack (Spiny Emex), Paradoxa Grass (Annual Phalaris), Wireweed, and suppression of Wild Radish and Mexican Poppy.	35 g	For best results for Paradoxa Grass control, apply to dry soil before the sowing rain.
	Annual Ryegrass, Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Hedge Mustard, Indian Hedge Mustard, Prickly Lettuce (Whipthistle), Smallflower Fumitory, Wild Turnip, Wireweed (Hogweed), Ward's Weed	(Southern NSW only) 10 g to 15 g plus 1 L of a 400 g/L trifluralin	This mixture to be only used on alkaline soils where pH is greater than 8 (1:5 Soil : Water suspension method). For best results apply mix to bare moist soil that has a minimum of trash and incorporate to a depth of 5 cm just prior to sowing. Incorporation should be made within 4 hours of application. Heavy rainfall (greater than 50 mm) within 7 days of application may affect efficacy especially at the 10 g/ha rate. Late germination of some weeds eg Annual Ryegrass will not be controlled in seasons of above average rainfall. Use the higher rate where heavy density of weeds is expected.

NEW SOUTH WALES ONLY**Post-emergent Application**

Restraints: DO NOT apply to crops undersown with legumes
 DO NOT spray when very dry conditions prevail.
 DO NOT spray under dry frosty conditions.
 DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season.
 If RAIN FALLS within 6 hours of application, the effect could be diminished.

Crop	Weeds Controlled	Rate/ha	Critical Comments
Barley, Oats, Wheat Late Post-emergent application	Wild Radish	15 g plus crop oil at 1 L per 100 L of spray mixture or surfactant at recommended label rates	Late Post-emergent Application: Spray during early flowering. DO NOT apply to the crop during crop anthesis or flowering (Zadoks 60-69) DO NOT spray on weeds under stress. WARNING: Application to oats may cause some yellowing.
Barley, Oats, Triticale, Wheat From 3 leaf to early tillering stage	Turnip Weed	6.5 g + 300 mL Terbutryn 500 SC	Spray when weeds are in the 2 to 6 leaf stage (up to 6 node/leaflet for Field Peas, 10 leaflet for Vetch), except 2 to 4 leaf for Doublegee. Use rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist. Spray only when weeds are actively growing. Spray only after good rain and when top soil is moist. Best results are obtained when good soil moisture has been present since planting.
	Hedge Mustard, Indian Hedge Mustard, London Rocket, Wild Turnip	6.5 g + 300 mL Terbutryn 500 SC to 10 g + 440 mL Terbutryn 500 SC	
	Black Bindweed (Climbing Buckwheat), Faba Bean (volunteer), Field Pea (volunteer), Medic, Shepherds Purse	10 g + 440 mL Terbutryn 500 SC	
	Deadnettle, Mexican Poppy, Wild Radish	10 g + 440 mL Terbutryn 500 SC to 13 g + 600 mL Terbutryn 500 SC	
	Coreopsis, Sunflower (volunteer), Vetch	13 g + 600 mL Terbutryn 500 SC	

DIRECTIONS FOR USE QUEENSLAND ONLY

Pre-emergent Application

Restrains: DO NOT apply to crops undersown with legumes.
DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season.

Crop	Weeds Controlled	Rate/ha	Critical Comments
Wheat	African Turnip Weed, Black Bindweed (Climbing Buckwheat), Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Hedge Mustard, Indian Hedge Mustard, New Zealand Spinach, Prickly Lettuce, Slender Celery, Smallflower Fumitory, Turnip Weed, Yellow Burrweed (Amsinckia)	30 g	Apply to weed free soil prior to sowing or at sowing and incorporate by the sowing operation using low profile 10cm combine points. Application should not be made to ridged or excessively cloddy soil. For best results apply when follow up rain is likely to occur within 7 to 10 days. In conservation tillage situations where weeds and grasses have emerged apply as a tank mixture with the recommended rate of knockdown herbicide prior to sowing.
	Annual Ryegrass, Common Peppergrass, Doublegee or Three-cornered Jack (Spiny Emex), London Rocket, Paradoxa Grass, Stagger Weed and suppression of Wild Radish and Wireweed.	35 g	For Paradoxa Grass control, apply to dry soil before the sowing rain. TO AVOID UNACCEPTABLE RESIDUES DO NOT graze or cut for stock food for 7 weeks after application.

QUEENSLAND ONLY**Post-emergent Application**

Restrains: DO NOT apply to crops undersown with legumes.
 DO NOT spray when very dry conditions prevail.
 DO NOT spray under dry frosty conditions.
 DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season. If RAIN FALLS within 6 hours of application, the effect could be diminished.

Crop	Weeds Controlled	Rate/ha	Critical Comments
Barley. Wheat From 3 leaf to early tillering stage	Turnip Weed	6.5 g + 300 mL Terbutryn 500 SC	Spray when weeds are in the 2 to 6 leaf stage, except 2 to 4 leaf for Doublegee. Use rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist. Spray only when weeds are actively growing. Best results are obtained when good soil moisture has been present since planting. TO AVOID UNACCEPTABLE RESIDUES DO NOT graze or cut for stock food for 14 days after application.
	Denseflower Fumitory, Indian Hedge Mustard, Smallflower Fumitory, Wild Turnip	6.5 g + 300 mL Terbutryn 500 SC to 10 g + 440 mL Terbutryn 500 SC	
	Black Bindweed (Climbing Buckwheat), London Rocket	10 g + 440 mL Terbutryn 500 SC	
	Wild Radish	10 g + 440mL Terbutryn 500 SC to 13 g + 600 mL Terbutryn 500 SC	
	Coreopsis, Corn Gromwell (Sheepweed or White Ironweed)	13 g + 600 mL Terbutryn 500 SC	

DIRECTIONS FOR USE VICTORIA ONLY

Pre-emergent Application

Restrains: DO NOT apply to crops undersown with legumes.
DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season

Crop	Weeds Controlled	Rate/ha	Critical Comments
Wheat	Annual Ryegrass, Burr Medic, Capeweed, Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Hedge Mustard, Indian Hedge Mustard, Prickly Lettuce (Whipthistle), Smallflower Fumitory, Volunteer Chickpeas, Faba Beans and Field Peas, Wild Turnip, Wireweed (Hogweed), Yellow Burrweed (Amsinckia), and suppression of Wild Radish and Skeleton Weed.	30 to 35 g	<p>Apply to bare moist soil prior to sowing or at sowing and incorporate by the sowing operation using low profile 10 cm combine points.</p> <p>Application should not be made to ridged or excessively cloddy soil. Use rates towards the lower end of the range where broadleaf weeds are the major problem.</p> <p>Use the higher rate where Capeweed, Volunteer Chickpeas, Faba Beans and Field Peas, Wild Radish and Wireweed are the problem KEN-GRAN 750 WG will provide good control of Volunteer Grain Legumes, however a small proportion of plants may survive and require an overspray to eliminate the potential for grain contamination.</p> <p>For best results apply to moist soil when follow up rain is likely to occur within 7 to 10 days.</p> <p>Use the lower rate (30 g/ha) on sandy clay loams with a pH greater than 8.5.</p> <p>For Skeleton Weed a significant degree of control will be achieved on soil types of a predominantly sandy clay loam mixture with a pH greater than 8. Best control is observed where Skeleton Weed germinates in the very early stages of the crop.</p> <p>Surviving plants will be stunted.</p>
	Annual Ryegrass, Corn Gromwell (White Ironweed, Sheepweed), Deadnettle, Hedge Mustard, Indian Hedge Mustard, Prickly Lettuce (Whipthistle), Smallflower Fumitory, Wild Turnip, Wireweed (Hogweed), Ward's Weed	10 g to 15 g plus 1 L of a 400 g/L trifluralin	This mixture to be only used on alkaline soils where pH is greater than 8 (1:5 Soil : Water suspension method). For best results apply mix to bare moist soil that has a minimum of trash and incorporate to a depth of 5 cm just prior to sowing. Incorporation should be made within 4 hours of application. Heavy rainfall (greater than 50 mm) within 7 days of application may affect efficacy especially at the 10 g/ha rate. Late germination of some weeds eg Annual Ryegrass will not be controlled in seasons of above average rainfall. Use the higher rate where heavy density of weeds is expected.

VICTORIA ONLY

Post-emergent Application

Restraints: DO NOT apply to crops undersown with legumes.
 DO NOT spray when very dry conditions prevail.
 DO NOT spray under dry frosty conditions.
 DO NOT use if another Group B herbicide (ALS inhibitor) has been used during the current season. If RAIN FALLS within 6 hours of application, the effect could be diminished.

Crop	Weeds Controlled	Rate/ha	Critical Comments
Barley, Oats, Wheat Late Post-emergent application	Wild Radish	15 g plus crop oil at 1 L per 100 L of spray mixture or surfactant at recommended label rates.	Late Post-emergent Application: Spray during early flowering. DO NOT apply to the crop during crop anthesis or flowering (Zadoks 60-69). DO NOT spray on weeds under stress. WARNING: Application to Oats may cause some yellowing.
Barley, Oats, Triticale, Wheat From 3 leaf to early tillering stage	Paterson's Curse, Turnip Weed	6.5 g + 300 mL Terbutryn 500 SC	Spray when weeds are in the 2 to 6 leaf stage (up to 6 node/leaflet for Field Peas, 10 leaflet for Vetch), except 2 to 4 leaf for Doublegee. Use rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist. Spray only when weeds are actively growing. Spray only after good rain and when top soil is moist. Best results are obtained when good soil moisture has been present since planting.
	Hedge Mustard, Indian Hedge Mustard, Wild Turnip	6.5 g + 300 mL Terbutryn 500 SC to 10 g + 440 mL Terbutryn 500 SC	
	Crassula, Faba Beans (volunteer), Hyssop, Loosestrife, Medic, Prickly Lettuce (Whipthistle)	10 g + 440 mL Terbutryn 500 SC	
	Deadnettle, Field Pea (volunteer), Wild Radish	10 g + 440 mL Terbutryn 500 SC to 13 g + 600 mL Terbutryn 500 SC	
	Lupins (volunteer), Vetch, Wireweed (less than 3 leaves) suppression	13 g + 600 mL Terbutryn 500 SC	

**ALL STATES
 NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
 UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

WITHHOLDING PERIOD
PRE-EMERGENT APPLICATION : DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 WEEKS AFTER APPLICATION.
POST-EMERGENT APPLICATION : DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS
 KEN-GRAN 750 WG Selective Herbicide is a water dispersible granular herbicide for the pre-plant, incorporated by sowing, control of annual ryegrass, paradoxa grass and certain broadleaf weeds in wheat and for post-emergent control of wild radish in wheat, oats and barley.
 Crops other than wheat, barley, oats, triticale and cereal rye can be very sensitive to low soil concentrations of KEN-GRAN 750 WG Selective Herbicide thus prior to using the product, careful consideration should be given to crop rotation plans.

Resistance Weeds Warning **GROUP B HERBICIDE**

KEN-GRAN 750 WG Selective Herbicide is a member of the sulfonylurea group of herbicides and has the ALS Inhibitor mode of action. For weed resistance management this is a Group B herbicide. Some naturally occurring weed biotypes resistant to KEN-GRAN 750 WG Selective Herbicide and other Group B herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by KEN-GRAN 750 WG Selective Herbicide or other Group B herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, accepts no liability for any losses that may result from the failure of KEN-GRAN 750 WG Selective Herbicide to control the resistant weeds.

Mixing

KEN-GRAN 750 WG Selective Herbicide is a water dispersible granular herbicide, which mixes readily with water and is applied as a spray.

- Partly fill the spray tank with water
- Start the agitation
- Add the correct amount of product to the spray tank with the agitation system running
- Continue agitation while topping up the tank with water and while spraying
- Use the spray mix within 24 hours of preparation

Application

Ground application: Apply by boom spray, applying 30 to 100 litres of water per hectare. Avoid overlapping of boom runs.

Aerial application: Always spray in a cross wind of less than 5 knots. Ensure good spray coverage is obtained. Apply 20 to 40 litres per hectare.

Sprayer Cleanup

Where the sprayer is being used to spray cereal crops, rinse the sprayer thoroughly with water. Where the sprayer is being used to spray crops other than cereals:

1. Drain tank and rinses tank and spray boom with clean water for at least 10 minutes.
2. Fill the tank with clean water and add to it 300 mL of household chlorine bleach (containing 4% chlorine) per 100 L of water. Rinse hoses and boom and leave in tank for 15 minutes whilst agitating. Drain through nozzle.
3. Repeat step 2 and then rinse thoroughly with clean water to remove all traces of chlorine bleach.
4. Nozzles and filters should be cleaned separately.
5. Dispose of all water used for cleaning.

Compatibility

When using a tank mix of KEN-GRAN 750 WG and Spray.Seed*, add KEN-GRAN 750 WG, with constant agitation to approximately half the total volume of water to be used.

Ensure that the KEN-GRAN 750 WG is fully dispersed.

Add the Spray.Seed*, fill the spray tank to full volume with water and mix thoroughly.

Apply tank mix immediately, under constant agitation.

KEN-GRAN 750 WG is compatible with Dual* Gold, Avadex* BW, Spray.Seed*, Ken-Up 450 CT, sodium molybdate, zinc sulphate, manganese sulphate, copper sulphate, 2,4-D Amine, Ken-Met 600, Kensban 500, Ambush*, bromoxynil, trifluralin.

CROP ROTATION GUIDELINES**(Pre-emergent Application)**

Where the product is applied at the rate of 30 to 35 g/ha: Unless otherwise specified (see table below), wheat, barley, oats, triticale and cereal rye can be planted the following season without restriction. For other specified crops the KEN-GRAN 750 WG treated area may be replanted after the interval indicated in the table below.

These recommendations are made on the assumption that KEN-GRAN 750 WG is applied to a wheat crop that reaches maturity in the season of application.

Soil pH (1:5 Soil:Water suspension method)	State	Replanting Interval	Minimum rainfall requirements between application and sowing the following crop	Crop	
6.5 or less	Qld, NSW, Vic, SA, WA only	12 months	300 mm	Field Peas, Linseed, Lucerne, Lupins, Medics ⁺ , Subterranean Clover ⁺ , Faba Beans, Chickpeas and Canola	
	Qld, NSW only	15 months	700 mm	Sorghum, Maize, Soybean, Cotton, Cowpea and Mung Bean	
		18 months	900 mm	Sunflowers	
6.6 to 7.5	Qld, NSW only	12 months	500 mm	Chickpeas and Canola	
		15 months	700 mm	Sorghum, Maize, Soybean, Cotton, Cowpea and Mung Bean	
		18 months	900 mm	Sunflowers	
	Vic, SA, WA only	22 months	500 mm	Field Peas, Linseed, Lucerne, Lupins, Medics ⁺ , Subterranean Clover ⁺ , Faba Beans, Chickpeas, Canola, Sorghum, Maize, Soybean and Cotton	
7.6 to 8.5	Vic, SA only	12 months	250 mm	Barley, Oats, Cereal Rye for grain crops	
			300 mm	Barley, Oats, Cereal Rye for hay crops	
	Qld, NSW only	12 months	500 mm	Chickpeas and Canola	
		18 months	700 mm	Sorghum, Maize, Soybean, Cotton, Cowpea and Mung Bean	
	Vic, SA, WA only	24 months	700 mm	Field Peas, Linseed, Lucerne, Lupins, Medics ⁺ , Subterranean Clover ⁺ , Faba Beans, Chickpeas, Canola, Sorghum, Maize, Soybean and Cotton	
	8.6 and above	Vic, SA only	12 months	250 mm	Barley, Oats, Cereal Rye for grain crops
				300 mm	Barley, Oats, Cereal Rye for hay crops
Qld, NSW, Vic, SA, WA, only		24 months	700 mm	Field Peas, Linseed, Lucerne, Lupins, Medics ⁺ , Subterranean Clover ⁺ , Faba Beans, Chickpeas, Canola, Sorghum, Maize, Soybean and Cotton	

⁺ Includes natural regeneration of Subterranean Clover and Medics. For all other crops seek advice from a Kenso Corporation (M) Sdn Bhd representative.

Where the product is applied at 10 to 15 g/ha plus 1 litre of 400 g/L trifluralin:

Where the pH is less than 7.5 the following crops can be replanted from 9 months after application of KEN-GRAN 750 WG providing 300 mm of rainfall has been recorded; field peas, canola, chickpeas, medics, clover, lucerne, safflower, lupins, cereal rye, barley, oats, wheat, triticale. Where rainfall is less than 300 mm for this period further advice should be sought from a Kenso Corporation (M) Sdn Bhd representative.

Where the pH is above 7.5 the following crops can be replanted 9 months after application of KEN-GRAN 750 WG - cereal rye, wheat, oats, barley, triticale.

Where the pH is above 7.5 the following crops can be replanted 12 months after application providing 350 mm of rainfall has been recorded; field peas, canola, chickpeas, medics, clover, lucerne, safflower, lupins. Where less than 350 mm of rain has fallen between application and sowing the crop, further advice should be sought from a Kenso Corporation (M) Sdn Bhd representative.

These recommendations are made on the assumption that KEN-GRAN 750 WG is applied to a wheat crop that reaches maturity in the season of application.

Where KEN-GRAN 750 WG is used on soil types with pH greater than 8 (1:5 soil:water suspension test), further advice should be sought from Kenso Agcare regarding crop rotation guidelines, except for barley, cereal rye, oats, triticale and wheat.

Crop Rotation Guidelines (Post-emergent Application)

Soil pH (1:5 Soil:Water suspension method)	State	Replanting Interval	Crop
6.5 or less	NSW, Vic, SA, WA only	7 months	Field Peas, Linseed, Lucerne, Lupins, Medics ⁺ , Subterranean Clover ⁺ , Faba Beans, Chickpeas and Canola
		14 months	Sorghum, Maize, Soybean, Cotton
6.6 to 8.0	NSW, Vic, SA, WA only	20 months	Chickpeas, Canola, Sorghum, Maize, Soybean, Cotton, Field Peas, Linseed, Lucerne, Medics ⁺ , Subterranean Clover ⁺ , Faba Beans

⁺ Includes natural regeneration of Subterranean Clover and Medics. For all other crops a replanting interval of 24 months has to be observed.

Crop Rotation Guidelines (tank mixture of KEN-GRAN 750 WG and Terbutryn)

The following crops can be planted after an application of KEN-GRAN 750 WG and Terbutryn:

In areas where pH is less than 7.5

- **9 months** after an application: barley, cereal rye, chickpeas, faba beans, field peas, lupins, medics, oats, canola, safflower, subclover, triticale, wheat In areas where pH is greater than 7.5

- **9 months** after an application: barley, cereal rye, oats, triticale, wheat.

- **14 months** after an application: cotton, maize, sorghum, soybeans, sunflowers. Where residual herbicides are applied following an application, the crop rotational guidelines for these products must be followed.

- **22 months** after an application: chickpeas, faba beans, field peas, lupins, medics, canola, safflower, subclover.

For all other crops advice should be sought from Kenso Agcare.

PRECAUTION

Some crop yellowing or crop retardation may occur where stress factors such as water logging, drought, excessive soil acidity or alkalinity, nutrient or trace element deficiency, disease - Rhizoctonia, Take-All, cereal cyst nematodes or soil insects are present or occur following application.

Special care should be taken with regard to the application of KEN-GRAN 750 WG to durum wheats as these may be more sensitive where the above stresses are present. In these situations crop recovery will be rapid provided the stress factors DO NOT continue exerting a negative effect on the crop's growth.

Crop retardation may also occur in some instances where considerable late summer/early autumn weed growth occurs. Weeds such as goosefoot, *Chenopodium* spp can release herbicidally active compounds into the soil.

Re-entry Period

DO NOT enter treated areas without protective clothing until spray has dried.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply on or near shrubs, trees, lawns or crops other than wheat, oats and barley.

DO NOT drain or flush equipment on, or near desirable trees or other plants, where their roots may extend or in situations where by movement of soil, or seepage, absorption of the herbicide may occur.

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift over nearby susceptible plants or crops, cropping lands or pastures.

DO NOT allow spray to drift onto adjacent crops and non-target desirable plants.

DO NOT allow spray to drift onto adjacent fallow land.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT store near food, feedstuffs, fertilizers or seed. Keep out of reach of children.

Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Ph.: 131126).


MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

WARRANTY AND EXCLUSION OF LIABILITY

Kenso Corporation (M) Sdn Bhd has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Kenso Corporation (M) Sdn Bhd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

* Registered trademarks



NRA Approval No. : 56839/0203



Kenso Corporation (M) Sdn Bhd
15a/243, Bradman Street, Acacia Ridge, 4110
Phone 07 3711 5900