



GENERAL INSTRUCTIONS

- For best results use high volume rates
- Spray to wet all foliage evenly. Apply when grasses are growing vigorously and before seed stems develop.
- Atlapon 2,2-DPA herbicide is slow in acting and it may take 6-8 weeks before full effect is seen.
- Atlapon 2,2-DPA in the soil may inhibit germination of sensitive plants for 30 days or more, depending on rate applied, soil type, temperature and amount of rainfall. Decomposition proceeds most rapidly in warm moist soils.

MIXING Atlapon 2,2-DPA dissolves readily in water and may be added directly to the spray tank or washed through a strainer.

WETTING AGENTS The addition of a Non-ionic wetting agent increases the efficiency of the spray.

RESISTANT WEEDS WARNING Agspray Atlapon 2,2-DPA Systemic Herbicide is a member of the Alkanoic acids group of herbicides. Atlapon has the inhibitors of fat synthesis mode of action. For weed resistance management, the product is a Group J herbicide. Some naturally occurring weed biotypes resistant to Atlapon and other inhibitors of fat synthesis herbicides may exist through normal; genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Agspray Atlapon or other inhibitors of fat synthesis herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Agspray Chemical Company Pty Ltd accepts no liability for any losses that may result from the failure of Agspray Atlapon to control resistant weeds.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT DO NOT contaminate streams, rivers or waterways with chemical or used container.

PROTECTION OF CROPS, NATIVE AND NON-TARGET PLANTS DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto nearby susceptible plants, adjacent crops, crop lands or pastures. DO NOT plant sensitive crops such as grass, small grains, corn or beans for at least 30 days after application.

DO NOT use or allow to drift onto pineapples or beneath fig trees or other sensitive crops.

DO NOT allow water movement from sprayed irrigation channels on to susceptible crops for 7 days.

DO NOT exceed rates specified.

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

AGSPRAY

ATLAPON 2,2-DPA

Systemic Herbicide

Active Constituent: 740g/kg 2,2-DPA present as the sodium salt

GROUP J HERBICIDE

A SYSTEMIC HERBICIDE FOR THE CONTROL OF ANNUAL AND PERENNIAL GRASSES INCLUDING PASPALUM, JOHNSON, COUCH, KIKUYU, CUMBUNGI, RUSHES AND SEDGES AS INDICATED IN THE DIRECTIONS FOR USE TABLE.

25 kg

AGSPRAY CHEMICAL COMPANY PTY. LTD. ACN 005 329 573
3/44 Fitzgerald Road, Laverton North, Vic. 3026 Ph: (03) 9369 8198

STORAGE AND DISPOSAL Store in the closed original container in a well-ventilated area, out of direct sunlight.

KEEP OUT OF REACH OF CHILDREN. Do not store for prolonged periods in direct sunlight. Shake bag contents into spray tank until bag is empty. Do not dispose of undiluted chemicals on site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers or product should not be burnt.

SAFETY DIRECTIONS Will irritate the eyes and skin. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with plenty of water. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID If poisoning occurs, contact a doctor or **Poisons Information Centre (131126).**

MATERIAL SAFETY DATA SHEET Additional information is listed on the Material Safety Data Sheet available from Agspray Chemical Company Pty. Ltd.

NOTICE This information is not intended to replace the product label. Failure to use any product in a manner consistent with the label may be illegal, prove dangerous and render the product ineffective. While every endeavour has been made to ensure that this information is accurate and up to date, Agspray Chemical Company Pty Ltd will not accept any responsibility for any use of this product based on the information herewith if this use is inconsistent with the label supplied at the time of purchase.

Batch No.:

Date of Manufacture:

APVMA Approval No.
31434/25/1203



DIRECTIONS FOR USE Note: Except where specified more effective control will be obtained using split application at half rates.

Situation and Crop	Weeds Controlled	State	Rate			Critical Comments
			General Rate/ha	Handgun kg/100L (1000L/ha)	10L Knapsack per 100m ²	
Non-Crop Areas	Annual Grasses	All States	5-10kg	500g-1kg	50-100g	Spray young seedlings and growing plants. Half rate 3-4 week intervals.
	Perennial Grasses		20-40kg	2-4kg	200-400g	
	Watsonia		10kg	1kg	100g	Apply at late pre flowering.
Irrigation Channels, Drains, Bore Drains	Rush and Sedge		10-20kg	1-2kg	100-200g	Where cumbungi or water couch is present, use higher rate in 2000L/ha. Cumbungi: Good results obtained if active regrowth is promoted by cutting and burning. Spray regrowth preferably when seeds are formed. Make sure channels and drains are dry before spraying or have no more 5-8cm of water. Overall rate should be not less than 40kg/ha. Spray application is preferable.
	Cumbungi, Water Couch					
Pastures	Serrated Tussock	Vic, NSW, WA, SA, Qld only	8kg	800g	80g	Apply between January and April and burn off 6-8 weeks later OR apply during January or February if burn off is not possible.
	Alternatively		28kg			
	Parramatta Grass	NSW only	5-10kg	1kg		Apply when plants are actively growing.
Lucerne	Annual Grasses	All States	3.5kg	300-500g	30-50g	Apply after cutting and grazing.
	Annual Rye Grass and Brown Top		5kg	500g	50g	DO NOT use more than 1kg/ha on seedling lucerne.
Orchards, Apples and Pears	Annual and Perennial Grasses		7.5-10kg	750g-1kg	75-100g	DO NOT use more 10kg/ha per annum. DO NOT apply to trees under 4 years old. Half rate at 4-6 week intervals.
Cherries, Peaches, Apricots	Annual and Perennial Grasses		6-8kg	600-800g	60-80g	DO NOT exceed 8kg/ha per annum. Apply to established trees only. Half rate at 4-6 week intervals.
Black and Red Currants	Annual and Perennial Grasses		10-15kg	1-1.5kg	100-150g	DO NOT use more than 15kg/ha per annum. DO NOT apply to bushes under 4 years old. Half rate at 4-6 week intervals.
Citrus (at least 4 years old)	Kikuyu, couch, Paspalum Dilatatum		2.5-4kg	500g	25-40g	Trees must be at least 4 years old. DO NOT exceed 12kg/ha per annum. Apply 500-800ml spray solution per 10m ² around each tree. Repeat 3 times at 10 day intervals.

Situation and Crop	Weeds Controlled	States	Rate			Critical Comments
			General Rate/ha	Handgun kg/100L (1000L/ha)	10L Knapsack per 100m ²	
Vines (at least 4 years old)	Kikuyu, couch, Johnson grass	All States	10kg	1kg	100g	Do not use more than 10kg/ha per annum. Apply direct spray into vine rows. Half rate at 10-14 day intervals.
Cotton, Potatoes, Tobacco, Sunflower, Maize, Soybean	Annual and Perennial Grasses	Qld only	2.5kg	200-500g	20-50g	Spray between rows.
		NSW, SA, WA, Tas, Vic only	5kg			
Bananas	Annual and Perennial Grasses	NSW and Qld only	10-15kg	1-1.5kg	100-150g	DO NOT exceed 30kg/ha per annum. DO NOT apply to plants especially suckers. Avoid spray drift in Lady Finger plantations. DO NOT spray while soil is wet or rain is likely. Half rate at 4-6 week intervals.
Tea-Trees (Melaleuca alternifolia)	Annual and Perennial Grasses	NSW only	2.5-10kg			Use higher rates for perennial weeds or where grasses are well established. Apply as an overall spray to seedling trees or immediately after harvest. At other times apply as a directed spray to the base of trees. Some burning of the leaf tips may occur at the 10kg/ha rate. A repeat application after 10 days may be necessary to control couch.
Sugar Cane	Annual and Perennial Grasses	NSW, Qld only	10kg	1kg	100g	Spray between rows only when canes 1.2-1.5 metres high. Repeat application may be necessary.
	Johnson Grass and Phragmites		10kg			Repeat application may be necessary.

NOT TO BE USED FOR ANY OTHER PURPOSE OR IN ANY OTHER MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

EDIBLE CROPS: Do not harvest for 7 days after application.

PASTURES & FORAGE CROPS: Do not graze pastures or forage crops or cut for stock food for 2 days after application.