



POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Baton[®] Low

Herbicide

ACTIVE CONSTITUENT: 800 g/kg 2,4-D present as the dimethylamine salt

GROUP I HERBICIDE

For the control of broadleaved weeds in fallow before direct drilling or sowing of cereals and pastures; and in cereal crops, pastures, sugar cane, peanuts and non agricultural areas as per the Directions for Use Table.

This is a PHENOXY HERBICIDE that can cause severe damage to susceptible crops such as cotton, grapes, tomatoes, oil seed crops and ornamentals.

READ THIS LEAFLET THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

© Baton is a registered trademark of Nufarm Australia Limited

nufarm.com.au

APVMA Approval No.:46314/55973

DIRECTIONS FOR USE

RESTRAINTS

DO NOT spray if rain seems likely within 6 hours or if strong winds prevail.

DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

1. FIELD CROPS

SITUATION & CROP	WEEDS	STATE	RATE	CRITICAL COMMENTS
Wheat	Refer Weed Table	NSW, SA only	250g-1.3kg/ha	Apply after the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller (NSW, SA only). Apply from tillering to boot stage (Vic only). Apply from mid tillering to before boot stage (Qld only). Apply from the 5 leaf to fully tillered (Tas only).
		Vic only	250g-1.1kg/ha	
		Qld only	400g-1.1kg/ha	
		Tas only	1.1kg/ha	
Barley		NSW, SA only	250g-1.1kg/ha	
		Vic only	250g-1.1kg/ha	
		Qld only	400g-1.1kg/ha	
		Tas only	1.1kg/ha	
Cereal rye, Triticale		NSW, SA only	250g-1.1kg/ha	
		Vic only	250g-1.1kg/ha	
Oats		NSW, SA only	250-600g/ha	
		Vic only	250-600g/ha	
Cereals: Wheat, Oats, Barley	Cape tulip	WA only	500g-1kg/ha	Apply from the 5 leaf stage up to jointing stage (Z15-33). Apply after the 6 leaf stage (Z16) for Cranbrook, Jjacup, Aroona and Spear wheat and Mortlock oats to avoid possible damage. DO NOT spray if lucerne is present. WEED STAGE: 10-15cm. Docks should be sprayed before 5 leaf stage. Cape tulip - low rate for cormils only.
	Dock, Saffron thistle		900g/ha	
	Indian hedge mustard, London rocket, Lupin, Rapistrum, Wild radish		600g/ha	
	Wild turnip		500g/ha	
	Capeweed, Doublegee, Erodium, London rocket, Lupin, Mustard, Rapistrum, Wild radish, Wild turnip		160g/ha plus 500mL/ha Nufarm Flowable Diuron (500g/L)	Apply when crop has 4-5 leaves and most weeds have germinated and are in 2-5 leaf stage. Crop and weeds should be dry at time of application. Some temporary yellowing of crop may occur after application. Undersown sub-clovers may be slightly retarded. DO NOT apply to undersown medics.

SITUATION & CROP	WEEDS	STATE	RATE	CRITICAL COMMENTS
Fallow, Stubble Spray prior to direct drilling or sowing a) Winter cereals b) Winter Cereals and peanuts	Refer Weed Table	Vic only	250g-1.1kg/ha	Observe plant back periods given in the table on this leaflet. Can be mixed with Glean*, Seige*, Nuquat® or SpraySeed* where grasses are present. For Skeleton weed, spraying should only be done 6-8 weeks before anticipated sowing date and subsequent cultivation limited to a minimum.
		NSW & Qld only	400g-1.3kg/ha	
Millet	Refer Weed Table	NSW, SA, Vic only	400-900g/ha	Spray when secondary roots have developed, when fully tillered and before heads start to form at the base of the tillers. DO NOT use on Panorama millet or Panicum.
		Qld only	400-700g/ha	
Saccaline, Broom Millet, Millet	Cape tulip, Dock, Saffron thistle, Indian hedge mustard, London rocket, Lupin, Rapistrum, Radish, Wild turnip	WA only	900g/ha	Spray when crop is 10-30cm high and secondary roots have developed and before tasselling. Apply as direct spray to weeds.
Sugar Cane	Bindy eye (Star burr), Blue top, Cobbler's pegs, Fleabanes, Jute, Leucas, Needle burr, Spear thistle, Water primrose, Ipomea vines, Convolvulus vines Chinese mint, Blue snakeweed	Qld only	1.4 -2.8kg/ha	Add 175-350mL Nufarm Activator®/100L of spray mixture. Agitate well. DO NOT use on Q63, Q67, Q80 or Q96 Varieties.
			2.8kg/ha	
Peanuts	Broadleaf weeds; except Noogoora burr. Grasses; except Mossman burr	Qld only	1.4 or 2.8kg/ha	LOWER RATE: Apply as BAND SPRAY as soon as possible after planting in a 55cm band. HIGHER RATE: Apply as OVERALL SPRAY after planting and before crop emergence. Some crop damage may occur if heavy rain falls between application and crop emergence.
Harvest Aid or Salvage Spray - Winter Cereals	Dessicate broadleaf weeds Refer to Weed Table	Qld only	900g-1.3kg/ha	Apply after dough stage.
		Qld & NSW only		

2. PASTURES, NON-AGRICULTURAL, RIGHTS OF WAY, INDUSTRIAL, LAWNS

SITUATION & CROP	WEEDS	STATE	RATE	CRITICAL COMMENTS
Pastures and Non-Agricultural	Refer Weed Table	NSW, Qld, SA, Tas only	400g-1.3kg/ha	Pasture legumes including lucerne, clovers and medics may be damaged unless well protected by grasses. Spot spraying is preferred.
	Amsinckia, Docks, Bindweed, Caltrop, Flatweed, Spear thistle, Capeweed, Doublegee, Saffron thistle, Mustard, Wild radish, Wild turnip, Annual thistles, Paterson's curse.	WA only	900g/ha	For pastures not containing legumes. Only seedling Docks, Spear thistle and Saffron thistle will be controlled.
	Afghan melons		1.3kg/ha plus 1% crop oil	Spray when plants are actively growing preferably before flowering or vining.
	Paddy melons		600-900g/ha	
	Prickly saltwort (Roly poly)		1.3kg/ha	Spray when plants are small.
	Stinkwort		1.3-2.5kg/ha plus surfactant	Best results are obtained when plants are small. Use high rate on larger plants.
	Dove weed		2.5kg/ha	Spray after good emergence of seedlings.
Pastures, Rights of Way and Industrial	Boxthorn, Boneseed, Hawthorn	Vic, SA only	500g/100L water 1kg in 1.6L water	Spot Spraying: For Boneseed only, thoroughly wet plants or seedlings. Apply to freshly cut stumps
	Groundsel	NSW, Qld, SA only	500g/10L water	MISTING: Lightly wet plants.
			250g/100L water	HIGH VOLUME: Thoroughly wet plants.
			125g/10L water	CUT STUMP: Swab the cut stump within one hour of cutting. Apply by a pouring can or knapsack spray.
			2.3-3.4kg/ha	AERIAL APPLICATION: Spray when Groundsel is actively growing.
	Lantana	NSW, Qld, SA only	250g/100L water	Use a coarse spray with sufficient pressure to penetrate canopy and wet stems as well as foliage. Spray at the end of a wet Summer (March to May). Defoliation should occur but respraying of new growth will be necessary in following Autumn. Broadcast grass seed and keep stock off following Summer to allow the pasture to establish. Damage may result to pasture legumes.
Mother of millions	NSW only	300g/100L water	Hand gun and Knapsack only. A thorough coverage of leaves and plantlets is necessary. Use Nufarm Chemwet 1000 at the rate of 1mL of surfactant/1L of mixture.	

SITUATION & CROP	WEEDS	STATE	RATE	CRITICAL COMMENTS
Pastures, Rights of Way and Industrial (cont)	Noogoora Burr, Weir Vine (Ipomea).	Qld only	250g/100L water	In all cases apply to young, actively growing weeds, ensuring thorough coverage. ^ Spray rosette stage. • Repeat spraying necessary.
	Annual and Perennial pigweed, Artichoke thistle, Bathurst burr, Billygoat weed, Blue snakeweed, Burr medic, Clockweed [^] , Fleabanes, Galvanised burr, Hemlock, Hoary cress*, Kyalinga weed (Whisker grass), Knobweed, Milky cotton bushes, Parthenium weed, Paterson's curse, Saffron thistle, Star burr, Thornapple, Variegated thistle [^]		250g/100L water	
	Rubber vine		125g/10L water	
Conservation Tillage - Direct Drilling, Surface Sowing or Fallow Maintenance	Charlock, Mustards, Shepherd's purse, Saffron, Slender, Spear and Variegated thistles, Turnip weed, Wild radish, Wild turnip	All States	400g-1.3kg/ha	Apply to actively growing young plants before sowing. Observe plant back periods given in the table on this leaflet.
	Clover, Sorrel		900g/ha plus 280-400g/ha Kamba® 500	Apply to actively growing plants in Autumn. DO NOT sow pasture seed for at least 30 days after application.
Pastures – Spray/Graze Techniques				PRECAUTION: An increased quantity of poisonous plants may be eaten by stock using Spray-Graze eg. Caltrop, Capeweed, Paterson's curse, Variegated thistle and deaths could result from causes such as nitrate poisoning. With Paterson's curse, preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock. DO NOT graze horses or pigs on Paterson's curse.
	Amsinckia, Thistles, Capeweed, Doublegee, Mustard, Paterson's curse, Wild turnip, Wild radish, Docks, Geranium, Erodium	SA only	400g/ha	Apply from 6 weeks after opening rains in Autumn until the end of August. Seven days after spraying stock paddock at 4-5 times normal rate, preferably with sheep. Maintain this level of grazing for 6 weeks or until pasture shows signs of over grazing. Then return to normal stocking levels. Use high stocking rates in following Spring to prevent weeds from flowering. Repeat treatments may be required for 2-3 years for complete control.
	Annual thistles, Capeweed, Doublegee, Mustards, Paterson's curse, Turnip, Saffron thistle, Spear thistle	Tas, Vic only		
	Amsinckia, Docks (seedling only), Capeweed, Doublegee, Mustard, Wild radish, Wild turnip, Paterson's curse, Annual thistles	WA only	500g/ha	Apply to saffron thistle at the end of September when plants are running up to flower. Sub-clovers may be damaged at this rate and use is not recommended for all medic pastures.
	Spear thistle, Saffron thistle		900g/ha	
	Melons		1.3kg/ha + 1% oil	
	Docks	Vic only	900g/ha	Apply in September only and follow other recommendations above.
	Caltrop, Capeweed, Charlock, Mustards, Paterson's curse, Shepherd's purse, Saffron, Slender, Spear or Variegated thistle, Turnip weed, Wild radish, Wild turnip	NSW, only	250-900g/ha	Spray actively growing 6-8 week old weeds. Introduce stock 7-10 days after spraying, preferably sheep (cattle are less effective). Stocking rate should be at least 5 times heavier than normal until weeds have been reduced, but before survival of desirable pasture species is threatened. Lucerne and medics may be damaged and should be grazed short before spraying. Other legumes may be affected.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

PASTURE, CEREAL CROPS - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

IN TASMANIA, THIS PRODUCT MAY ONLY BE USED FROM 15 APRIL TO 15 SEPTEMBER UNLESS OTHERWISE PERMITTED BY THE REGISTRAR OF PESTICIDES.

WEED TABLE:

NOTE: Where weeds are to be sprayed in a CROP or PASTURE use only the rates given for the particular crop or situation indicated under the Directions for Use table.

In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed. The rates listed in the Weed Table below are for use where weeds only are present and no crop or pasture is involved. Use at these rates in a crop or pasture other than for spot spraying may cause damage.

This product should only be used in those states where a rate or range of rates is indicated for the particular weeds listed.

WEED TABLE

WEEDS CONTROLLED	APPLICATION RATE							CRITICAL COMMENTS
	CROP						PASTURE	
	Vic	NSW	SA	Qld	Tas	WA	NSW, SA, Qld, Tas only	
Amaranthus spp.	-	400-900g/ha	-	700g/ha	-	-	-	Spray young plants.
Apple of Peru	-	400-900g/ha	-	700g/ha	-	-	-	Spray young plants. Susceptible when young.
Bathurst burr	-	600-900g/ha	-	700g/ha	-	-	600-900g/ha not SA	Spray seedlings only.
Blackberry nightshade	-	400-900g/ha	-	700g/ha	-	-	-	
California burr	-	600-900g/ha	-	700g/ha	-	-	600-900g/ha not SA	Spray seedlings only.
Cape tulip	-	-	-	-	-	500g-1kg/ha	-	Low rate for cormils only
Capeweed	1.1kg/ha	-	1.3kg/ha	-	1.1kg/ha	-	1.3-2.1kg/ha	Spray seedlings to rosette stage.
Caltrop	-	600g-1.3kg/ha	-	700g/ha	-	-	-	Moderately susceptible.
Charlock	400-600g/ha	400-900g/ha	400g/ha	-	1.1kg/ha	-	600-900g/ha	Spray at rosette stage.
Clover	-	900g/ha	-	-	-	-	-	
Common ice plant	-	-	900g/ha	-	-	-	-	
Docks	1.1kg/ha	-	1.1kg/ha	1.1kg/ha	1.1kg/ha	900g/ha	2.5kg/ha SA only	Spray at multiple leaf stage. - effective only on seedlings.
Fat hen	-	400g-1.3kg/ha	-	700g/ha	1.1kg/ha	-	-	Spray pre-flowering.
Fumitory - Red	-	-	1.3kg/ha	-	-	-	-	
Fumitory - White	600g/ha	-	400g/ha	-	-	-	-	Spray at multiple leaf stage.
Hexham scent/Melilotus	1.1kg/ha	-	900g/ha	1.1kg/ha	-	-	900g-1.3kg/ha	Spray multiple leaf stage before seeding
Hoary cress	700g-1.1kg/ha	900g-1.3kg/ha	1.1kg/ha	1.1kg/ha	-	-	1.1-1.3kg/ha	Spray rosettes and pre-flowering.
Hogweed/Wireweed	1.1kg/ha	-	-	1.1kg/ha	-	-	-	Spray at multiple leaf stage (Vic). Spray at seedling and young plant stage (Qld).
Horehound	-	-	1.1kg/ha	-	-	-	1.8-2.5kg/ha SA only	Spray seedlings.
Khaki weed	-	-	-	-	-	-	900g-1.8kg/ha not SA	Spray seedlings only.
Lincoln weed	-	-	1.3kg/ha	-	-	-	-	Spray early rosettes.
London rocket	-	-	-	-	-	600g/ha	-	
Lupins	-	600g-1.3kg/ha	-	-	-	-	-	
Mexican poppy	-	-	-	1.1kg/ha	-	-	-	Spray seedlings – plants become more resistant with age.
Mintweed	-	900g/ha	-	700g/ha	-	-	-	Spray seedlings – resistant in later stages.
Mustards	250-400g/ha	400-900g/ha	400g-1.1kg/ha	700g/ha	-	600g/ha	400-900g/ha	Spray at 2-4 leaf up to rosette stage.
New Zealand spinach	-	900g-1.3kg/ha	-	-	-	-	-	
Noogoora burr	-	600-900g/ha	-	700g/ha	-	-	600-900g/ha not SA	Spray seedlings only.
Paterson's curse	-	900g-1.3kg/ha	-	1.1kg/ha	-	1.1kg/ha	1.3-1.8kg/ha	Spray rosettes or before plants have 10 leaves. Later stages harder to kill.
Potato weed	-	400-900g/ha	-	700g/ha	-	-	-	
Rapeseed	-	600-1.3kg/ha	-	-	-	-	-	
Rough poppy	-	900g/ha	-	-	-	-	-	
Safflower	-	400-900g/ha	-	-	-	-	-	
Shepherds purse	-	900g-1.3kg/ha	-	-	1.1kg/ha	-	600-900g/ha	Spray young rosettes.
Skeleton weed	1.1kg/ha	900g-1.3kg/ha	1.1kg/ha	-	-	-	1.1-1.8kg/ha	Spray rosettes before aerial growth commences.
Sorrel	1.1kg/ha	1.3kg/ha	1.1kg/ha	-	-	-	-	Only moderately susceptible.
Speedwell - Ivy leaf	-	-	900g/ha	-	-	-	-	

WEEDS CONTROLLED	APPLICATION RATE							CRITICAL COMMENTS
	CROP						PASTURE	
	Vic	NSW	SA	Qld	Tas	WA	NSW, SA, Qld, Tas only	
Spiny emex	-	-	-	1.1kg/ha	-	-	-	Only young plants are susceptible.
Stinkwort	-	600g-1.1kg/ha	-	-	-	-	-	
Storkbill/Erodium	-	-	-	-	1.1kg/ha	-	1.3-2.5kg/ha	Spray seedlings to young rosettes.
Sunflower (seedlings)	1.1kg/ha	400g-1.1kg/ha	-	700g/ha	-	-	-	
Thistles: -Californian	-	-	-	-	2.1kg/ha	-	2.5-2.9kg/ha	Repeated applications may be necessary. (NSW, Tas only)
- Saffron	900g/ha	400g-1.3kg/ha	1.1kg/ha	1.1kg/ha	800g/ha	900g/ha	900g-1.3kg/ha	Low rate only sufficient to control weeds in crops at rosette stage when sprayed early.
- Slender/Shore	-	600g-1.3kg/ha	-	-	1.1kg/ha	-	900g/ha	Suppression only.
- Soldier	1.1kg/ha	-	-	-	-	-	900g-1.3kg/ha not NSW, Tas	Spray young rosette.
- Spear	400g/ha	-	-	-	1.1kg/ha	-	900g-1.3kg/ha	Spray young rosettes.
- Star	-	-	-	-	-	-	1.3-1.9kg/ha SA only	Use higher rate as flower stalk appears.
- Variegated	-	400g-1.3kg/ha	-	700g/ha	1.1kg/ha	-	900g-1.3kg/ha	Spray at rosette stage.
Thornapple	-	600-900g/ha	-	-	-	-	1.3-1.9kg/ha not SA	Spray seedlings only.
Turnip weed/Rapistrum	-	400-900g/ha	-	400g/ha	-	600g/ha	400-900g/ha	
Ward's weed	-	-	900g/ha	-	-	-	-	
Wild cabbage	1.1kg/ha	-	-	-	-	-	-	Spray multiple leaves.
Wild poppy	400g/ha	-	-	-	-	-	800g-1.3kg/ha	Spray rosettes.
Wild radish	1.1kg/ha	1.1-1.3kg/ha	1.1kg/ha	700g/ha	1.1kg/ha	600g/ha	600-900g/ha	Spray up to young rosette stage.
Wild turnip	250-400g/ha	400-900g/ha	250g/ha	-	1.1kg/ha	500g/ha	400-900g/ha	Spray 2-4 leaf up to rosette stage.
Vetches/Tares	1.1kg/ha	-	900g/ha	-	-	-	-	Spray at multiple leaf stage.

IMPORTANT

WHEN APPLIED TO DRY SOILS AT LEAST 15mm (1/2 inch) OF RAIN MUST FALL PRIOR TO THE COMMENCEMENT OF THE PLANT BACK PERIOD.

NOTES:

% In Queensland, no rainfall is required to fall prior to commencement of Plant Back Period for Wheat, Barley and Triticale.

In Queensland, planting of Canola/Rapeseed, Chickpeas and Safflower must be delayed for at least 14 days following rainfall of at least 15mm.

@ In Central Queensland, when using 1L/ha or less of Baton Low, the Plant Back Period for Sorghum and Sunflower is 1 day irrespective of rainfall.

+ In WA the Plant Back Period for Lupins at all rates is 28 days.

Plant Back Periods (days) for Baton Low

CROP	PLANT BACK DAYS FOR NUFARM BATON LOW		
	RATES		
	Up to 400g	400-900g	900g-1.3kga
Balansa clover	7	7	10
Barley %	1	1	3
Chickpeas #	7	14	21
Cotton	10	14	21
Faba beans	7	7	10
Field peas	7	14	14
Lentils	7	7	10
Linseed	7	7	14
Lucerne	7	7	10
Lupins +	7	14	21
Medic	7	7	10
Narbon beans	7	7	10
Navybean	10	10	14
Oats	3	3	7
Perennial ryegrass	7	7	10
Persian clover	7	7	10
Phalaris	7	7	10
Canola / Rapeseed #	14	21	28
Rice	7	7	14
Safflower #	7	14	21
Sorghum @	3	7	10

CROP	PLANT BACK DAYS FOR NUFARM BATON LOW		
	RATES		
	Up to 400g	400-900g	900g-1.3kga
Soybean	14	14	21
Sub-clover	7	7	10
Sunflower @	7	10	14
Triticale %	1	3	7
Vetch	7	7	10
Wheat %	1	3	7
White clover	7	7	10

GENERAL INSTRUCTIONS

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions and First Aid Instructions.

MIXING

1kg Water Soluble Bag:

Mix product only with water. Just add the recommended water to your spray tank then remove the water soluble packet from outer bag, add the appropriate amount into the spray tank, agitate for three to five minutes. This product will make a stable water mixture, it is non-flammable. Adding oil, wetting agent or other surfactant to the spray may be used to increase effectiveness on weeds, but doing so may reduce selectivity to crops, resulting in crop damage.

5kg Measure Pack:

This product mixes readily with water, no pre-mixing is required. During mixing the spray vat should be no more than 60% full of water. Vigorous agitation of vat contents is required during the addition of this product. Pour product steadily when adding the required quantity to the spray vat. DO NOT DUMP all required product into vat at once. When the product has been thoroughly dispersed, add surfactant (for post-emergence application only). Fill tank and agitate to ensure thorough mixing. Continue agitation while spraying. Agitate vigorously from the bottom if allowed to stand. DO NOT allow partly filled spray tank to stand overnight. Reseal part used containers immediately.

APPLICATION INFORMATION

BOOM SPRAYING - Use 30-120L/ha of water.

AERIAL SPRAYING - Use 10-90L/ha of water.

EQUIPMENT MAINTENANCE AND USAGE

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants, unless it has been well washed out with hot soapy water or 1% solution of ammonia, followed by several clear water rinses.

COMPATIBILITY

This product may be tank mixed with Kamba® 500 (dicamba), Bromicide® 200, Flowable Diuron, Glean*, Nuquat®, Propon*, Nu-trazine, Lusta®, SpraySeed*/Revolver® and Trooper® 75-D.

RESISTANT WEEDS WARNING

GROUP	I	HERBICIDE
--------------	----------	------------------

Nufarm Baton Low Herbicide ('Baton Low') is a member of the Phenoxys group of herbicides. Baton Low has the Disruptors of plant cell growth mode of action. For weed resistance management Baton Low is a Group I herbicide. Some naturally occurring weed biotypes resistant to Baton Low and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Baton Low or other Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from the failure of Baton Low to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT spray in high winds. DO NOT spray cereals if lucerne is present. DO NOT spray crops or weeds outside the stages indicated in "Critical Comments" as damage, loss of yield or inadequate weed control may result.

Drift Warning: DO NOT use unless wind speed is more than 3 kilometres per hour and less than 15 kilometres per hour as measured at the application site. DO NOT apply with smaller than coarse to very coarse spray droplets according to ASAE S572 definition for standard nozzles. DO NOT apply under weather conditions, or from spraying equipment that may cause spray to drift onto nearby susceptible plant/crops, cropping lands or pastures. Avoid spray drift onto susceptible crops such as cotton, tobacco, tomatoes, vines, fruit trees, vegetables, legume crops and pastures, oilseed crops, susceptible trees (e.g. Kurrajongs, Belahs, Eucalypts) and ornamentals.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used container.

STORAGE AND DISPOSAL

Store in the closed, original container, in a dry, cool, well ventilated area away out of direct sunlight. DO NOT dispose of undiluted chemicals on site. Puncture or shred and bury empty outer containers in a local authority landfill. If not available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray, wear PVC or rubber apron, elbow length PVC gloves and face shield. When using the prepared spray, wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone 13 11 26).

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS), which can be obtained from your supplier or from the Nufarm website – www.nufarm.com.au

In case of emergency: Phone 1600 033 498 Ask for shift supervisor. Toll free 24 hours.

CONDITIONS OF SALE

"Any provisions or rights under the Trade Practices 1974 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute."

Nufarm Australia Limited,
ACN 004 377 780
103-105 Pipe Road,
Laverton North Victoria 3026
Tel: (03) 9282 1000
Fax: (03) 9282 1001

© Nuquat, Bromicide, Kamba, Lusta, Revolver and Trooper are registered trademarks of Nufarm Australia Limited.

* Other registered trademarks