

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm
Estercide[®] 800
Herbicide

ACTIVE CONSTITUENT: 800 g/L 2,4-D present as the ethyl ester

GROUP  HERBICIDE

For the control of broadleaved weeds before direct drilling or sowing of cereals and pastures; and in cereal crops (not oats), pastures, sugar cane and non-agricultural areas as per the Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET
BEFORE USING THIS PRODUCT.

Contents: 5 Litres
2 Litres
20 Litres
110 Litres
200 Litres
1000 Litres

Nufarm Australia Limited
ACN 004 377 780
103-105 Pipe Road
Laverton North Victoria 3026
Tel: (03) 9282 1000
Fax: (03) 9282 1001



® Estercide is a registered trademark
of Nufarm Australia Limited.

GENERAL INSTRUCTIONS

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions and First Aid Instructions.

DO NOT spray in high winds. DO NOT spray crops or weeds outside the stages indicated in the Critical Comments as damage, loss of yield or inadequate weed control may result.

STORAGE AND DISPOSAL (2L, 5L, 20L, 200L only)

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and products should not be burnt.

Refillable containers (1000L only):

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Envirodrum 110L Mini Bulk Returnable Container

Store the original sealed Envirodrum in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Do not tamper with the Micro Matic valve or the security seal. Do not contaminate the Envirodrum with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Nufarm Australia Limited.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray, wear PVC or rubber apron, elbow length PVC gloves and face shield. When using the prepared spray, wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone 13 11 26).

MATERIAL SAFETY DATA SHEET

For further information, refer to the Material Safety Data Sheet (MSDS)

CONDITIONS OF SALE

"Nufarm Australia Limited ('Nufarm') shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Nufarm's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Nufarm has any authority to add to or alter these conditions."

NRA Approval No.: 31209/0902

In case of emergency: Phone 1800 033 498
Ask for shift supervisor. Toll free 24 hours.

Batch No.:

Date of Manufacture:

DrumMuster Logo (2L, 5L, 20L, 200L only)

Envirodrum logo (110L only)

Barcode

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm
Estercide[®] 800
Herbicide

ACTIVE CONSTITUENT: 800 g/L 2,4-D present as the ethyl ester

GROUP  HERBICIDE

For the control of broadleaved weeds before direct drilling or sowing of cereals and pastures; and in cereal crops (not oats), pastures, sugar cane and non-agricultural areas as per the Directions for Use table.

READ COMPLETE DIRECTIONS FOR USE
BEFORE USING THIS PRODUCT.

Nufarm Australia Limited
ACN 004 377 780
103-105 Pipe Road
Laverton North Victoria 3026
Tel: (03) 9282 1000
Fax: (03) 9282 1001

NRA Approval No.: 31209/0902

© Estercide is a registered trademark
of Nufarm Australia Limited.



DIRECTIONS FOR USE:**RESTRAINTS:** Avoid spraying if rain is likely within 6 hours or if strong winds prevail.

DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

1. FIELD CROPS

Situation & Crop	Weeds	State	Rate	With-holding Period (days)	Critical Comments
Wheat, Barley	Refer Weed Table	Vic only	180-700mL/ha	7	<p>CROP STAGES: ALL CEREALS Apply at tillered to boot stages. (Vic only) Apply after when the first node can be felt at the base of the tiller and before swelling of the head can be felt in a tiller. (NSW only) Apply from mid-tillering to before boot stage. (Qld only) Apply from completion of tillering to early jointing stage. (SA only) Apply from the 5 leaf stage up to jointing stage (Zadoks 15-33). Apply only at 6 leaf stage for cranbrook and jacup wheats (Zadoks 16) to avoid possible damage (WA only).</p>
		Qld, NSW only	350-700mL/ha		
		SA only	200-700mL/ha		
		WA only	700mL/ha		
Triticale		Qld, NSW only	350-700mL/ha		
		SA only	200-700mL/ha		
		Vic only	180-700mL/ha		
Cereal Rye		NSW, Qld only	350-700mL/ha		
		Vic only	180-700mL/ha		
Sugar Cane		Qld only	2.1-4L/ha	Pre-emergence	
Fallow- Stubble spray prior to Direct Drilling or Sowing Winter cereals, Grain legumes (Peanuts – Qld only), Canola		All States except Tasmania	180-700mL/ha		<p>Observe the plant back periods given in the table in this leaflet. Must be tank mixed with a knockdown herbicide such as Roundup[®]Max, Credit[®] plus Bonus[®], Roundup[®]CT, Glyphosate CT, Nuquat[®] 250 or Spray Seed*. Select appropriate rate from the Weed Table. For skeleton weed, spraying should only be done 6-8 weeks before anticipated sowing date and subsequent cultivation limited to a minimum.</p>
Harvest Aid or Salvage Spray - Winter Cereals	Broadleaf Weeds	All States except Tasmania	1.4L/ha		<p>Apply after dough stage of crop. Interval between application and effectiveness is 10-20 days. For dessication of green matter, estimate harvest date and spray approximately 14 days earlier. Rain between spraying and actual harvest can negate results. NB. Where thistles are tall and branching above crop, spraying can turn the branches down into the crop, presenting more stalks to cause header comb blockages. Spraying may increase seed contamination of harvest by accelerating maturity. Do not use with undersown legumes that have not set seed.</p>
	Refer Weed Table				

2. PASTURES, NON-AGRICULTURAL, INDUSTRIAL

Situation & Crop	Weeds	State	Rate	With-holding Period (days)	Critical Comments
Improved Pasture containing Clover	Refer Weed Table	Qld, NSW, SA only	350-700mL/ha	7	Clover must be well covered by the grass or extensive damage may result.
Pastures - non legumes, Rights of Way, Industrial		Qld, NSW, SA, WA only	700mL-4L/ha		Control of most perennial weeds, but due to the rooting habits of most species control may take a number of years. Damage may result to legumes in pasture.
		Vic only	700mL-5.6L 60-500mL/ha		Boom spray. Spot spraying.
Pastures - Direct Drilling or Surface Sowing	Charlock, Clover, Medics, Mustards, Paterson's Curse, Saffron, Slender, Variegated and Spear Thistles, Turnip Weed, Wild Radish, Wild Turnip	Qld, NSW, WA, Vic only	700mL-1.25L/ha (Aerial application)		Apply to young, actively growing weeds. SOWING: Do not sow pasture seeds for at least 21 days after application. If soil moisture is dry, delay sowing for at least 30 days.
	As above plus: Capeweed, Wireweed, Storksbill / Erodium, Flatweed, Horehound (seedlings), Skeleton Weed, Nodding or Star Thistles		700mL-1L/ha (Ground application)		
	St Johns Wort		2.8-4L/ha (Aerial or Ground)		
	All of the above plus grasses		As above plus Propon® or Roundup ^ Max or Credit® plus Bonus® or Glyphosate CT or Roundup^ CT or Weedmaster® Duo		

3. SPOT SPRAYING

Situation & Crop	Weeds	State	Rate	With-holding Period (days)	Critical Comments
Spot spraying (All situations)	Refer to Weed Table	NSW, Qld, Vic, SA, WA only	1/100th of rate on Weed Table per 10L water per 100m ²	7	Apply through Knapsack. Thorough wetting of weed is essential.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: PASTURE, CEREAL CROPS - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

WEED TABLE:

NOTE: Where weeds are to be sprayed in a CROP or PASTURE use only the rates given for the crop in the table below. In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed. This product should only be used in those states where a rate or range of rates is indicated for the particular weeds listed.

Weeds	Application Rate per Hectare							Critical Comments
	Crop					Pasture – Non-legumes		
	Vic	SA	NSW	Qld	WA	Vic	NSW, SA, Qld, WA only	
Amaranthus spp.	-	-	700mL	-	-	-	-	
Angled Onion	-	-	-	-	-	2.8L	700mL-1.4L	Spray when buds forming or early flowering.
Apple of Sodom	-	-	-	-	-	-	2.5-2.8L	
Bathurst Burr	-	-	700mL	-	-	1.4-2.8L	1-1.4L	Spray from seedling to pre-flowering. Use higher rate as plant matures. In SA use 1.4 - 2.8L/ha.
Black Knapweed	-	-	-	-	-	1.8L	-	Spray before flowering. DO NOT cultivate these infestations.
Buffalo Burr	-	-	-	-	-	-	700mL-1L (not Qld and WA)	Spray from seedling to pre-flowering. Use higher rate as plant matures.
California Burr	-	-	700mL	-	-	1.4-2.8L	1-1.4L (not SA)	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Caltrop	-	-	500-700mL	-	-	1.4-2.8L	-	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Cape Tulip	-	-	-	-	700mL	2.8L	1.4-2.8L	Spray before flowering.
Capeweed	700mL	700mL	450-700mL	-	-	-	1-1.4L	Spray up to rosette stage.
Charlock	350mL	350mL	350mL	-	-	-	700mL	Spray up to rosette stage.
Clover	-	-	500-700mL	-	-	-	700mL	
Colocynth	-	-	-	-	-	2.8L	-	Spray at seedling stage only.
Deadnettle	-	-	700mL	-	-	-	-	
Devil's Claw	-	-	700mL	-	-	1.1L	1-1.4L (not SA)	Spray prior to pods forming.
Dock	700mL	700mL	-	700mL	700mL	-	1.4-2.1L	Spray at rosette stage to kill top growth only.
Fat Hen	-	-	350-700mL	-	-	-	-	
Flatweed	-	-	700mL	-	-	-	700mL-1L	In SA use 2.1 - 2.8L/ha.
Fumitory - red	-	700mL	700mL	-	-	-	2.1-2.8L	Spray up to rosette stage.
Fumitory - white	700mL	350mL	700mL	-	-	-	2.1-2.8L	Spray up to rosette stage.
Galvanised Burr	-	-	-	-	-	4L	4L (not Qld and WA)	Spray from seedling to pre-flowering.
Goosefoots	-	-	700mL	-	-	-	-	
Hard Head or Russian Knapweed	-	-	-	-	-	2.8-4.4L	-	Spray before flowering.
Hoary Cress Whiteweed	-	700mL	700mL	-	-	1.4-2.8L	1.4-1.8L	Spray from late rosette to pre-flowering.
Hogweed / Wireweed	700mL	700mL	700mL	700mL	-	-	1-1.4L	Spray up to rosette stage.

Weeds	Application Rate per Hectare							Critical Comments
	Crop					Pasture – Non-legumes		
	Vic	SA	NSW	Qld	WA	Vic	NSW, SA, Qld, WA only	
Horehound (seedlings)	-	700mL	-	-	-	-	1.4-2.8L	Late Autumn to early Spring.
Iron Weed, Corn Gromwell	-	-	700mL	-	-	-	1-1.4L	
Khaki Weed	-	-	-	-	-	-	700mL-1L (not SA)	Spray young seedlings only.
Lincoln Weed	-	700mL	-	-	-	-	-	Autumn spray before sowing improves control.
London Rocket	-	-	-	-	500mL	-	-	
Lupins	700mL	-	350-700mL	-	-	-	-	Spray up to rosette stage.
Melilotus /Hexham Scent	700mL	700mL	-	700mL	-	-	1-1.4L	Spray up to rosette stage.
Melons - Camel, Paddy	-	-	350-700mL	-	-	-	-	
Mustards	280mL	200-700mL	350-700mL	500mL	500mL	2.8L	700mL-1L	Spray up to rosette stage.
Mexican Poppy	-	-	700mL	-	-	-	700mL-1L	Spray rosette stage and before flowering. In SA use 2-3L/ha.
Mintweed	-	-	700mL	500mL	-	-	700mL-1L	
Muskweed	700mL	-	-	-	-	-	-	Spray up to rosette stage.
New Zealand Spinach	-	-	700mL	-	-	-	-	
Noogoora Burr	-	-	700mL	-	-	1.4-2.8L	1.4-2.8L	Spray seedling to pre-flowering.
Nut Grass	-	-	-	-	-	2.8-4.4L	-	Spray within 4 weeks of foliage emergence, repeat spray necessary.
Paterson's Curse	-	-	700mL	-	-	1.4-2.8L	700mL-1.4L	Spray seedling to rosette stage.
Poppy Wild	350mL	-	-	-	-	-	1.8-2.5L	Spray up to rosette stage.
Ragwort	-	-	-	-	-	2.8L	2.8L	Spray at rosette to cabbage stage.
Rapeseed	700mL	-	350-700mL	-	-	-	-	Spray up to rosette stage.
Rough Poppy	-	350mL	350-700mL	-	-	-	700mL	Spray young seedlings only.
St Johns Wort	-	-	-	-	-	2.8-4.4L	2.8-4L	Spray before flowering. Spray before plants 40cm high.
Safflower	-	-	350-700mL	-	-	-	-	
Sand Mustard / Sand Rocket	-	-	-	-	-	2.8L	-	Spray before flowering.
Shepherds Purse	-	-	700mL	-	-	-	-	
Silverleaf Nightshade	-	-	-	-	-	2.8L	-	Spray at flowering. Fallowland: controls top growth only.
Skeleton Weed	-	700mL	700mL	-	-	2.8L	1-1.4L	Spray rosettes before aerial growth commences.
Stingless Nettle (Deadnettle)	-	700mL	-	-	-	-	1.8-2.1L	
Stinging Nettle	700mL	-	-	-	-	-	-	Spray up to rosette stage.
Stinkwort	-	-	700mL	-	-	1.4-2.8L	1.4-2.8L	Spray younger plants, use higher rate as plants mature.
Storksbill / Erodium	-	-	700mL	-	-	-	-	

Weeds	Application Rate per Hectare							Critical Comments
	Crop					Pasture – Non-legumes		
	Vic	SA	NSW	Qld	WA	Vic	NSW, SA, Qld, WA only	
Sunflower seedlings	700mL	-	350-700mL	500mL	-	-	-	Spray multiple leaves.
Thistle								
- Golden	-	-	-	-	-	2.8L	2.8L	Spray up to rosette stage.
- Nodding	-	-	-	-	-	2.8L	1-1.4L	Spray rosette to pre-flowering.
- Saffron	550mL	700mL	350-700mL	700mL	700mL	700mL-1.4L	700mL-2.1L	Spray up to rosette stage.
- Slender / Shore	-	-	700mL	-	-	1.4-2.8L	700mL-2.8L	Spray at rosette stage.
- Soldier	-	-	-	-	-	2.8L	-	Spray at rosette stage.
- Spear	700mL	-	-	-	-	700mL-2.1L	1-1.8L	Spray at seedling to rosette stage. Use higher rate as plants mature (pastures).
- Stemless	-	-	-	-	-	2.8L	2.1-2.8L	Spray rosette stage to flowering.
- St Barnaby's	-	-	-	-	-	-	1-1.4L	
- Star	-	-	700mL	-	-	1.4-2.8L	1-1.4L	Spray seedling to rosette stage. Use higher rate as plants mature.
- Variegated	-	-	350-700mL	500mL	-	700mL-2.1L	700mL-1L	Spray at rosette stage. Can cause stock poisoning.
Thornapple	-	-	350-700ml	-	-	2.8L	700mL-1.4L	Spray at seedling stage. In SA use 3L/ha.
Tree Hogweed	700mL	-	-	-	-	-	-	Spray up to rosette stage.
Turnip Weed	-	350mL	350-700mL	350mL	500mL	-	700mL	Spray seedlings only.
Vetches / Tares	700mL	550mL	-	-	-	-	-	
Wards Weed	-	350mL	-	-	-	-	-	Spray at seedling stage.
Wild Cabbage	700mL	-	-	-	-	-	-	Spray up to rosette stage.
Wild Garlic	-	-	-	-	-	5.6L	-	Suppresses aerial growth only.
Wild Mignonette	-	-	-	-	-	2.8L	-	Spray at rosette stage.
Wild Radish	700mL	700mL	350-700mL	700mL	500mL	-	700mL	Spray up to rosette stage.
Wild Sage	-	-	-	-	-	-	2.1-2.8mL	
Wild Teasel	-	-	-	-	-	1.4-2.8L	-	Spray at rosette stage. Use higher rate as plants mature.
Wild Turnip	180mL	200mL	350-700mL	-	350mL	-	700mL	Spray up to rosette stage.

Plant back days for Nufarm Estercide 800

Crop	Rates		
	Up to 440mL/ha	440mL to 860mL/ha	860 to 1.3L/ha
Balansa Clover	7	7	10
Barley %	1	1	3
Chickpeas #	7	14	21
Cotton	10	14	21
Faba Beans	7	7	10
Field Peas	7	14	14
Lentils	7	7	10
Linseed	7	7	14
Lucerne	7	7	10
Lupins +	7	14	21
Medics	7	7	10
Narbon Beans	7	7	10
Navy Beans	10	10	14
Oats	3	3	7
Perennial Ryegrass	7	7	10
Persian Clover	7	7	10
Phalaris	7	7	10
Canola / Rapeseed #	14	21	28
Rice	7	7	14
Safflower #	7	14	21
Sorghum @	3	7	10
Soybean	14	14	21
Sub - Clover	7	7	10
Sunflower @	7	10	14
Triticale %	1	3	7
Vetch	7	7	10
Wheat %	1	3	7
White Clover	7	7	10

IMPORTANT:

WHEN APPLIED TO DRY SOILS AT LEAST 15MM (1/2 INCH) OF RAIN MUST FALL PRIOR TO THE COMMENCEMENT OF THE PLANT BACK PERIOD.

NOTES:

- % In Queensland, no rainfall is required to fall prior to commencement of Plant Back Period for wheat, barley and triticale.
- # In Queensland, planting of canola / rapeseed, chickpeas and safflower must be delayed for at least 14 days following rainfall of at least 15mm
- @ In Central Queensland, when using 625ml/ha or less of Estercide 800, the Plant Back Period for sorghum and sunflower is 1 day irrespective of rainfall.
- + In WA the Plant Back Period for lupins at all rates is 28 days.

GENERAL INSTRUCTIONS

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions and First Aid Instructions.

DO NOT spray in high winds. DO NOT spray crops or weeds outside the stages indicated in the Critical Comments as damage, loss of yield or inadequate weed control may result.

APPLICATION INFORMATION

This product may be used in either high or low volume sprays. Just pour into water and stir.
Boom Spraying - Use 30-120 litres water/ha.
Aerial Spraying - Use 10-90 litres water/ha.

EQUIPMENT MAINTENANCE AND USAGE

Keep the spray unit for herbicides only if possible. Otherwise wash out the unit with hot soapy water followed by several clear water rinses. DO NOT use wooden spray vats as they cannot be cleaned. Hoses cannot be cleaned and new hoses should be fitted when the unit is to be used for any other purpose.

COMPATIBILITY

Kamba[®] 500, Glean[®]/Lusta[™], Nuquat[®] 250, Propon[®], Nu-trazine (900DF + Flowable), Spray•Seed^{*}, Credit[®]/Bonus[®], Weedmaster[®] Duo, Roundup[®]Max, Roundup[®]CT, Nufarm Glyphosate CT.

RESISTANT WEEDS WARNING

Nufarm Estercide 800 Herbicide ("Estercide 800") is a member of the Phenoxys group of herbicides.

Estercide 800 has the Disruptors of plant cell growth mode of action. For weed resistance management Estercide 800 is a Group I herbicide.

Some naturally occurring weed biotypes resistant to Estercide 800 and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Estercide 800 or Group I herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Limited accepts no liability for any losses that may result from the failure of Estercide 800 to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Legume Tolerance: If clovers are present, care should be taken to ensure that they have reached the 3-4 leaf stage before spraying. Rates above 350mL of this product per hectare will destroy most clovers, whilst lucerne and medics are susceptible at any strength.

Drift Warning: DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray drift onto nearby susceptible plants, adjacent crops, crop lands or pastures. In particular, avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, fruit trees, ornamentals, oil seed and legume crops and other susceptible plants and trees (eg kurrajongs, belahs, eucalypts).

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied on any plants at any time as recommended.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT.

DO NOT contaminate streams, rivers or waterways with the chemical or used container.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Envirodrum 110L Mini Bulk Returnable Container

Store the original sealed Envirodrum in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Do not tamper with the Micro Matic valve or the security seal. Do not contaminate the Envirodrum with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Nufarm Australia Limited.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray, wear PVC or rubber apron, elbow length PVC gloves and face shield. When using the prepared spray, wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone 13 11 26).

MATERIAL SAFETY DATA SHEET

For further information, refer to the Material Safety Data Sheet (MSDS)

CONDITIONS OF SALE

“Nufarm Australia Limited ('Nufarm') shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Nufarm's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Nufarm has any authority to add to or alter these conditions.”

NRA Approval No.: 31209/0902

In case of emergency: Phone 1800 033 498

Ask for shift supervisor. Toll free 24 hours.

® Kamba, Nuquat, Propon, Credit, Bonus and Weedmaster are registered trademarks of Nufarm Australia Limited.

^ Roundup is a registered trademark of Monsanto Technology LLC used under licence by Nufarm Australia Limited.

* Other registered trademarks

™ Trademark pending