

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm
LV Estercide[®] 600

Herbicide

**ACTIVE CONSTITUENT: 600g/L 2,4-D
present as the 2-ethylhexyl ester**

GROUP II HERBICIDE

A specially formulated low volatile herbicide for selective control of various weeds in crops, pastures and non-agricultural areas as per the Directions For Use Table.

This is a PHENOXY HERBICIDE that can cause severe damage to susceptible crops such as cotton, grapes, tomatoes, oilseed crops and ornamentals.

**IMPORTANT: READ THE ATTACHED LEAFLET
BEFORE USING THIS PRODUCT.**

Contents: 5 Litres
20 Litres
110 Litres
200 Litres
500 Litres
1000 Litres

Nufarm Australia Limited
ACN 004 377 780
103-105 Pipe Road
Laverton North Victoria 3026
Tel: (03) 9282 1000
Fax: (03) 9282 1001



® Estercide is a registered trademark of Nufarm Australia Limited

GENERAL INSTRUCTIONS

Before opening, carefully read Directions For Use, Precautionary Statements, Safety Directions and First Aid Instructions.

STORAGE AND DISPOSAL

(5L, 20L, 200L only)

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and products should not be burnt.

Refillable containers (500L, 1000L only): Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Envirodrum 110L Mini Bulk Returnable Container

Store the original sealed Envirodrum in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Do not tamper with the Micro Matic valve or the security seal. Do not contaminate the Envirodrum with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Nufarm Australia Limited.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray wear PVC or rubber apron, elbow-length PVC gloves and face shield. When using the prepared spray, wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone 13 11 26).

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS).

CONDITIONS OF SALE

"Nufarm Australia Limited ('Nufarm') shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Nufarm's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Nufarm has any authority to add to or alter these conditions."

APVMA Approval No.: 42229/5L/0206

APVMA Approval No.: 42229/20L/0206

APVMA Approval No.: 42229/110L/0206

APVMA Approval No.: 42229/200L/0206

APVMA Approval No.: 42229/500L/0206

APVMA Approval No.: 42229/1000L/0206

In case of emergency: Phone 1800 033 498

Ask for shift supervisor. Toll free 24 hours.

Batch No.:

Date of Manufacture:

DrumMuster logo (5L, 20L, 200L only)

Envirodrum logo (110 only)

[Barcode]

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm
LV Estercide[®] 600

Herbicide

**ACTIVE CONSTITUENT: 600g/L 2,4-D
present as the 2-ethylhexyl ester**

GROUP II HERBICIDE

A specially formulated low volatile herbicide for selective control of various weeds in crops, pastures and non-agricultural areas as per the Directions For Use Table.

This is a PHENOXY HERBICIDE that can cause severe damage to susceptible crops such as cotton, grapes, tomatoes, oilseed crops and ornamentals.

**READ COMPLETE DIRECTIONS FOR USE
BEFORE USING THIS PRODUCT**

Nufarm Australia Limited
ACN 004 377 780
103-105 Pipe Road
Laverton North Victoria 3026
Tel: (03) 9282 1000
Fax: (03) 9282 1001



APVMA Approval No.: 42229/0206

® Estercide is a registered trademark of Nufarm Australia Limited

DIRECTIONS FOR USE:

RESTRAINTS: AVOID spraying if rain is likely within 6 hours or if strong winds prevail.

DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

DO NOT use when breeze is blowing towards nearby susceptible plants.

1. FIELD CROPS

REFER TO SECTION “SPRAY APPLICATIONS AND DRIFT RISK ASSESSMENT” BEFORE APPLICATION.

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Wheat, Barley,	Refer to Weed Table	Vic only	240-900mL	CROP STAGES: ALL CEREALS Variations between varieties do occur. Check sensitivity and growth stages of varieties before applying. Damage may result if applied too early. Apply at tillered to boot stages. (Vic only). Apply after when the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller. (NSW only). Apply from mid-tillering (5 to 6 fully emerged main stem leaves plus one or more tillers) to before boot stage (visible swelling of the head at the top of the main stem. (Qld only). Apply from completion of tillering to early jointing stage. (SA, Tas only). Apply from the 5 leaf stage up to jointing stage (Zadoks 15-33). Apply only at 6 leaf stage for cranbrook and jacup wheats (Zadoks 16) to avoid possible damage (WA only).
		SA only	260-900mL	
		Qld, NSW only	460-900mL	
		Tas only	700-900mL	
		WA only	900mL	
Triticale	Refer to Weed Table	Qld, NSW only	460-900mL	
		SA only	270-930mL	
		Vic only	240-900mL	
Cereal Rye	Refer to Weed Table	NSW, Qld only	460-900mL	
		Vic only	240-900mL	
Sugar Cane	Refer to Weed Table	Qld only	1.3-2.7L	Post-emergence.
Stubble/Fallow Spray Prior to Direct Drilling or Sowing. Winter Cereals, Grain legumes (Peanuts Qld only), Canola	Refer to Weed Table	All States	240-900mL/ha	Observe the plant back periods given in the table in this leaflet. Must be tank mixed with a knockdown herbicide such as Roundup [^] PowerMAX, Credit [®] plus Bonus [®] , Roundup [^] CT, Glyphosate CT, Nuquat [®] 250 or Spray•Seed [®] /Revolver [®] . Select appropriate rate from the weed table. For skeleton weed, spraying should only be done 6-8 weeks before anticipated sowing date and subsequent cultivation limited to a minimum.
Harvest Aid or Salvage Spray -Winter Cereals	Broadleaf Weeds Refer to Weed Table	All States	1.9L	Apply after dough stage of crop. Interval between application and effectiveness is 10-20 days. For desiccation of green matter, estimate harvest date and apply spray approximately 14 days earlier. Rain between spraying and actual harvest can negate results. NB. Where thistles are tall and branching above the crop, spraying can turn the branches down into the crop, presenting more stalks to cause header comb blockages. Spraying may increase seed contamination of harvest by accelerating maturity. Do not use with undersown legumes that have not set seed.
Potatoes Pre-harvest Preparation	Broadleaf Weeds such as Clover, Variegated Thistle & Cruciferous weeds	Vic, Tas only	1.3-2.7L	Apply approximately 4 to 5 weeks before harvest after the potato haulms have dried off. Use the highest rate where weeds are more than 30cm in height. For boom spraying apply at least 100 litres of spray mixture per hectare. If grasses such as Rye grass and Winter grass are also present add Amitrole T Herbicide.

**2. PASTURES, NON-AGRICULTURAL, INDUSTRIAL
REFER TO SECTION “SPRAY APPLICATIONS AND DRIFT RISK ASSESSMENT”
BEFORE APPLICATION.**

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Improved Pasture containing Clovers	Refer to Weed Table	NSW, Tas, SA, Qld only	460-900mL	Clover must be well covered by the grass or extensive damage may result.
Pastures – non legumes, Rights of Way & Industrial		NSW, Tas, SA, WA, Qld only	900mL-5.3L	Control of most perennial weeds, but due to the rooting habits of most species control may take a number of years. Damage may result to legumes in pasture.
		Vic only	900mL-7.5L 80-700mL	Boom spray. Spot spraying.
Pastures – Direct Drilling or Surface Sowing	Charlock, Clover, Medics, Mustards, Paterson’s Curse, Saffron, Slender, Variegated and Spear Thistles, Turnip Weed, Wild Radish, Wild Turnip	NSW, Qld, WA, Vic, SA, Tas only	900mL-1.7L (Aerial Application)	Apply to young, actively growing weeds. SOWING: Do not sow pasture seed for at least 21 days after application. If soil moisture is dry, delay sowing for at least 30 days.
	As Above plus: Capeweed, Wireweed, Storksbill/Erodium, Flatweed, Horehound (seedlings), Skeleton Weed, Nodding or Star Thistles		900mL-1.3L (Ground Application)	
	St John’s Wort		3.7-5.3L (Aerial or Ground)	
	All of Above plus grasses		As above plus Propon or Roundup [^] PowerMAX or Credit [®] plus Bonus [®] or Glyphosate CT or Roundup [^] CT or Weedmaster [®] Duo	

3. SPOT SPRAYING

Situation & Crop	Weeds Controlled	State	Rate/ha	Critical Comments
Spot Spraying (All Situations)	Refer to Weed Table	All States	1/100th of rate on Weed Table per 10L water per 100m ²	Apply through Knapsack. Thorough wetting of weed is essential.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

IN TASMANIA, THIS PRODUCT MAY ONLY BE USED FROM 15TH APRIL TO 15TH SEPTEMBER UNLESS OTHERWISE PERMITTED BY THE REGISTRAR OF PESTICIDES.

WITHHOLDING PERIOD (WHP): PASTURES, CEREAL CROPS – DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

HARVEST WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

Before opening, carefully read Directions For Use, Precautionary Statements, Safety Directions and First Aid Instructions.

APPLICATION INFORMATION

This product may be used in either high or low volume sprays. Just pour into water and stir.
BOOM SPRAYING - Use 30-100 litres water per hectare.
AERIAL SPRAYING - Use 10-90 litres water per hectare.
NOTE: Refer to the Department of Agriculture/ Primary Industries in your state for the current restricted spraying areas.

SPRAY APPLICATIONS AND DRIFT RISK ASSESSMENT

USE ONLY when wind speed is more than 3 kilometres per hour or less than 15 km/h, as measured by an anemometer at the application site.

USE ONLY coarse to very coarse spray according to the ASAE S572 definition for standard nozzles.

Checklist:

- Have you cleaned/decontaminated your boom sprayer?
- Have you contacted your neighbour prior to spraying?
- Is your sprayer set-up correctly for the particular application?
- Check
 - boom calibration - at nozzle
 - nozzle choice - low drift/what spray quality - coarse to very coarse droplet?
 - boom height
 - speed of intended application
- You must check, determine and record the weather conditions immediately prior to, and immediately after the spray application is made.
- Record
 - Temperatures
 - Relative Humidity
 - Delta T
 - Wind speed (min 3km/hr, max 15km/hr)
 - Is there a temperature inversion?
- Night Spraying - Extra care is required to ensure that inversion conditions are not present. Use smoke generator to determine wind direction and presence of inversion conditions.

EQUIPMENT MAINTENANCE AND USAGE

Keep the spray unit for herbicides only if possible. Otherwise wash out the unit with hot soapy water followed by several clear water rinses. Do not use wooden spray vats as they can not be cleaned. Hoses cannot be cleaned and new hoses should be fitted when the unit is to be used for any other purpose.

COMPATIBILITY

This product can be tank mixed with Kamba[®] 500, Glean[®]/Lusta[™], Nuquat[®] 250, Propon[®], Nu-trazine (900DF & Flowable), Credit[®]/Bonus[®], Weedmaster[®] Duo, Roundup[^] PowerMAX, Roundup[^] CT, Nufarm Glyphosate CT, Ally[®]/Associate[®], Logran[®]/Nugran, Spray-Seed[®]/Revolver[®].

WEED TABLE:

NOTE: Where weeds are to be sprayed in a CROP or PASTURE, use only the rates given for the crop in the table below. In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed. eg. Rumex spp. (docks) and Polygonum spp. (wireweed, climbing buckwheat) are killed to ground level only.

APPLICATION RATE PER HECTARE

WEEDS CONTROLLED	CROP						PASTURES – NON LEGUMES		
	Vic	SA	Tas	NSW	Qld	WA	Vic	NSW, TAS, SA, QLD, WA only	Critical Comments
Amaranthus spp	-	-	-	900mL	-	-	-	-	
Angled Onion	-	-	-	-	-	-	3.7L	0.9-1.9L	Spray when buds forming or early flowering.
Apple of Sodom	-	-	-	-	-	-	-	3.3-3.7L	
Bathurst Burr	-	-	-	900mL	-	-	1.9-3.7L	1.9-3.7L	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Black Knapweed	-	-	-	-	-	-	3.7L	-	Spray before flowering. DO NOT cultivate these infestations.
Buffalo Burr	-	-	-	-	-	-	-	900mL-1.3L (Not Qld & WA)	Spray from seedling to pre-flowering. Use higher rate as plant matures.
California Burr	-	-	-	900mL	-	-	1.9-3.7L	1.3-1.9L (Not SA)	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Caltrop	-	-	-	700-900mL	-	-	1.9-3.7L	-	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Cape Tulip	-	-	-	-	-	1.3L	3.7L	1.9-3.7L	Spray before flowering.
Capeweed	900mL	900mL	900mL	600-900mL	-	-	-	2.8-3.7L	Spray up to rosette stage.
Charlock	460mL	460mL	900mL	460mL	-	-	-	900mL	Spray up to rosette stage.
Clover	-	-	-	700-900mL	-	-	-	900mL	
Colocynth	-	-	-	-	-	-	3.7L	-	Spray at seedling stage only.
Deadnettle	-	-	-	900mL	-	-	-	-	
Devil's Claw	-	-	-	900mL	-	-	1.5L	1.3-1.9L (Not SA)	Spray prior to pods forming.
Dock	900mL	900mL	-	-	900mL	900mL	-	1.9-2.8L	Spray at rosette stage to kill top growth only.
Fat Hen	-	-	-	460-900mL	-	-	-	-	
Flatweed	-	-	-	900mL	-	-	-	2.8-3.7L	
Fumitory (red)	-	900mL	-	900mL	-	-	-	2.8-3.7L	Spray up to rosette stage.
Fumitory (white)	900mL	460mL	-	900mL	-	-	-	2.8-3.7L	Spray up to rosette stage.
Galvanised Burr	-	-	-	-	-	-	5.3L	5.3L (Not Qld & WA)	Spray from seedling to pre-flowering.
Goosefoots	-	-	-	900mL	-	-	-	-	
Hard Head or Russian Knapweed	-	-	-	-	-	-	3.7-5.9L	-	Spray before flowering.
Hogweed, Wireweed	900mL	900mL	-	900mL	900mL	-	-	1.3-1.9L (Not SA)	Spray up to rosette stage.

APPLICATION RATE PER HECTARE continued

WEEDS CONTROLLED	CROP						PASTURES – NON LEGUMES		
	Vic	SA	Tas	NSW	Qld	WA	Vic	NSW, TAS, SA, QLD, WA only	Critical Comments
Hoary Cress, Whiteweed	-	900mL	900mL	900mL	-	-	1.9-3.7L	1.9-2.4L	Spray from late rosette to pre-flowering.
Horehound (seedlings)	-	900mL	-	-	-	950mL	-	1.9-3.7L	Late Autumn to early Spring.
Ironweed, Corn Gromwell,	-	-	-	900mL	-	-	-	1.3-1.9L	
Khaki Weed	-	-	-	-	-	-	-	900mL-1.3L (Not SA)	Spray young seedlings only.
Lincoln Weed	-	900mL	-	-	-	-	-	-	Autumn spray before sowing improves control.
London Rocket	-	-	-	-	-	650mL	-	1.8-2.8L (WA only)	
Lupins	900mL	-	-	460mL-900mL	-	-	-	-	Spray up to rosette stage.
Mellilotus/Hexham Scent	900mL	900mL	-	-	900mL	-	-	1.3-1.9L	Spray up to rosette stage.
Melons – camel, paddy	-	-	-	460mL-900mL	-	-	-	-	
Mustards	370mL	260mL-900mL	900mL	460mL-900mL	700mL	700mL	3.7L	1.9-2.8L	Spray up to rosette stage.
Mexican Poppy	-	2.6-4L	-	900mL	-	950mL	-	900mL-1.3L (1.3-1.7L WA only)	Spray rosette stage and before flowering.
Mintweed	-	-	-	900mL	700mL	-	-	900mL-1.3L	Spray active seedlings only.
Muskweed	900mL	-	-	-	-	-	-	-	Spray up to rosette stage.
New Zealand Spinach	-	-	-	900mL	-	-	-	-	
Noogoora Burr	-	-	-	900mL	-	-	1.9-3.7L	1.9-3.7L	Spray seedling to pre-flowering.
Nut Grass	-	-	-	-	-	-	3.7-5.9L	-	Spray within 4 weeks of foliage emergence, repeat spray necessary.
Paterson's Curse	-	-	-	900mL	-	950mL	1.9-3.7L	900mL-1.9L (1.3-1.7L WA only)	Spray seedling to rosette stage.
Poppy Wild	460mL	-	-	-	-	-	-	2.4-3.3L	Spray up to rosette stage.
Ragwort	-	-	-	-	-	-	3.7L	3.7L	Spray at rosette to cabbage stage.
Rapeseed	900mL	-	-	460mL-900mL	-	-	-	-	Spray up to rosette stage.
Rapistrum	-	-	-	-	-	650mL	-	950mL (WA only)	
Rough Poppy	-	460mL	-	460mL-900mL	-	-	-	900mL	Spray young seedlings only.
St John's Wort	-	-	-	-	-	-	3.7-5.9L	3.7- 5.3L	Spray before flowering. Spray before plants are 40cm high.
Safflower	-	-	-	460mL-900mL	-	-	-	-	

APPLICATION RATE PER HECTARE continued

WEEDS CONTROLLED	CROP						PASTURES – NON LEGUMES		
	Vic	SA	Tas	NSW	Qld	WA	Vic	NSW, TAS, SA, QLD, WA only	Critical Comments
Sand Mustard/Sand Rocket	-	-	-	-	-	-	3.7L	-	Spray before flowering.
Shepherd's Purse	-	-	-	900mL	-	-	-	-	
Silver Leaf Nightshade	-	-	-	-	-	-	3.7L	-	Spray at flowering. Fallowland: controls top growth only.
Skeleton Weed	-	900mL	-	900mL	-	-	3.7L	1.3-1.9L	Spray rosettes before aerial growth commences.
Stingless Nettle (Deadnettle)	-	900mL	900mL	-	-	-	-	2.4-2.8L	
Stinging Nettle	900mL	-	-	-	-	-	-	-	Spray up to rosette stage.
Stinkwort	-	-	-	900mL	-	-	1.9-3.7L	1.9-3.7L	Spray younger plants. Use higher rate as plants mature.
Storksbill/Erodium	-	-	-	900mL	-	-	-	-	
Sunflower Seedlings	900mL	-	-	460mL-900mL	700mL	-	-	-	Spray multiple leaves.
Thistles:									
- Golden	-	-	-	-	-	-	3.7L	3.7L	Spray at rosette stage.
- Nodding	-	-	-	-	-	-	3.7L	1.3-1.9L	Spray rosette stage to pre-flowering.
- Saffron	700mL	900mL	-	460mL-900mL	900mL	900mL	900mL-1.9L	900mL-2.8L	Spray up to rosette stage.
- Sheep	-	-	-	-	-	950mL	-	950mL-3.7L (WA only)	
- Slender, Shore	-	-	900mL	900mL	-	-	1.9-3.7L	900mL-3.7L	Spray at rosette stage.
- Soldier	-	-	-	-	-	-	3.7L	-	Spray at rosette stage.
- Spear	900mL	-	900mL	-	-	-	900mL-2.8L	1.3-2.4L	Spray at seedling to rosette stage. Use higher rate as plants mature (pastures).
- Stemless	-	-	-	-	-	-	3.7L	2.8-3.7L	Spray rosette stage to flowering.
- St Barnaby's	-	-	-	-	-	-	-	1.3-1.9L	
- Star	-	-	-	900mL	-	-	1.9-3.7L	1.3-1.9L	Spray seedling to rosette stage. Use higher rate as plants mature.
- Variegated	-	-	900mL	460mL-900mL	700mL	-	900mL-2.8L	900mL-3.7L	Spray at rosette stage. Can cause stock poisoning
Thornapple	-	4L	-	460mL-900mL	-	-	3.7L	900mL-1.9L	Spray at seedling stage.
Tree Hogweed	900mL	-	-	-	-	-	-	-	Spray up to rosette stage.
Turnip Weed	-	460mL	-	460mL-900mL	460mL	700mL	-	900mL	Spray seedlings only.
Vetches/Tares	900mL	700mL	900mL	-	-	-	-	-	
Wards Weed	-	460mL	-	-	-	-	-	-	Spray at seedling stage.
Wild Cabbage	900mL	-	-	-	-	-	-	-	Spray up to rosette stage.

APPLICATION RATE PER HECTARE continued

WEEDS CONTROLLED	CROP						PASTURES – NON LEGUMES		
	Vic	SA	Tas	NSW	Qld	WA	Vic	NSW, TAS, SA, QLD, WA only	Critical Comments
Wild Garlic	-	-	-	-	-	-	7.5L	-	Suppresses aerial growth only.
Wild Mignonette	-	-	-	-	-	950mL	3.7L	-	Spray at rosette stage.
Wild Mustard	-	-	-	-	-	650mL	-	1.8-2.8L (WA only)	
Wild Radish	900mL	900mL	900mL	460mL-900mL	900mL	650mL	-	900mL (950mL WA only)	Spray up to rosette stage.
Wild Sage	-	-	-	-	-	-	-	2.8-3.7L	
Wild Teasel	-	-	-	-	-	-	1.9-3.7L	-	Spray at rosette stage. Use higher rate as plants mature.
Wild Turnip	240mL	260mL	900mL	460mL-900mL	-	450mL	-	900mL (950mL WA only)	Spray up to rosette stage.

Plant back days for Nufarm LV Estercide 600

Crop	Rates		
	Up to 580mL/ha	580mL - 1.15L/ha	1.15- 1.8L/ha
Balansa Clover	7	7	10
Barley %	1	1	3
Chickpeas #	7	14	21
Cotton	10	14	21
Faba Beans	7	7	10
Field Peas	7	14	14
Lentils	7	7	10
Linseed	7	7	14
Lucerne	7	7	10
Lupins +	7	14	21
Medics	7	7	10
Narbon Beans	7	7	10
Navybean	10	10	14
Oats	3	3	7
Perennial Ryegrass	7	7	10
Persian Clover	7	7	10
Phalaris	7	7	10
Canola / Rapeseed #	14	21	28
Rice	7	7	14
Safflower #	7	14	21
Sorghum @	3	7	10
Soybean	14	14	21
Sub-Clover	7	7	10
Sunflower @	7	10	14
Triticale %	1	3	7
Vetch	7	7	10
Wheat %	1	3	7
White Clover	7	7	10

IMPORTANT:

WHEN APPLIED TO DRY SOILS AT LEAST 15mm ($\frac{1}{2}$ inch) OF RAIN MUST FALL PRIOR TO THE COMMENCEMENT OF THE PLANT BACK PERIOD.

NOTES:

- % In Queensland, no rainfall is required to fall prior to commencement of Plant Back Period for wheat, barley and triticale.
- # In Queensland, planting of canola / rapeseed, chickpeas and safflower must be delayed for at least 14 days following rainfall of at least 15mm.
- @ In Central Queensland, when using 830mL/ha or less of LV Estercide 600, the Plant Back Period for sorghum and sunflower is 1 day irrespective of rainfall.
- + In WA the Plant Back Period for lupins at all rates is 28 days.

RESISTANT WEEDS WARNING

GROUP I HERBICIDE

Nufarm LV Estercide 600 Herbicide ('LV Estercide 600') is a member of the Phenoxys group of herbicides. LV Estercide 600 has the Disruptors of plant cell growth mode of action. For weed resistance management LV Estercide 600 is a Group I herbicide. Some naturally-occurring weed biotypes resistant to LV Estercide 600 and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by LV Estercide 600 or other Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from the failure of LV Estercide 600 to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT spray in high winds. DO NOT spray crops or weeds outside the stages indicated in "Critical Comments" as damage, loss of yield or inadequate weed control may result.

Drift Warning:

- DO NOT use unless wind speed is more than 3 kilometres per hour and less than 15 kilometres per hour as measured at the application site.
- DO NOT apply with smaller than coarse to very coarse spray droplets according to the ASAE S572 definition for standard nozzles.
- Direct spray contact or even slight drift may cause severe injury or destruction of any growing crop or other desirable plants including trees and native vegetation.
- DO NOT use when breeze is blowing towards nearby desirable plants.
- DO NOT apply under meteorological conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants, adjacent crops, crop lands or pastures.
- Equipment settings which produce fine droplets (150 microns or less), "still air" and hot dry days all contribute to drift.
- This is a phenoxy herbicide that can cause severe damage to susceptible crops such as cotton, tobacco, tomatoes, flowers, vegetables, vines, fruit trees, legume crops and pastures, oil seed crops or other susceptible crops and trees (eg Kurrajongs, Belahs, Eucalypts).

Legume Tolerance: If clovers are present, care should be taken to ensure that they have reached the 3-4 leaf stage before spraying. Rates above 460mL of this product per hectare will destroy most clovers, whilst lucerne and medics are susceptible at any strength.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used container.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray

tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Envirodrum 110L Mini Bulk Returnable Container

Store the original sealed Envirodrum in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Do not tamper with the Micro Matic valve or the security seal. Do not contaminate the Envirodrum with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Nufarm Australia Limited.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray wear PVC or rubber apron, elbow-length PVC gloves and face shield. When using the prepared spray, wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone 13 11 26).

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS).

CONDITIONS OF SALE

"Nufarm Australia Limited ('Nufarm') shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Nufarm's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Nufarm has any authority to add to or alter these conditions."

APVMA Approval No.: 42229/0206

In case of emergency: Phone 1800 033 498

Ask for shift supervisor. Toll free 24 hours.

® Associate, Kamba, Propon, Credit, Bonus, Revolver, Weedmaster and Nuquat are registered trademarks of Nufarm Australia Limited.

^ Roundup PowerMAX is a registered trademark of Monsanto Technology LLC used under licence by Nufarm Australia Limited.

* Other trademarks
™ Trademark pending