

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Dow AgroSciences

Esteron™ LV

Herbicide

ACTIVE CONSTITUENT: 680 g/L 2,4-D present as the ethylhexyl ester

GROUP I HERBICIDE

A low volatile herbicide for selective control of various weeds in crops, pastures and non-agricultural areas according to the Directions for Use.

This is a PHENOXY HERBICIDE that can cause severe damage to susceptible crops. Please read the Drift Warning Statement.

Dow AgroSciences Australia Limited ABN 24 003 771 659
20 Rodborough Road FRENCHS FOREST NSW 2086
www.dowagrosciences.com.au

CUSTOMER SERVICE TOLL FREE 1-800 700 096

®™ Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow



Dow AgroSciences

DIRECTIONS FOR USE

RESTRAINTS:

Avoid spraying if rain is likely within 6 hours or if strong winds prevail.

DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

1. FIELD CROPS

SITUATION & CROP	WEEDS CONTROLLED	STATE	RATE (per ha)	WHP (days)	CRITICAL COMMENTS
Wheat, Barley	Refer to Weed Table	Vic only	210-800 mL	7	<p>CROP STAGES: ALL CEREALS</p> <p>WA, Vic only: Apply at tillered to boot stages.</p> <p>NSW only: Apply after when the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller.</p> <p>Qld only: Apply from mid-tillering (5 to 6 fully emerged main stem leaves plus one or more tillers) to before boot stage (visible swelling on the head at the top of the main stem).</p> <p>SA, Tas only: Apply from completion of tillering to early jointing stage.</p>
		QLD, NSW only	405-800 mL		
		SA only	230-800 mL		
		Tas only	620-800 mL		
		WA only	800 mL		
Triticale	Refer to Weed Table	QLD, NSW, SA only	405-800 mL	7	<p>CROP STAGES: ALL CEREALS</p> <p>WA, Vic only: Apply at tillered to boot stages.</p> <p>NSW only: Apply after when the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller.</p> <p>Qld only: Apply from mid-tillering (5 to 6 fully emerged main stem leaves plus one or more tillers) to before boot stage (visible swelling on the head at the top of the main stem).</p> <p>SA, Tas only: Apply from completion of tillering to early jointing stage.</p>
Vic only		210-800 mL			
Cereal Rye	Refer to Weed Table	NSW, QLD, only	405-800 mL	7	<p>CROP STAGES: ALL CEREALS</p> <p>WA, Vic only: Apply at tillered to boot stages.</p> <p>NSW only: Apply after when the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller.</p> <p>Qld only: Apply from mid-tillering (5 to 6 fully emerged main stem leaves plus one or more tillers) to before boot stage (visible swelling on the head at the top of the main stem).</p> <p>SA, Tas only: Apply from completion of tillering to early jointing stage.</p>
		Vic only	210-800 mL		
Sugar Cane	Refer to Weed Table	QLD only	1.15-2.38 L	7	Post-emergence
Stubble/Fallow Spray prior to Direct Drilling or Sowing Winter cereals, Grain legumes (Peanuts - Qld only), Canola	Refer to Weed Table	All States	210-800 mL	N/A	<p>Observe the plantback periods given in the table in this leaflet. Must be tank mixed with a knockdown herbicide such as glyphosate, paraquat or paraquat/diquat (e.g. Spray Seed®). Select appropriate rate from the Weed Table. For skeleton weed, spraying should only be done 6-8 weeks before anticipated sowing date and subsequent cultivation limited to a minimum.</p>
Harvest Aid or Salvage Spray	Broadleaf Weeds	All States	1.68 L	7	<p>Apply after dough stage of crop. Interval between application and effectiveness is 10-20 days. For desiccation of green matter, estimate harvest date and spray approximately 14 days earlier. Rain between spraying and actual harvest can negate results.</p> <p>NB. Where thistles are tall and branching above crop, spraying can turn the branches down into the crop, presenting more stalks to cause header comb blockages.</p> <p>Spraying may increase seed contamination of harvest by accelerating maturity.</p> <p>Do not use with undersown legumes that have not set seed.</p>
Winter Cereals	Refer to Weed Table				



1. FIELD CROPS (continued)

SITUATION & CROP	WEEDS CONTROLLED	STATE	RATE (per ha)	WHP (days)	CRITICAL COMMENTS
Potatoes Pre-harvest Preparation	Broadleaf weeds such as Clover, Variegated thistle & Cruciferous weeds	Vic, Tas only	1.15-2.38 L	N/A	Apply approximately 4 to 5 weeks before harvest after the potato haulms have dried off. Use the highest rate where weeds are more than 30 cm in height. For boom spraying apply at least 100 L of spray mixture per hectare. If grasses such as rye grass and winter grass are also present add amitrole.

2. PASTURES, NON-AGRICULTURAL, INDUSTRIAL

SITUATION & CROP	WEEDS CONTROLLED	STATE	RATE (per ha)	WHP (days)	CRITICAL COMMENTS
Improved Pasture containing Clover	Refer to Weed Table	Qld, NSW, SA, Tas only	405-800 mL	7	Clover must be well covered by the grass or extensive damage may result.
Pastures - non legumes, Rights-of-Way, Industrial		Qld, NSW, SA, Tas, WA only	800 mL-4.68 L		Control of most perennial weeds, but due to the rooting habits of most species control may take a number of years. Damage may result to legumes in pasture.
		Vic only	800 mL-4.68 L 70-620 mL		Boom spray. Spot spraying.
Pasture - Direct Drilling or Surface Sowing	Charlock, Clover, Medics, Mustards, Paterson's Curse, Saffron, Slender, Variegated and Spear Thistles, Turnip Weed, Wild Radish, Wild Turnip	NSW only	800 mL-1.5 L (Aerial application)		Apply to young, actively growing weeds. SOWING: Do not sow pasture seeds for at least 21 days after application. If soil moisture is dry, delay sowing for at least 30 days.
	As above plus: Capeweed, Wireweed, Storksbill/ Erodium, Flatweed, Horehound (seedlings), Skeleton Weed, Nodding or Star Thistles		800 mL-1.15 L (Ground application)		
	St. John's Wort		3.26-4.68 L (Aerial or Ground)		
	All of the above plus grasses		As above plus glyphosate		



3. SPOT SPRAYING

SITUATION & CROP	WEEDS CONTROLLED	STATE	RATE (per ha)	WHP (days)	CRITICAL COMMENTS
Spot spraying (All situations)	Refer to Weed Table	All States	1/100th of rate on Weed Table per 10 L water per 100 m ²	7	Apply with a knapsack. Thorough wetting of weed is essential.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION. IN TASMANIA, THIS PRODUCT MAY ONLY BE USED FROM 15TH APRIL TO 15TH SEPTEMBER UNLESS OTHERWISE PERMITTED BY THE REGISTRAR OF PESTICIDES.

WITHHOLDING PERIOD: Pasture, Cereal Crops - **DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION**

HARVEST WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED

WEED TABLE:

NOTE: Where weeds are to be sprayed in a CROP or PASTURE, use only the rates given for the crop in the table below. In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed, e.g. *Rumex* spp. (docks) and *Polygonum* spp. (wireweed, climbing buckwheat) are killed to ground level only.

WEEDS CONTROLLED	APPLICATION RATE (per Hectare)								CRITICAL COMMENTS
	CROP						PASTURE – NON LEGUMES		
	VIC	SA	Tas	NSW	QLD	WA	VIC	QLD, NSW, SA, Tas, WA only	
<i>Amaranthus</i> spp.	-	-	-	800 mL	-	-	-	-	
Angled Onion	-	-	-	-	-	-	3.26 L	800 mL-1.68 L	Spray when buds forming or early flowering.
Apple of Sodom	-	-	-	-	-	-	-	2.9-3.26 L	
Bathurst Burr	-	-	-	800 mL	-	-	1.68-3.26 L	1.68-3.26 L	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Black Knapweed	-	-	-	-	-	-	3.26 L	-	Spray before flowering. DO NOT cultivate these infestations.
Buffalo Burr	-	-	-	-	-	-	-	800 mL-1.15 L (not QLD & WA)	Spray from seedling to pre-flowering. Use higher rate as plant matures.
California Burr	-	-	-	800 mL	-	-	1.68-3.26 L	-	
Caltrop	-	-	-	620-800 mL	-	-	1.68-3.26 L	-	
Cape Tulip	-	-	-	-	-	1.15 L	3.26 L	1.68-3.26 L	Spray before flowering.
Capeweed	800 mL	800 mL	800 mL	530-800 mL	-	-	-	2.47-3.26 L	Spray up to rosette stage.
Charlock	405 mL	405 mL	800 mL	405 mL	-	-	-	800 mL	
Clover	-	-	-	620-800 mL	-	-	-	800 mL	
Colocynth	-	-	-	-	-	-	3.26 L	-	Spray at seedling stage only.



WEED TABLE (continued):

WEEDS CONTROLLED	APPLICATION RATE (per Hectare)								CRITICAL COMMENTS
	CROP						PASTURE – NON LEGUMES		
	VIC	SA	Tas	NSW	QLD	WA	VIC	QLD, NSW, SA, Tas, WA only	
Deadnettle	-	-	-	800 mL	-	-	-	-	
Devil's Claw	-	-	-	800 mL	-	-	1.32 L	-	Spray prior to pods forming.
Dock	800 mL	800 mL	-	-	800 mL	800 mL	-	1.68-2.47 L	Spray at rosette stage to kill top growth only.
Fat Hen	-	-	-	405-800 mL	-	-	-	-	
Flatweed	-	-	-	800 mL	-	-	-	2.47-3.26 L	
Fumitory - red	-	800 mL	-	800 mL	-	-	-	2.47-3.26 L	Spray up to rosette stage.
Fumitory - white	800 mL	405 mL	-	800 mL	-	-	-	2.47-3.26 L	Spray up to rosette stage.
Galvanized Burr	-	-	-	-	-	-	4.68 L	4.68 L (not Qld & WA)	Spray from seedling to pre-flowering.
Goosefoots	-	-	-	800 mL	-	-	-	-	
Hard Head or Russian Knapweed	-	-	-	-	-	-	3.48-5.2 L	-	Spray before flowering.
Hoary Cress, Whiteweed	-	800 mL	800 mL	800 mL	-	-	1.68-3.26 L	1.68-2.12 L	Spray from late rosette to pre-flowering.
Hogweed / Wireweed	800 mL	800 mL	-	800 mL	800 mL	-	-	1.15-1.68 L (not SA)	Spray up to rosette stage.
Horehound (seedlings)	-	800 mL	-	-	-	840 mL	-	1.68-3.26 L	Late autumn to early spring.
Iron Weed, Corn Gromwell	-	-	-	800 mL	-	-	-	1.15-1.68 L	
Khaki Weed	-	-	-	-	-	-	-	800 mL-1.15 L (not SA)	Spray young seedlings only.
Lincoln Weed	-	800 mL	-	-	-	-	-	-	Autumn spray before sowing improves control.
London Rocket	-	-	-	-	-	575 mL	-	1.59-2.47 L (WA only)	
Lupins	800 mL	-	-	405-800 mL	-	-	-	-	Spray up to rosette stage.
Mellilotus / Hexham Scent	800 mL	800 mL	-	-	800 mL	-	-	1.15-1.68 L	Spray up to rosette stage.
Melons-Camel, Paddy	-	-	-	405-800 mL	-	-	-	-	
Mustards	330 mL	230-800 mL	800 mL	405-800 mL	620 mL	620 mL	3.26 L	1.68-2.47 L	Spray up to rosette stage.



WEED TABLE (continued):

WEEDS CONTROLLED	APPLICATION RATE (per Hectare)								CRITICAL COMMENTS
	CROP						PASTURE – NON LEGUMES		
	VIC	SA	Tas	NSW	QLD	WA	VIC	QLD, NSW, SA, Tas, WA only	
Mexican Poppy	-	-	-	800 mL	-	840 mL	-	800 mL-1.15 L (1.15-1.5 L WA only)	Spray rosette stage and before flowering.
Mintweed	-	-	-	800 mL	620 mL	-	-	800 mL-1.15 L	Spray active seedlings only.
Muskweed	800 mL	-	-	-	-	-	-	-	Spray up to rosette stage.
New Zealand Spinach	-	-	-	800 mL	-	-	-	-	
Noogoora Burr	-	-	-	800 mL	-	-	1.68-3.26 L	1.68-3.26 L	Spray seedling to pre-flowering.
Nut Grass	-	-	-	-	-	-	3.26 -5.2 L	-	Spray within 4 weeks of foliage emergence, repeat spray necessary.
Paterson's Curse	-	-	-	800 mL	-	840 mL	1.68-3.26 L	800 mL-1.68 L (1.15-1.5 L WA only)	Spray seedling to rosette stage.
Poppy Wild	405 mL	-	-	-	-	-	-	2.12-3.3 L	Spray up to rosette stage.
Ragwort	-	-	-	-	-	-	3.26 L	3.26 L	Spray at rosette to cabbage stage.
Rapeseed	800 mL	-	-	405-800 mL	-	-	-	-	Spray up to rosette stage.
Rapistrum spp.	-	-	-	-	-	650 mL	-	840 mL (WA only)	
Rough Poppy	-	405 mL	-	405-800 mL	-	-	-	800 mL	Spray young seedlings only.
St. John's Wort	-	-	-	-	-	-	3.26-5.9 L	3.26-4.68 L	Spray before flowering. Spray before plants 40cm high.
Safflower	-	-	-	405-800 mL	-	-	-	-	
Sand Mustard / Sand Rocket	-	-	-	-	-	-	3.26 L	-	Spray before flowering.
Shepherds Purse	-	-	-	800 mL	-	-	-	-	
Silverleaf Nightshade	-	-	-	-	-	-	3.26 L	-	Spray at flowering. Fallow land: controls top growth only.
Skeleton Weed	-	800 mL	-	800 mL	-	-	3.26 L	1.15-1.68 L	Spray rosettes before aerial growth commences.
Stingless Nettle (Deadnettle)	-	800 mL	-	-	-	-	-	2.12-2.47 L	
Stinging Nettle	800 mL	-	-	-	-	-	-	-	Spray up to rosette stage.



WEED TABLE (continued):

WEEDS CONTROLLED	APPLICATION RATE (per Hectare)								CRITICAL COMMENTS
	CROP						PASTURE – NON LEGUMES		
	VIC	SA	Tas	NSW	QLD	WA	VIC	QLD, NSW, SA, Tas, WA only	
Stinkwort	-	-	-	800 mL	-	-	1.68-3.26 L	1.68-3.26 L	Spray younger plants, use higher rate as plants mature.
Storksbill / Erodium	-	-	-	800 mL	-	-	-	-	
Sunflower seedlings	800 mL	-	-	405-800 mL	620 mL	-	-	-	Spray multiple leaves.
Thistle: - Golden	-	-	-	-	-	-	3.26 L	3.26 L	Spray up to rosette stage.
- Nodding	-	-	-	-	-	-	3.26 L	1.15-1.68 L	Spray rosette to pre-flowering.
Thistle: - Saffron	620 mL	800 mL	-	405-800 mL	800 mL	800 mL	800 mL-1.68 L	800 mL-2.47 L	Spray up to rosette stage.
- Sheep	-	-	-	-	-	840 mL	-	840 mL-3.26 L (WA only)	
- Slender / Shore	-	-	800 mL	800 mL	-	-	1.68-3.26 L	800 mL-3.26 L	Spray at rosette stage.
- Soldier	-	-	-	-	-	-	3.26 L	-	Spray at rosette stage.
- Spear	800 mL	-	800 mL	-	-	-	0.8-2.47 L	1.15-2.12 L	Spray at seedling to rosette stage. Use higher rate as plants mature (pastures).
- Stemless	-	-	-	-	-	-	3.26 L	2.47-3.26 L	Spray rosette stage to flowering.
- St Barnaby's	-	-	-	-	-	-	-	1.15-1.68 L	
- Star	-	-	-	800 mL	-	-	1.68-3.26 L	-	Spray seedling to rosette stage. Use higher rate as plants mature.
- Variegated	-	-	800 mL	405-800 mL	620 mL	-	800 mL-2.47 L	800 mL-3.26 L	Spray at rosette stage. Can cause stock poisoning.
Thornapple	-	-	-	405-800 mL	-	-	3.26 L	800 mL-1.68 L	Spray at seedling stage.
Tree Hogweed	800 mL	-	-	-	-	-	-	-	Spray up to rosette stage.
Turnip Weed	-	405 mL	-	405-800 mL	405 mL	620 mL	-	800 mL	Spray seedlings only.
Vetches/Tares	800 mL	620 mL	800 mL	-	-	-	-	-	
Wards Weed	-	405 mL	-	-	-	-	-	-	Spray at seedling stage.
Wild Cabbage	800 mL	-	-	-	-	-	-	-	Spray up to rosette stage.



WEED TABLE (continued):

WEEDS CONTROLLED	APPLICATION RATE (per Hectare)								CRITICAL COMMENTS
	CROP						PASTURE – NON LEGUMES		
	VIC	SA	Tas	NSW	QLD	WA	VIC	QLD, NSW, SA, Tas, WA only	
Wild Garlic	-	-	-	-	-	-	6.62 L	-	Suppresses aerial growth only.
Wild Mignonette	-	-	-	-	-	840 mL	3.26 L	-	Spray at rosette stage.
Wild Mustard	-	-	-	-	-	650 mL	-	1.8-2.47 L (WA only)	
Wild Radish	800 mL	800 mL	800 mL	405-800 mL	800 mL	650 mL	-	800 mL (840 mL WA only)	Spray up to rosette stage.
Wild Sage	-	-	-	-	-	-	-	2.47-3.26 L	
Wild Teasel	-	-	-	-	-	-	1.68-3.26 L	-	Spray at rosette stage. Use higher rate as plants mature.
Wild Turnip	210 mL	230 mL	800 mL	405-800 mL	-	450 mL	-	800 mL (840 mL WA only)	Spray up to rosette stage.

Plantback Periods (days) for Esteron LV

CROP	RATES		
	Up to 510 mL/ha	510 mL - 1.15 L/ha	1.15 L - 1.59 L/ha
Balansa Clover	7	7	10
Barley ①	1	1	3
Chickpeas ②	7	14	21
Cotton	10	14	21
Faba Beans	7	7	10
Field Peas	7	14	14
Lentils	7	7	10
Linseed	7	7	14
Lucerne	7	7	10
Lupins ③	7	14	21
Medics	7	7	10
Narbon Beans	7	7	10
Navy Bean	10	10	14
Oats	3	3	7
Perennial Ryegrass	7	7	10
Persian Clover	7	7	10



Plantback Periods (days) for Esteron LV (continued)

CROP	RATES		
	Up to 510 mL/ha	510 mL - 1.15 L/ha	1.15 L - 1.59 L/ha
Phalaris	7	7	10
Canola / Rapeseed ②	14	21	28
Rice	7	7	14
Safflower ②	7	14	21
Sorghum ③	3	7	10
Soybean	14	14	21
Sub-Clover	7	7	10
Sunflower ③	7	10	14
Triticale ①	1	3	7
Vetch	7	7	10
Wheat ①	1	3	7
White Clover	7	7	10

IMPORTANT: WHEN APPLIED TO DRY SOILS AT LEAST 15 mm OF RAIN MUST FALL PRIOR TO THE COMMENCEMENT OF THE PLANT BACK PERIOD.

NOTES:

- ① In Queensland, no rainfall is required to fall prior to commencement of Plantback Period for barley, triticale and wheat.
- ② In Queensland, planting of canola/rapeseed, chickpeas and safflower must be delayed for at least 14 days following rainfall of at least 15 mm.
- ③ In Central Queensland, when using 730 mL/ha or less of Dow AgroSciences Esteron LV, the Plantback Period for sorghum and sunflower is 1 day irrespective of rainfall.
- ④ In WA the Plantback Period for lupins at all rates is 28 days.

GENERAL INSTRUCTIONS

Before opening, carefully read DIRECTIONS FOR USE, PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS, SAFETY DIRECTIONS and FIRST AID Instructions.

DO NOT spray in high winds.

DO NOT spray crops or weeds outside the stages indicated in the CRITICAL COMMENTS as damage, loss of yield or inadequate weed control may result.

APPLICATION INFORMATION

This product may be used in either high or low volume sprays. Just pour into water and stir.

Boom Spraying - Use 30-120 litres water/ha.

Aerial Spraying - Use 10-90 litres water/ha.

Note: Refer to the Department of Agriculture / Primary Industries in your state for the current restricted spraying areas.

EQUIPMENT MAINTENANCE AND USAGE

Keep the spray unit for herbicides only if possible.

Otherwise wash out the unit with hot soapy water followed by several clear water rinses. **DO NOT** use wooden spray vats as they cannot be cleaned. Hoses cannot be cleaned and new hoses should be fitted when the unit is to be used for any other purpose.

COMPATIBILITY

This product can be tank mixed with atrazine, dicamba, chlorsulfuron, Garlon™ 600, glyphosate, metsulfuron, paraquat, paraquat/diquat (e.g. SpraySeed®), Starane™ Advanced and triasulfuron.

NOTE:

1. As formulations of other manufacturers' products are beyond the control of Dow AgroSciences all mixtures should be tested on a small scale before mixing in the spray tank.
2. Tank mixing instructions:
Fill the spray tank 1/4 full of water and agitate. Add wettable powders and water dispersible granules first. Agitate until these are uniformly dispersed, meanwhile adding water until the tank is 90% full. Add suspension concentrates (flowables) then soluble concentrates. Emulsifiable concentrates go in last. Top off the tank with water and continue agitation until all the ingredients are properly mixed.

Observe any mixing sequence instructions mentioned on the tank mix products.

RESISTANT WEEDS WARNING

GROUP I HERBICIDE

Dow AgroSciences Esteron LV Herbicide (Esteron LV) is a member of the phenoxy group of herbicides. Esteron LV has the disruptors of plant cell growth mode of action. For weed resistance management Esteron LV is a Group I herbicide. Some naturally occurring weed biotypes resistant to Esteron LV and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Esteron LV or Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Dow AgroSciences Australia Limited accepts no liability for any losses that may result from the failure of Esteron LV to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Legume Tolerance: If clovers are present, care should be taken to ensure that they have reached the 3-4 leaf stage before spraying. Rates above 405 mL of this product per hectare will destroy most clovers, whilst lucerne and medics are susceptible at any strength.

Drift Warning: DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray drift onto nearby susceptible plants, adjacent crops, crop lands or pastures. In particular, avoid spray drift and vapour movement onto susceptible crops such as: cotton, tobacco, tomatoes, vines, fruit trees, ornamentals, oil seed and legume crops and other susceptible plants and trees (e.g. Kurrajongs, Belahs, Eucalypts).

- **DO NOT** use unless wind speed is more than 3 kilometres per hour and less than 15 kilometres per hour as measured at the application site.
- **DO NOT** apply with smaller than coarse to very coarse spray droplets according to the ASABE S572 definition for standard nozzles.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied on any plants at any time as recommended in the DIRECTIONS FOR USE.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used container.



STORAGE AND DISPOSAL

Storage of all containers:

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.

DO NOT store near food, feedstuffs, fertilisers or seed.

This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate by adding to the spray tank. Do not dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management site. The cap should not be replaced but may be taken separately. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

SPILL AND LEAK MANAGEMENT

Do not touch or walk through spilled material. Wear a face shield or goggles, overalls buttoned to neck and wrist, chemical resistant gloves and footwear. Stop leak when safe to do so. Dam area and prevent entry into waterways, and drains.

Small spills/leaks: Absorb with material such as sand, soil or sawdust. Collect spilled product and place in sealable container for disposal. Spill residues may be cleaned using water and detergent. Contain and absorb wash water for disposal. Absorb and collect washings and place in the same sealable container for disposal. Dam the area of large spills and report them to Dow AgroSciences Emergency Services at 1-800 033 882.

SAFETY DIRECTIONS

- Poisonous if swallowed.
- Avoid contact with eyes and skin.
- DO NOT inhale spray mist.
- When preparing spray, wear PVC or rubber apron, elbow length PVC gloves and face shield. When using the prepared spray, wear face shield.
- If product on skin, immediately wash area with soap and water.
- After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water.
- After each day's use wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone: *Australia* 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed on the Material Safety Data Sheet for Dow AgroSciences **ESTERON™ LV HERBICIDE** which is available from Dow AgroSciences on request.

Call Customer Service Toll Free on 1-800 700 096 or visit www.dowagrosciences.com.au

©™ Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

APVMA Approval No: 60216/0214

EMERGENCY RESPONSE

(ALL HOURS)

RING FROM ANYWHERE IN
AUSTRALIA
1800 033 882
(LOCAL CALL FEE ONLY)

IN A TRANSPORT
EMERGENCY ONLY
DIAL 000
FOR POLICE OR
FIRE BRIGADE



Dow AgroSciences