

CAUTION

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

Genfarm LV Ester 600 Herbicide

Active Constituent: 600 g/L 2,4-D present as the ethylhexyl ester

GROUP	I	HERBICIDE
--------------	----------	------------------

A low volatile herbicide for selective control of various weeds in crops, pastures and non-agricultural areas according to the Directions for Use

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE.

This is a PHENOXY HERBICIDE that can cause severe damage to susceptible crops. Please read the Drift Warning Statement.

Contents: 20 L, 100L, 110L, 1000L

Genfarm Crop Protection Pty Ltd

Suite 2, Level 3, 64 Talavera Road,

Macquarie Park NSW, 2113

Tel: (02) 9889 5400

BACK PANEL

STORAGE AND DISPOSAL

Storage of all containers:

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.

DO NOT store near food, feedstuffs, fertilisers or seed.

Disposal (20L, 100L) This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or preferably pressure rinse container for disposal. Dispose of rinsate by adding to the spray tank. Do not dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap and the closed system valve present in 100L containers removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management site. The cap should not be replaced but may be taken separately. The closed system valve should be disposed of in a local authority landfill.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Disposal (110L)

Do not tamper with the dry valves or security seal. Do not contaminate the drum with water or any other foreign matter. After each use of the product ensure that the dry valve coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained. Add the rinsings to the spray tank. When the drum is empty close all valves and return to the point of purchase. The drum remains the property of Genfarm and must be returned.

1000L - Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray, wear PVC or rubber apron, elbow length PVC gloves and face shield. When using the prepared spray, wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. (Phone: Australia 13 11 26).

MATERIAL SAFETY DATA SHEET

For further information, refer to the Material Safety Data Sheet (MSDS) which is available from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Genfarm Crop Protection Pty Ltd. accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd. will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

APVMA Approval No: 62347/20L/1107, 62347/100L/1107, 62347/110L/1107, 62347/1000L/1107

Batch No:

Date of Manufacture:

In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS AUSTRALIA WIDE
--	---



(20L, 100L labels)

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Genfarm LV Ester 600 Herbicide

Active Constituent: 600 g/L 2,4-D present as the ethylhexyl ester

GROUP	I	HERBICIDE
-------	----------	-----------

A low volatile herbicide for selective control of various weeds in crops, pastures and non-agricultural areas according to the Directions for Use

IMPORTANT: READ THIS BOOKLET BEFORE USE.

This is a PHENOXY HERBICIDE that can cause severe damage to susceptible crops. Please read the Drift Warning Statement.

APVMA APPROVAL NO: 62347/1107

Genfarm Crop Protection Pty Ltd
Suite 2, Level 3, 64 Talavera Road,
Macquarie Park NSW, 2113
Tel: (02) 9889 5400

DIRECTIONS FOR USE

RESTRAINTS: Avoid spraying if rain is likely within 6 hours or if strong winds prevail.
DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

1. FIELD CROPS

Situation & Crop	Weeds Controlled	State	Rate (per ha)	WHP (days)	Critical Comments
Wheat, Barley	Refer to Weed Table	Vic only	240-900 mL	7 (G)	<p>CROP STAGES: ALL CEREALS Variations between varieties do occur. Check sensitivity and growth stages of varieties before applying. Damage may result if applied too early.</p> <p>Vic only: Apply at tillered to boot stages. ACT, NSW only: Apply after when the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller.</p> <p>Qld only: Apply from mid-tillering (5 to 6 fully emerged main stem leaves plus one or more tillers) to before boot stage (visible swelling on the head at the top of the main stem).</p> <p>SA, Tas only: Apply from completion of tillering to early jointing stage.</p> <p>WA only: Apply from the 5 leaf stage up to jointing stage (Zadoks 15-33).</p> <p>Apply only at 6 leaf stage for Cranbrook and Jacup wheats (Zadoks 16) to avoid possible damage.</p>
		Qld, NSW, ACT only	460-900 mL		
		SA only	260-900 mL		
		Tas only	700-900 mL		
		WA only	900 mL		
Triticale	Refer to Weed Table	Qld, NSW, ACT	460-900 mL	7 (G)	<p>Qld only: Apply from mid-tillering (5 to 6 fully emerged main stem leaves plus one or more tillers) to before boot stage (visible swelling on the head at the top of the main stem).</p> <p>SA, Tas only: Apply from completion of tillering to early jointing stage.</p> <p>WA only: Apply from the 5 leaf stage up to jointing stage (Zadoks 15-33).</p> <p>Apply only at 6 leaf stage for Cranbrook and Jacup wheats (Zadoks 16) to avoid possible damage.</p>
		SA only	270-930 mL		
		Vic only	240-900 mL		
Cereal Rye	Refer to Weed Table	ACT, NSW, Qld, only	460-900 mL	7 (G)	<p>Qld only: Apply from mid-tillering (5 to 6 fully emerged main stem leaves plus one or more tillers) to before boot stage (visible swelling on the head at the top of the main stem).</p> <p>SA, Tas only: Apply from completion of tillering to early jointing stage.</p> <p>WA only: Apply from the 5 leaf stage up to jointing stage (Zadoks 15-33).</p> <p>Apply only at 6 leaf stage for Cranbrook and Jacup wheats (Zadoks 16) to avoid possible damage.</p>
		Vic only	240-900 mL		
Sugar Cane	Refer to Weed Table	Qld only	1.3-2.7 L		Post-emergence
Stubble/Fallow Spray prior to Direct Drilling or Sowing Winter cereals, Grain legumes (Peanuts - Qld only), Canola		All States	240-900 mL	N/A	<p>Observe the plant back periods given in the table in this leaflet. Must be tank mixed with a knockdown herbicide such as glyphosate, paraquat or paraquat/diquat (e.g. Genfarm Di-Par 250). Select appropriate rate from the Weed Table.</p> <p>For skeleton weed, spraying should only be done 6-8 weeks before anticipated sowing date and subsequent cultivation limited to a minimum.</p>
Harvest Aid or Salvage Spray	Broadleaf Weeds	All States	1.9 L	7 (G)	<p>Apply after dough stage of crop. Interval between application and effectiveness is 10-20 days. For desiccation of green matter, estimate harvest date and spray approximately 14 days earlier. Rain between spraying and actual harvest can negate results.</p> <p>NB. Where thistles are tall and branching above crop, spraying can turn the branches down into the crop, presenting more stalks to cause header comb blockages.</p> <p>Spraying may increase seed contamination of harvest by accelerating maturity.</p> <p>Do not use with undersown legumes that have not set seed.</p>
Winter Cereals	Refer to Weed Table				
Potatoes Pre-harvest Preparation	Broadleaf weeds such as Clover, Variegated thistle & Cruciferous weeds	Vic, Tas only	1.3-2.7 L	N/A	<p>Apply approximately 4 to 5 weeks before harvest after the potato haulms have dried off. Use the highest rate where weeds are more than 30 cm in height.</p> <p>For boom spraying apply at least 100 L of spray mixture per hectare. If grasses such as rye grass and winter grass are also present add amitrole.</p>

2. PASTURES, NON-AGRICULTURAL, INDUSTRIAL

Situation & Crop	Weeds Controlled	State	Rate (per ha)	WHP (days)	Critical Comments
Improved Pasture containing Clover	Refer to Weed Table	Qld, NSW, ACT, SA, Tas only	460-900 mL	7 (G)	Clover must be well covered by the grass or extensive damage may result.
Pastures – non legumes, Rights-of-Way, Industrial		Qld, NSW, ACT, SA, Tas, WA only	900 mL-5.3 L	7 (G)	Control of most perennial weeds, but due to the rooting habits of most species control may take a number of years. Damage may result to legumes in pasture.
		Vic only	900 mL-7.5 L		Boom spray.
Pasture - Direct Drilling or Surface Sowing	Charlock, Clover, Medics, Mustards, Paterson's Curse, Saffron, Slender, Variegated and Spear Thistles, Turnip Weed, Wild Radish, Wild Turnip	ACT, NSW, Qld, WA, SA, Vic, Tas only	900 mL-1.7 L (Aerial application)	NA	Apply to young, actively growing weeds. SOWING: Do not sow pasture seeds for at least 21 days after application. If soil moisture is dry, delay sowing for at least 30 days.
	As above plus: Capeweed, Wireweed, Storksbill/Erodium, Flatweed, Horehound (seedlings), Skeleton Weed, Nodding or Star Thistles		900 mL-1.3 L (Ground application)		
	St. John's Wort		3.7-5.3 L (Aerial or Ground)		
	All of the above plus grasses		As above plus glyphosate		

3. SPOT SPRAYING

Situation & Crop	Weeds Controlled	State	Rate	WHP (days)	Critical Comments
Spot spraying (All situations)	Refer to Weed Table	All States	1/100th of rate on Weed Table per 10 L water per 100 m ²	7 (G)	Apply with a knapsack. Thorough wetting of weed is essential.

Note: Grazing (G), Harvesting (H).

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

IN TASMANIA, THIS PRODUCT MAY ONLY BE USED FROM 15TH APRIL TO 15TH SEPTEMBER UNLESS OTHERWISE PERMITTED BY THE REGISTRAR OF PESTICIDES.

WITHHOLDING PERIOD: PASTURE, CEREAL CROPS - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

HARVEST WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED

WEED TABLE:

NOTE: Where weeds are to be sprayed in a CROP or PASTURE, use only the rates given for the crop in the table below. In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed, e.g. *Rumex* spp. (docks) and *Polygonum* spp. (wireweed, climbing buckwheat) are killed to ground level only.

Weeds Controlled	Application Rate (per Hectare)								Critical Comments
	CROP						PASTURE – NON-LEGUMES		
	Vic	SA	Tas	NSW, ACT	QLD	WA	Vic	QLD, NSW, ACT, SA, Tas, WA only	
Amaranthus spp.	-	-	-	900 mL	-	-	-	-	
Angled Onion	-	-	-	-	-	-	3.7 L	900 mL-1.9 L	Spray when buds forming or early flowering.
Apple of Sodom	-	-	-	-	-	-	-	3.3-3.7 L	
Bathurst Burr	-	-	-	900 mL	-	-	1.9-3.7 L	1.9-3.7 L	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Black Knapweed	-	-	-	-	-	-	3.7 L	-	Spray before flowering. DO NOT cultivate these infestations.
Buffalo Burr	-	-	-	-	-	-	-	900 mL-1.3 L (not Qld & WA)	Spray from seedling to pre-flowering. Use higher rate as plant matures.
California Burr	-	-	-	900 mL	-	-	1.9-3.7L	1.3-1.9 L (Not SA)	
Caltrop	-	-	-	700-900 mL	-	-	1.7-3.7 L	-	
Cape Tulip	-	-	-	-	-	1.3 L	3.7 L	1.9-3.7 L	Spray before flowering.
Capeweed	900 mL	900 mL	900 mL	600-900 mL	-	-	-	2.8-3.7 L	Spray up to rosette stage.
Charlock	460 mL	460 mL	900 mL	460 mL	-	-	-	900 mL	
Clover	-	-	-	700-900 mL	-	-	-	-	
Colocynth	-	-	-	-	-	-	3.7 L	-	Spray at seedling stage only.
Deadnettle	-	-	-	900 mL	-	-	-	-	
Devil's Claw	-	-	-	900 mL	-	-	1.5 L	-	Spray prior to pods forming.
Dock	900 mL	900 mL	-	-	900 mL	900 mL	-	1.9-2.8 L	Spray at rosette stage to kill top growth only.
Fat Hen	-	-	-	460-900 mL	-	-	-	-	
Flatweed	-	-	-	900 mL	-	-	-	2.8-3.7 L	
Fumitory – red	-	900 mL	-	900 mL	-	-	-	2.8-3.7 L	Spray up to rosette stage.
Fumitory – white	900 mL	460mL	-	900 mL	-	-	-	2.8-3.7 L	Spray up to rosette stage.
Galvanized Burr	-	-	-	-	-	-	5.3 L	5.3 L (not Qld & WA)	Spray from seedling to pre-flowering.
Goosefoots	-	-	-	900 mL	-	-	-	-	
Hard Head or Russian Knapweed	-	-	-	-	-	-	3.7-5.9 L	-	Spray before flowering.

Hoary Cress, Whiteweed	-	900 mL	900 mL	900 mL	-	-	1.9-3.7 L	1.9-2.2 L	Spray from late rosette to pre-flowering.
Hogweed / Wireweed	900 mL	900 mL	-	900 mL	900 mL	-	-	1.3-1.9 L (not SA)	Spray up to rosette stage.
Horehound (seedlings)	-	900 mL	-	-	-	950 mL	-	1.9-3.7 L	Late Autumn to early Spring.
Iron Weed, Corn Gromwell	-	-	-	900 mL	-	-	-	1.3-1.9 L	
Khaki Weed	-	-	-	-	-	-	-	900 mL- 1.3 L (not SA)	Spray young seedlings only.
Lincoln Weed	-	900 mL	-	-	-	-	-	-	Autumn spray before sowing improves control.
London Rocket			-	-	-	650 mL	-	1.8-2.8 L (WA only)	
Lupins	900 mL	-	-	460- 900 mL	-	-	-	-	Spray up to rosette stage.
Melilotus / Hexham Scent	900 mL	900 mL	-	-	900 mL	-	-	1.3-1.9 L	Spray up to rosette stage.
Melons-Camel, Paddy	-	-	-	460- 800 mL	-	-	-	-	
Mustards	370 mL	260- 900 mL	900 mL	460- 900 mL	700 mL	700 mL	3.7 L	1.9-2.8 L	Spray up to rosette stage.
Mexican Poppy	-	2.6-4 L	-	900 mL	-	950 mL	-	900 mL- 1.3 L (1.3-1.7 L WA only)	Spray rosette stage and before flowering.
Mintweed	-	-	-	900 mL	700 mL	-	-	900 mL- 1.3 L	Spray active seedlings only.
Muskweed	900 mL	-	-	-	-	-	-	-	Spray up to rosette stage.
New Zealand Spinach	-	-	-	900 mL	-	-	-	-	
Noogoora Burr	-	-	-	900 mL	-	-	1.9-3.7 L	1.9-3.7 L	Spray seedling to pre-flowering.
Nut Grass	-	-	-	-	-	-	3.7 -5.9 L	-	Spray within 4 weeks of foliage emergence, repeat spray necessary.
Paterson's Curse	-	-	-	900 mL	-	950 mL	1.9-3.7 L	900 mL- 1.9 L (1.3-1.7 L WA only)	Spray seedling to rosette stage.
Poppy Wild	460 mL	-	-	-	-	-	-	2.4-3.3 L	Spray up to rosette stage.
Ragwort	-	-	-	-	-	-	3.7 L	3.7 L	Spray at rosette to cabbage stage.
Rapeseed	900 mL	-	-	460- 900 mL	-	-	-	-	Spray up to rosette stage.
Rapistrum spp.	-	-	-	-	-	650 mL	-	950 mL (WA only)	
Rough Poppy	-	460 mL	-	460- 900 mL	-	-	-	900 mL	Spray young seedlings only.
St. John's Wort	-	-	-	-	-	-	3.7-5.9 L	3.7-5.3 L	Spray before flowering. Spray before plants 40cm high.
Safflower	-	-	-	460- 900 mL	-	-	-	-	
Sand Mustard / Sand Rocket	-	-	-	-	-	-	3.7 L	-	Spray before flowering.
Shepherds Purse	-	-	-	900 mL	-	-	-	-	

Silverleaf Nightshade	-	-	-	-	-	-	3.7 L	-	Spray at flowering. Fallow land: controls top growth only.
Skeleton Weed	-	900 mL	-	900 mL	-	-	3.7 L	1.3-1.9 L	Spray rosettes before aerial growth commences.
Stingless Nettle (Deadnettle)	-	900 mL	900 mL	-	-	-	-	2.4-2.8 L	
Stinging Nettle	900 mL	-	-	-	-	-	-	-	Spray up to rosette stage.
Stinkwort	-	-	-	900 mL	-	-	1.9-3.7 L	1.9-3.7 L	Spray younger plants, use higher rate as plants mature.
Storksbill / Erodium	-	-	-	900 mL	-	-	-	-	
Sunflower seedlings	900 mL	-	-	460-900 mL	700 mL	-	-	-	Spray multiple leaves.
Thistle:									
- Golden	-	-	-	-	-	-	3.7 L	3.7 L	Spray up to rosette stage.
- Nodding	-	-	-	-	-	-	3.7 L	1.3-1.9 L	Spray rosette to pre-flowering.
- Saffron	700 mL	900 mL	-	460-900 mL	900 mL	900 mL	900 mL-1.9 L	900 mL-2.8 L	Spray up to rosette stage.
- Sheep	-	-	-	-	-	950 mL	-	950 mL-3.7 L (WA only)	
- Slender / Shore	-	-	900 mL	900 mL	-	-	1.9-3.7 L	900-3.7 L	Spray at rosette stage.
Thistles									
- Soldier	-	-	-	-	-	-	3.7 L	-	Spray at rosette stage.
- Spear	900 mL	-	900 mL	-	-	-	900 mL-2.8 L	1.3-2.4 L	Spray at seedling to rosette stage. Use higher rate as plants mature (pastures).
- Stemless	-	-	-	-	-	-	3.7 L	2.8-3.7 L	Spray rosette stage to flowering.
- St Barnaby's	-	-	-	-	-	-	-	1.3-1.9 L	
- Star	-	-	-	900 mL	-	-	1.9-3.7 L	1.3-1.9 L	Spray seedling to rosette stage. Use higher rate as plants mature.
- Variegated	-	-	900 mL	460-900 mL	620 mL	-	900 mL-2.8 L	900 mL-3.7 L	Spray at rosette stage. Can cause stock poisoning.
Thornapple	-	4 L	-	460-900 mL	-	-	3.7 L	900 mL-1.9 L	Spray at seedling stage.
Tree Hogweed	900 mL	-	-	-	-	-	-	-	Spray up to rosette stage.
Turnip Weed	-	460 mL	-	460-900 mL	460 mL	700 mL	-	900 mL	Spray seedlings only.
Vetches/Tares	900 mL	700 mL	900 mL	-	-	-	-	-	
Wards Weed	-	460 mL	-	-	-	-	-	-	Spray at seedling stage.
Wild Cabbage	900 mL	-	-	-	-	-	-	-	Spray up to rosette stage.
Wild Garlic	-	-	-	-	-	-	7.5 L	-	Suppresses aerial growth only.
Wild Mignonette	-	-	-	-	-	950 mL	3.7 L	-	Spray at rosette stage.
Wild Mustard	-	-	-	-	-	650 mL	-	1.8-2.8 L (WA only)	
Wild Radish	900 mL	900 mL	900 mL	460-900 mL	900 mL	650 mL	-	900 mL (950 mL WA only)	Spray up to rosette stage.
Wild Sage	-	-	-	-	-	-	-	2.8-3.7 L	
Wild Teasel	-	-	-	-	-	-	1.9-3.7 L	-	Spray at rosette stage. Use higher rate as plants mature.
Wild Turnip	240 mL	260 mL	900 mL	460-900 mL	-	450 mL	-	900 mL (950 mL WA only)	Spray up to rosette stage.

Plant back Period (days) for Genfarm LV Ester 600

Crop	Rates		
	Up to 580 L/ha	580L to 1.15 L/ha	1.15 to 1.8 L/ha
Balansa Clover	7	7	10
Barley ①	1	1	3
Chickpeas ②	7	14	21
Cotton	10	14	21
Faba Beans	7	7	10
Field Peas	7	14	14
Lentils	7	7	10
Linseed	7	7	14
Lucerne	7	7	10
Lupins ④	7	14	21
Medics	7	7	10
Narbon Beans	7	7	10
Navy Beans	10	10	14
Oats	3	3	7
Perennial Ryegrass	7	7	10
Persian Clover	7	7	10
Phalaris	7	7	10
Canola/Rapeseed ②	14	21	28
Rice	7	7	14
Safflower ②	7	14	21
Sorghum ③	3	7	10
Soybean	14	14	21
Sub - Clover	7	7	10
Sunflower ⑤	7	10	14
Triticale ①	1	3	7
Vetch	7	7	10
Wheat ①	1	3	7
White Clover	7	7	10

IMPORTANT

WHEN APPLIED TO DRY SOILS AT LEAST 15 MM OF RAIN MUST FALL PRIOR TO THE COMMENCEMENT OF THE PLANT BACK PERIOD.

NOTES:

- ① In Queensland, no rainfall is required to fall prior to commencement of Plant back Period for barley triticale, and wheat.
- ② In Queensland, planting of canola / rapeseed, chickpeas and safflower must be delayed for at least 14 days following rainfall of at least 15 mm
- ③ In Central Queensland, when using 830 mL/ha or less of Genfarm LV Ester 600, the Plant back Period for sorghum and sunflower is 1 day irrespective of rainfall.
- ④ In WA the Plant back Period for lupins at all rates is 28 days.

GENERAL INSTRUCTIONS

Before opening, carefully read Directions for Use, PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS, SAFETY DIRECTIONS and FIRST AID Instructions.

DO NOT spray in high winds.

DO NOT spray crops or weeds outside the stages indicated in the CRITICAL COMMENTS as damage, loss of yield or inadequate weed control may result.

APPLICATION INFORMATION

This product may be used in either high or low volume sprays. Just pour into water and stir.

Boom Spraying - Use 30-100 litres water/ha.

Aerial Spraying - Use 10-90 litres water/ha.

Note: Refer to the Department of Agriculture / Primary Industries in your state for the current restricted spraying areas.

EQUIPMENT MAINTENANCE AND USAGE

Keep the spray unit for herbicides only if possible. Otherwise wash out the unit with hot soapy water followed by several clear water rinses. DO NOT use wooden spray vats as they cannot be cleaned. Hoses cannot be cleaned and new hoses should be fitted when the unit is to be used for any other purpose.

COMPATIBILITY

This product can be tank mixed with atrazine, dicamba, Genfarm Huron 750WG, Genfarm Trident 600, glyphosate, Genfarm Muron 600, paraquat, paraquat/diquat (e.g. Genfarm Di-Par), Genfarm Fluroxypyr 200 and triasulfuron.

NOTE:

1. As formulations of other manufacturers' products are beyond the control of Genfarm all mixtures should be tested on a small scale before mixing in the spray tank.
2. Tank mixing instructions:

Fill the spray tank 1/4 full of water and agitate. Add wettable powders and water dispersible granules first. Agitate until these are uniformly dispersed, meanwhile adding water until the tank is 90% full. Add suspension concentrates (flowables) then soluble concentrates. Emulsifiable concentrates go in last. Top off the tank with water and continue agitation until all the ingredients are properly mixed.

Observe any mixing sequence instructions mentioned on the tank mix products.

RESISTANT WEEDS WARNING

GROUP	I	HERBICIDE
-------	----------	-----------

Genfarm LV Ester 600 Herbicide (LV Ester 600) is a member of the phenoxy group of herbicides. LV Ester 600 has the disruptors of plant cell growth mode of action. For weed resistance management LV Ester 600 is a Group I herbicide. Some naturally occurring weed biotypes resistant to LV Ester 600 and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by LV Ester 600 or Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Genfarm Crop Protection Pty Ltd accepts no liability for any losses that may result from the failure of LV Ester 600 to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Legume Tolerance: If clovers are present, care should be taken to ensure that they have reached the 3-4 leaf stage before spraying. Rates above 460 mL of this product per hectare will destroy most clovers, whilst lucerne and medics are susceptible at any strength.

Drift Warning: DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray drift onto nearby susceptible plants, adjacent crops, crop lands or pastures. In particular, avoid spray drift and vapour movement onto susceptible crops such as: cotton, tobacco, tomatoes, vines, fruit trees, ornamentals, oil seed and legume crops and other susceptible plants and trees (e.g. Kurrajongs, Belahs, Eucalypts).

DO NOT use unless wind speed is more than 3 kilometres per hour and less than 15 kilometres per hour as measured at the application site.

DO NOT apply with smaller than coarse to very coarse spray droplets according to the ASAE S572 definition for standard nozzles.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied on any plants at any time as recommended in the Directions For Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used container.

STORAGE AND DISPOSAL

Storage of all containers:

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.

DO NOT store near food, feedstuffs, fertilisers or seed.

Disposal (20L, 100L) This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or preferably pressure rinse container for disposal. Dispose of rinsate by adding to the spray tank. Do not dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap and the closed system valve present in 100L containers removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management site. The cap should not be replaced but may be taken separately. The closed system valve should be disposed of in a local authority landfill.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Disposal (110L)

Do not tamper with the dry valves or security seal. Do not contaminate the drum with water or any other foreign matter. After each use of the product ensure that the dry valve coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained. Add the rinsings to the spray tank. When the drum is empty close all valves and return to the point of purchase. The drum remains the property of Genfarm and must be returned.

1000L - Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray, wear PVC or rubber apron, elbow length PVC gloves and face shield. When using the prepared spray, wear face shield. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. (Phone: Australia 13 11 26).

MATERIAL SAFETY DATA SHEET

For further information, refer to the Material Safety Data Sheet (MSDS) which is available from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Genfarm Crop Protection Pty Ltd. accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd. will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

Genfarm Crop Protection Pty Ltd
Suite 2, Level 3, 64 Talavera Road,
North Ryde,
NSW, 2113
Tel: (02) 9889 5400