

Main panel

---

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**Blazer<sup>®</sup>**  
**Selective Herbicide**

**ACTIVE CONSTITUENT: 224 g/L ACIFLUORFEN (present as the sodium salt)**

GROUP	<b>G</b>	HERBICIDE
-------	----------	-----------

**For the selective control of certain broadleaf weeds and grasses in mung beans, peanuts, soybeans, green beans and seed crops of Siratro and Stylo as specified in the DIRECTIONS FOR USE table**

<sup>®</sup> Registered Trademark

**UNITED PHOSPHORUS LTD**

ABN 76 066 391 384

**Suite 416, 14 Lexington Drive, Norwest Business Park,  
Bella Vista, NSW 2153 AUSTRALIA**

TELEPHONE 02 8824 7277

FACSIMILE 02 8814 6469

**Contents: 10 Litres**

APVMA Approval No. 41650/10/0409

*Ancillary panel*

**DIRECTIONS FOR USE:**

**RESTRAINTS:**

DO NOT mix with oils, fertilisers or other pesticides, except where indicated.

DO NOT apply when crops or weeds are under stress conditions.

DO NOT apply as a post-emergence treatment to control hard to kill weeds when rainfall is expected within 6 hours.

**POST-EMERGENCE TREATMENTS:**

<b>CROP</b>	<b>WEEDS CONTROLLED</b>	<b>WEED SIZE</b>	<b>RATE</b>	<b>CRITICAL COMMENTS</b>
Soybeans (Qld, NSW, WA only) Peanuts (Qld, WA only) Mung beans (Qld only) Adzuki beans ( <i>Vigna angularis</i> ) (All States)	Apple of Peru, turnip weed, wild radish	Cotyledon to 2 true leaves	1.0 L/ha	Apply with thorough coverage to young actively growing weeds.
		2 to 6 true leaves	1.5 L/ha	
	Prince of Wales Feather, redshank, thornapple	Cotyledon to 2 true leaves	2.0 L/ha	
		2 to 6 true leaves		
	Pigweed	Cotyledon to 50 mm diameter	1.0 L/ha	
		50 to 100 mm diameter	1.5 L/ha	
	Annual ground cherry, wild gooseberry	Cotyledon to 4 true leaves	1.0 L/ha	
		4 to 8 true leaves	1.5 L/ha	
		8 leaves to 200 mm high	2.0 L/ha	
	Blackberry nightshade, bladder ketmia, capeweed, fumitory, Noogoora burr	Cotyledon to 4 true leaves	1.5 L/ha	
	Bellvine	2 to 4 true leaves		
	Bellvine	4 to 6 true leaves	2.0 L/ha	
		Giant pigweed	Cotyledon to 2 true leaves	
	Jute	Cotyledon to 4 true leaves	1.5 L/ha	
		4 to 6 true leaves	2.0 L/ha	
Sesbania	Cotyledon to 200 mm high	2.0 L/ha		
Mung beans (NSW only)	bladder ketmia, Noogoora burr		Cotyledon to 4 true leaves	
Mung beans (Qld, NSW only)	See weeds as above	Apply at the weed stage as indicated above	1.0 L/ha plus Hasten* at 1 L/100 L water (1% v/v)	Do NOT exceed 1 L/ha BLAZER with this mixture.

Green beans (All States)	Prince of Wales Feather ( <i>Amaranthus powellii</i> )	2 to 4 leaf stage	250 mL to 500 mL/ha	DO NOT exceed 2 L of product per hectare per season. Apply up to 3 times by boom spray <ul style="list-style-type: none"> <li>after the fully expanded unfoliate leaf stage of the crop.</li> <li>when crop is at 6 to 7 leaf stage and/or when crop is at early flowering stage.</li> </ul> <b>See Application section below.</b>
			250 mL to 750 mL/ha	
Peanuts (Qld only)	Bellvine	2 to 6 true leaves	1.0 L BLAZER <b>PLUS</b> 1.0 to 1.5 L of 400 g/L formulation of 2,4- DB/ha	Apply with thorough coverage as a tank mix to young actively growing weeds. Use the higher rate of 2,4-DB where larger weeds are present. The mixture will give additional control of larger weeds than given by BLAZER alone.
	Noogoora burr	Cotyledon to 6 true leaves		
Seed crops of Siratro and common and fine- stem Stylo ( <i>Stylosanthes guianensis</i> only) (Qld only)	Billygoat weed, gambia pea, glycine, greenleaved desmodium, five-leaved cassia	Cotyledon to 4 true leaves	2.0 L/ha	Apply with thorough coverage to young actively growing weeds. Do NOT use on other Stylos.

Treatment of weeds larger than those listed is not recommended because control may be incomplete due to increased weed tolerance and poor spray coverage. However, in a crop salvage situation, BLAZER may be used at dosages up to 4 L/ha, providing the application is not made within 50 days of harvest.

**PRE-EMERGENCE TREATMENTS: Soybeans and peanuts only. Not recommended for mungbeans.**

CROP	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Peanuts, Soybeans (Qld, WA only)	Apple of Peru, black bindweed, blackberry nightshade, fat hen, fumitory, jute, narrow leaf plantain, Noogoora burr, pigweed, potato weed, redshank, scarlet pimpernel, turnip weed, wild gooseberry, wild radish	3.0 L/ha	Apply as a broadcast spray after sowing but before the crops and weeds emerge. Applications should be made to a well prepared, clod free soil of relatively fine tilth. Mechanical soil incorporation is not recommended as this reduces the activity of BLAZER.
	Annual ground cherry, barnyard grass, bladder ketmia, liverseed grass, Mossman river grass	4.0 L/ha	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIODS:**

**GREEN BEANS, ADZUKI BEANS: DO NOT HARVEST FOR 28 DAYS AFTER APPLICATION.**

**CROP SALVAGE SITUATION WHEN BLAZER IS USED AT UP TO 4 L/HA: DO NOT HARVEST FOR 50 DAYS AFTER APPLICATION.**

**GENERAL INSTRUCTIONS**

**RESISTANT WEEDS WARNING:**



BLAZER Selective Herbicide is a member of the Diphenyl ether group of herbicides. The product has the inhibitor of protoporphyrinogen oxidase mode of action. For weed resistance management the product is a Group G herbicide. Some naturally occurring weed biotypes resistant to the product and other Group G herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group G herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, United Phosphorus Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

### **MIXING**

BLAZER mixes readily with water. Fill the spray tank about two-thirds full with water and add the required amount of BLAZER. Fill the spray tank and mix. Maintain sufficient agitation during mixing and spraying to ensure a uniform spray mixture.

Use of Surfactant: The use of a low foam, non-ionic surfactant at the labelled rate should be used for maximum weed control. This surfactant should be added to the spray mixture after adding the required amount of BLAZER.

Use of crop oil adjuvants: BLAZER may be applied to mung beans at the reduced rate of 1 L/ha in combination with Hasten at 1 L/100 L (1 %v/v). The adjuvant should be added to the spray mixture after adding the required amount of BLAZER. Use of this mixture may increase crop phytotoxicity, however the plants will soon recover and yield will be unaffected.

### **APPLICATION**

Spray Equipment: BLAZER can be applied by ground equipment using standard low pressure herbicide sprayers, equipped with hollow cone or flat fan nozzles. The spray equipment should be calibrated to deliver between 100 and 300 litres of spray mixture per hectare at a pressure of 250 to 300 kPa at the nozzle tips.

BLAZER may be applied as a banded treatment directly over the crop row. When this method of application is used, it is important that the spray equipment be calibrated to give the correct dosage per sprayed hectare, and that the spray pattern gives maximum coverage of the weed foliage present. Before using BLAZER, it is important to check that the equipment is clean and that all materials from previous uses are completely flushed from the system.

Aerial Application: QUEENSLAND AND Northern NSW ONLY. BLAZER can be applied from a fixed-wing aircraft or helicopter, using a minimum of 48 litres of spray per hectare. Nozzles on spray booms should not be placed any closer to wing tips than three-quarters of the wing span; this will minimise the formation of spray or wing tip vortice roll. A spraying height of 2 to 3 metres over the crop is recommended. Swirl jet or hollow cone nozzles are suggested. Keep pressures below 275 kPa. Nozzles should point to the rear of the aircraft and not be pointed downward more than 20 degrees.

DO NOT USE MICRONAIRE EQUIPMENT FOR BLAZER APPLICATION.

### **GREEN BEAN CROPS**

Early application is important. Treatment later than the recommended weed growth stage may result in inadequate control of Prince of Wales Feather (*Amaranthus powellii*).

The application of BLAZER may cause some crop phytotoxicity: e.g. some tip burning and stunting. The beans usually grow out of this, though they may remain about 10 cm shorter than unsprayed beans. This does not appear to reduce yield. It is recommended that users consult an experienced field agronomist prior to using this product. Additionally it is recommended to apply in the evening if possible. Increased phytotoxicity may occur under warm humid conditions. A non-ionic wetting agent may be added at 0.1% v/v. Thorough coverage is essential. Do not apply earlier than fully-expanded unifoliate leaf stage. Use higher rate after the beans have reached the first fully-expanded trifoliate leaf stage, to target *Amaranthus* which is greater than the 2-leaf stage. Multiple applications should only be used if necessary to treat the weed problem.

## **POST-EMERGENCE TREATMENTS**

Best results are obtained when the crop and weeds are free from moisture and temperature stress, and when the relative humidity is around 65% or higher. DO NOT apply BLAZER if soil temperature is below 15°C or air temperature is below 20°C. Crop damage will also be increased when BLAZER is applied when air temperatures are above 30°C.

Since it is important that the maximum possible weed coverage is obtained, application should be made before the crop canopy shields the weeds from spray contact but after the maximum number of weed seedlings have emerged. This will usually occur between the 2nd and 5th trifoliate stages of the crop, or about one month after sowing. Thorough coverage is essential for optimum weed control. After treatment, soybean leaves may exhibit leaf cupping, crinkling and speckling, particularly on the youngest leaves present at the time of application. Soybeans recover from this injury and continue growing at a normal rate with no adverse effect on crop yield.

## **COMPATIBILITY:**

BLAZER is compatible with BASAGRAN\*, haloxyfop, fluazifop-P, Select\*, and Sertin\*.

## **PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Wash sprayer thoroughly after use.

## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT apply when weather conditions favour run off or drift from areas treated.

## **STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

## **SAFETY DIRECTIONS**

May irritate the eyes and skin. Avoid contact with eyes and skin. If product on skin, immediately wash area with soap and water.

## **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

## **MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet.

**CONDITIONS OF SALE:** United Phosphorus accepts responsibility for the consistent quality of the product; however since the use and application of the product is beyond control, the company accepts no responsibility whatsoever for any loss, damage or other result following the use of the product whether used in accordance with directions or not; other than those mandatorily imposed by statutes, the liability is limited to the replacement of the goods and is conditional upon a claim made in writing and, where necessary, a sufficient part of the goods being returned for proper examination by the company within thirty days of sale.

\* trademarks

**Batch No.:**

**DOM:**

**Bar Code:**