

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CONQUEST
ALPHA FORTE 250 SC

INSECTICIDE

Active Constituent: 250 g/L ALPHA-CYPERMETHRIN

GROUP	3A	INSECTICIDE
--------------	-----------	--------------------

For the control of certain insect pests, including redlegged earth mite on certain field crops and pastures and certain insect pests on fruit and vegetable crops as indicated in the DIRECTIONS FOR USE table in the ATTACHED BOOKLET

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

Contents:

20 L

(5 L)

Conquest Crop Protection Pty Ltd
ABN 84 098 814 932
Level 1, 4 Collingwood Street
Osborne Park, WA 6017

Telephone (08) 9347 0500
Facsimile: (08) 9347 0551

APVMA Approval No: 65245/1021102



PRECAUTION

Do NOT use human flaggers unless they are protected by engineering controls such as vehicles with cabs.

RE-ENTRY PERIOD

Do NOT allow entry into treated areas for 12 hours after application. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist, a washable hat and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF LIVESTOCK

Dangerous to bees. Do NOT spray on any plants in flower while bees are foraging. CONQUEST ALPHA FORTE 250 SC INSECTICIDE is known to have a deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in early morning and late evening while bees are not foraging.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish and aquatic invertebrates. Do NOT contaminate fish ponds, dams, drains, rivers or streams with product or used containers. Drift and run-off from treated areas may be hazardous to fish in adjacent sites.

STORAGE AND DISPOSAL

Store in the closed, original container, in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Harmful if inhaled. Will irritate the eyes and skin. Facial skin contact may cause temporary facial numbness. Avoid contact with eyes and skin. Do NOT inhale spray mist. When opening the container, preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Telephone 131126 Australia-wide.

MSDS

Additional information is listed in the Material Safety Data Sheet. A copy of the MSDS is available from our website: www.conquestag.com.au

CONDITIONS OF SALE

Conquest Crop Protection Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CONQUEST

ALPHA FORTE 250 SC

INSECTICIDE

Active Constituent: 250 g/L ALPHA-CYPERMETHRIN

GROUP	3A	INSECTICIDE
--------------	-----------	--------------------

For the control of certain insect pests, including redlegged earth mite on certain field crops and pastures and certain insect pests on fruit and vegetable crops as indicated in the DIRECTIONS FOR USE table in this LEAFLET

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

Conquest Crop Protection Pty Ltd
ABN 84 098 814 932
Level 1, 4 Collingwood Street
Osborne Park, WA 6017

Telephone (08) 9347 0500
Facsimile: (08) 9347 0551



DIRECTIONS FOR USE

RESTRAINT: Do NOT apply if rain is expected within 6 hours after application.

Asparagus: DO NOT apply more than 6 times per season.

NOTE: This product is ineffective against synthetic pyrethroid resistant *Helicoverpa armigera* larvae longer than 5 mm. All *Helicoverpa armigera* in NSW and Qld should be treated as being resistant to synthetic pyrethroids. Refer to RESISTANCE MANAGEMENT under GENERAL INSTRUCTIONS. This product is ineffective against synthetic pyrethroid resistant *Plutella xylostella*.

CROP	INSECT PESTS	STATE	RATE	WHP	CRITICAL COMMENTS
Asparagus (not for use on white asparagus)	Garden weevil (<i>Phlyctinus callosus</i>)	WA only	40 mL/100 L	1 day	Apply in spring after weevil emergence, at up to 500 L spray solution per hectare. Day time spraying is effective but superior control may be achieved if spray is applied at night. Repeat applications as required, depending on pest pressure. Application to fern, after spear harvest may reduce carry-over of garden weevil for the following season. Caution: Not for use on white asparagus, there have been reports of some phytotoxicity when using alpha-cypermethrin.
Banksias	Banksia moth (<i>Danima banksiae</i>)	WA only	8 mL/100 L	-	Apply on a regular program at 2 weeks intervals at early flower development. Commence spraying when blooms are immature and continue until flowers are fully developed.
Broccoli, Brussels sprouts, Cabbages, Cauliflowers, Chinese cabbage, Kale, Kohlrabi, Turnips	Cabbage moth (<i>plutella xylostella</i>), cabbage white butterfly (<i>Pieris rapae</i>), <i>Helicoverpa punctigera</i> , <i>Helicoverpa armigera</i>	All States	<p>LOW VOLUME 160 mL/ha</p> <p>HIGH VOLUME 20 mL/100 L</p>	1 day (Harvest)	<p>Apply according to pest incidence. When reinfestation is continuous, treatment every 7-10 days may be required. Add a non-ionic wetting agent at recommended label rate.</p> <p>LOW VOLUME: Ground Rig Application: Apply in 100 to 600 L of water per hectare as a fine spray, (ie. A droplet size of 100 to 200 microns).</p> <p>Aerial Application: Apply in 20 to 60 L of water per hectare as a spray of 100 to 150 microns droplet size.</p> <p>HIGH VOLUME: Gradually increase the spray volume as the plants grow, from 600 L/ha just after transplanting to 1000 L/ha at maturity. Apply as a medium spray (ie. A droplet size of 200 to 400 microns VMD).</p> <p><i>Helicoverpa armigera</i> in NSW and Qld. Follow the application directions for the pest above. Apply as required according to pest incidence. Thorough and frequent crop checks are essential. Preferably apply to eggs. Apply to larvae only if they are less than 5 mm long.</p>
	Cluster caterpillar (<i>Spodoptera litura</i>)	All States			
Canola	Native budworm (<i>Helicoverpa punctigera</i>)	NSW, SA, Tas, Vic, WA only	80 or 120 mL/ha	21 days (cutting for harvest or stockfeed or grazing)	<p>Do NOT apply more than a total 160 mL/ha per season to any one crop.</p> <p>Inspect the crop regularly during and immediately after flowering. Apply when damaging pest numbers first appear on the crop and repeat if necessary.</p>
	Tobacco looper (<i>Chrysodeixis argentifera</i>)	NSW, SA, Tas, Vic only	80 or 120 mL/ha		For aerial application, use a total volume of 30-35 L/ha and apply in the cooler part of the day. Use the higher rate if larvae longer than 10 mm are present.
	Vegetable weevil (<i>Listroderes difficilis</i>)	NSW, ACT, Vic, Tas, SA, WA only	160 mL/ha		Crops should be inspected as they emerge. Border sprays are required to control invading adults. Apply when cotyledons and leaves are being eaten or the plant lopped. Repeat as necessary.
	Redlegged earth mite (<i>Halotydeus destructor</i>)	All States except NT and Qld	40 mL/ha		Pre-emergence: Apply by ground rig only. Treat infested paddocks after sowing but prior to crop emergence when soil is moist. Monitor redlegged earth mite numbers and re-treat if necessary.
	Redlegged earth mite, blue oat mite (<i>Penthaleus major</i>)		20 mL/ha		Apply when mite numbers reach damaging levels. Do NOT apply as a pre-emergence treatment.

CROP	INSECT PESTS	STATE	RATE	WHP	CRITICAL COMMENTS	
Chickpeas	Native budworm (<i>Helicoverpa punctigera</i>)	WA only	64 mL/ha	21 days (Harvest) 35 days (Grazing)	Apply to open, less dense crops when numbers of newly hatched larvae first appear on the crop and repeat as necessary.	
		NSW, Vic, SA, WA only	80 or 120 mL/ha		Apply when pest numbers reach damaging levels and repeat if necessary. Use the higher rate if larvae longer than 10 mm are present. Best results will be obtained by spraying at egg hatch.	
	Redlegged earth mite (<i>Halotydeus destructor</i>)	NSW, Vic, Tas, SA, WA only	40 mL/ha		Pre-emergence: Apply by ground rig only. Treat infested paddocks after sowing but prior to crop emergence when soil is moist. Monitor redlegged earth mite numbers and re-treat if necessary. Do NOT apply as a ULV application.	
			20 mL/ha		Apply when mite numbers reach damaging levels. Do NOT apply as a pre-emergence treatment. Do NOT apply as a ULV application.	
	Cutworm (<i>Agrotis</i> spp.)		30 mL/ha		Check emerging or establishing crops in the late afternoon and evening for caterpillars crawling on the soil surface and feeding on the seedlings. Spray in the late afternoon and evening.	
Cotton	Native budworm (<i>Helicoverpa punctigera</i>)	NSW, NT, Qld, WA only	120 mL/ha	14 days (Harvest)	Apply as indicated by field checks using rates appropriate for the infestation level determined. Application should be timed to coincide with egg hatching and before larvae are in protected feeding sites. Apply when there are up to 75 eggs and/or up to 5 larvae less than 5 mm long per 100 terminals.	
			160 mL/ha		Apply when there are up to 150 eggs and/or up to 10 larvae less than 5 mm long per 100 terminals and/or when larvae between 5 and 10 mm are present.	
			200 mL/ha		Apply when there are up to 150 eggs and/or more than 10 larvae less than 5 mm long per 100 terminals and/or when larvae longer than 10 mm are present.	
			Cotton bollworm (<i>Helicoverpa armigera</i>)		120 mL/ha	Preferably apply to eggs. Apply to larvae only if they are less than 5 mm long.
					160 mL/ha	Apply when there are up to 75 eggs and/or up to 5 larvae less than 5 mm long per 100 terminals.
					200 mL/ha	Apply when there are up to 150 eggs and/or up to 10 larvae less than 5 mm long per 100 terminals.
	Rough bollworm (<i>Earias huegeli</i>)		120 or 160 mL/ha		Apply when there are more than 150 eggs and/or more than 10 larvae less than 5 mm long per 100 terminals.	
					Apply when an average of 2 or more larvae are present per 100 bolls. It is essential to detect and treat infestations in the early stages before larvae are established or concealed in bolls deep in the canopy. Use the higher rate if larvae longer than 10 mm are present. Best results will be obtained by applying at egg hatch.	

CROP	INSECT PESTS	STATE	RATE	WHP	CRITICAL COMMENTS
Cereals (Winter)	Cutworm (<i>Agrotis</i> spp.)	NSW, ACT, Vic, SA, WA only	30 mL/ha	7 days (Harvest) 14 days (Stubble grazing)	Do NOT apply more than a total of 216 mL/ha per season to any one crop. Check emerging and establishing crops in the late afternoon and evening for caterpillars crawling on the soil surface and feeding on the seedlings. Spray in late afternoon or evening.
		Qld only	30 or 60 mL/ha		In Qld, use the higher rate when the infestation is severe, or when there are larvae longer than 10 mm, or when longer residual activity is required.
	Webworm (<i>Hednota</i> spp.)	NSW, Vic, SA, WA only	30 mL/ha		Pre planting: May be applied with knock-down herbicides prior to planting. Apply from the last week in May when the larvae have emerged. Do NOT apply to dense pasture. All pasture should be closely grazed prior to application to ensure adequate spray penetration. Apply in minimum of 100 L of water per hectare. Repeat as required. Post crop emergence: Inspect crop regularly from emergence and apply at the first sign of pest activity. Repeat as required.
	Common armyworm (<i>Mythimna convecta</i>), southern armyworm (<i>Persectania ewingii</i>)	All States	96 mL/ha		Apply before "head lopping" occurs and when there are 2 or more larvae per square metre. Spray in the cool of the day (usually late afternoon) when larvae are most active. Ensure the spray penetrates the crop. This rate is effective on larvae up to 20 mm in length. Monitor crop closely and re-treat if necessary. Poor control may occur in crops that have lodged. See application section for correct water rates.
	Redlegged earth mite (<i>Halotydeus destructor</i>)	NSW, ACT, Vic, Tas, SA, WA only	40 mL/ha		Pre-emergence: Apply by ground rig only. Treat infested paddocks after sowing but prior to crop emergence when soil is moist. Monitor redlegged earth mite numbers and re-treat if necessary.
	Redlegged earth mite, blue oat mite (<i>Penthaleus major</i>)	NSW, ACT, Vic, Tas, SA, WA only	20 mL/ha		Apply when mite numbers reach damaging levels. Spray seedling crops if silvering or whitening (bleaching) of the leaves is causing a reduction in crop growth. If possible, spray on a calm, mild morning when mites are actively feeding on crop leaves. Do NOT apply as a pre-emergence treatment.
	Aphids (<i>Rhopalosiphum</i> spp.) (barley yellow dwarf virus vectors)	NSW, ACT, Vic, Tas, SA, WA only	50 mL/ha		To control aphids, sprays should be applied at 3 and 7 weeks after emergence to reduce aphid colonisation and the spread of Barley Yellow Dwarf Virus. This will also reduce the effect of feeding aphid damage.
Eucalypt plantations	Adults and larvae of Tasmanian eucalyptus leaf beetle (<i>Chrysophtharta bimaculata</i>)	Tas only	100 mL/ha	-	Apply by fixed wing aircraft or by helicopter using hydraulic or Micronair equipment, to the crowns of eucalypt trees. Micronair application in 5 litres of water/ha has proved effective. Apply before insect damage causes severe defoliation. Treatment will control small and large larvae as well as adult beetles.

CROP	INSECT PESTS	STATE	RATE	WHP	CRITICAL COMMENTS
Faba beans	Native budworm (<i>Helicoverpa punctigera</i>)	WA only	64 mL/ha	4 weeks (Harvest) 35 days (Grazing)	Apply to open, less dense crops when numbers of newly hatched larvae first appear on the crop and repeat as necessary.
		NSW, Vic, Tas, SA, WA only	80 or 120 mL/ha		Apply when pest numbers reach damaging levels and repeat if necessary. Use the higher rate if larvae longer than 10 mm are present. Best results will be obtained by spraying at egg hatch.
	Regged earth mite (<i>Halotydeus destructor</i>)	40 mL/ha	Pre-emergence: Apply by ground rig only. Treat infested paddocks after sowing but prior to crop emergence when soil is moist. Monitor redlegged earth mite numbers and re-treat if necessary. Do NOT apply as a ULV application.		
	Redlegged earth mite (<i>Halotydeus destructor</i>), Blue oat mite (<i>Penthaleus major</i>)	20 mL/ha	Post-emergence: Apply to established crops when mite numbers reach damaging levels. Do NOT apply as a pre-emergence treatment. Do NOT apply as a ULV application.		
	Cutworm (<i>Agrotis</i> spp.)	30 mL/ha	Check emerging or establishing crops in the late afternoon and evening for caterpillars crawling on the soil surface and feeding on the seedlings. Spray in the late afternoon and evening.		
Field peas	Native budworm (<i>Helicoverpa punctigera</i>)	WA only	64 mL/ha	4 weeks (Harvest)	Apply to open, less dense crops when damaging numbers of newly hatched larvae first appear on the crop and repeat as necessary.
	Native budworm (<i>Helicoverpa punctigera</i>)	NSW, Vic, Tas, SA, WA only	80 or 120 mL/ha		Apply when pest numbers reach damaging levels and repeat if necessary. Use the higher rate if larvae longer than 10 mm are present. Best results will be obtained by spraying at egg hatch.
	Pea weevil (<i>Bruchus pisorum</i>)	NSW, Vic, Tas, SA, WA only	64 or 80 mL/ha		Apply during flowering prior to egg laying when the adult weevil population reaches on or more per 25 sweeps of a sweep net. Use the higher rate for longer residual protection.
	Cutworm (<i>Agrotis</i> spp.)		30 mL/ha		Check emerging and establishing crops in the late afternoon and evening for caterpillars crawling on the soil surface and feeding on the seedlings. Spray in late afternoon and evening.
	Redlegged earth mite (<i>Halotydeus destructor</i>)	NSW, Vic, Tas, SA, WA only	40 mL/ha		Pre-emergence: Apply by ground rig only. Treat infested paddocks after sowing but prior to crop emergence when soil is moist. Monitor redlegged earth mite numbers and re-treat if necessary.
	Redlegged earth mite, blue oat mite (<i>Penthaleus major</i>)		20 mL/ha		Apply when mite numbers reach damaging levels. Do NOT apply as a pre-emergence treatment.
Grapevines (non bearing)	Pink cutworm (<i>Agrotis munda</i>), apple weevil (<i>Curculio beetle</i>) (<i>Otiorynchus cribricollis</i>), garden weevil (<i>Phlyctinus callosus</i>)	NSW, Vic, Tas, SA, WA only	40 mL/100 L	-	Monitor young vines during spring and early summer and apply at the first signs of leaf damage. Spray the leaves, canes and the soil around each vine to a diameter of 30 cm. 70-80 mL of dilute spray should be sufficient for each vine. If pest infestation persists, a second application may be required after three weeks.
Lettuce	<i>Helicoverpa</i> spp.	All States	LOW VOLUME 160 mL/ha HIGH VOLUME 20 mL/100 L	3 days (Harvest)	Thoroughly and regularly check the crop. Apply at the first sign of pest activity. Preferably apply to eggs. Apply to <i>H. armigera</i> ONLY if larvae are less than 5 mm long. Repeat according to pest incidence.

CROP	INSECT PESTS	STATE	RATE	WHP	CRITICAL COMMENTS
Linola	Native budworm (<i>Helicoverpa punctigera</i>)	WA only	64 or 80 mL/ha	12 weeks (Harvest)	Do NOT apply more than a total 160 mL/ha per season to any one crop. Inspect the crop regularly during and immediately after flowering. Apply when damaging pest numbers first appear on the crop. For aerial application, apply during the cooler part of the day in a total volume of 30-35 L/ha. Use the higher rate if larvae longer than 10 mm are present. Refer to application section for water rates.
Linseed	Native budworm (<i>Helicoverpa punctigera</i>)	NSW, Vic, Tas, SA, WA only	80 or 120 mL/ha	14 days (Harvest)	Inspect the crop regularly during and immediately after flowering. Apply when damaging pest numbers first appear on the crop and repeat if necessary. Use the higher rate if larvae longer than 10 mm are present. Best results will be obtained by spraying at egg hatch. Refer to application section for water rates.
	Cutworm (<i>Agrotis</i> spp.)	NSW, ACT, Tas, SA only	30 mL/ha		Check emerging and establishing crops in the late afternoon and evening for caterpillars crawling on the soil surface and feeding on seedlings. Spray in late afternoon and evening.
Lucerne (seed and forage crops)	Native budworm (<i>Helicoverpa punctigera</i>)	NSW, Vic, Tas, SA, WA only	64 mL/ha	14 days (grazing or cutting for stockfeed)	Do NOT apply more than one application per cut or grazing for animal feed. Apply when pest populations reach economically damaging levels. Apply to larvae less than 5 mm in length.
	Green mirid (<i>Creontiades dilutus</i>)				Do NOT apply more than one application per cut or grazing for animal feed. Apply when pest populations reach economically damaging levels.
Lupins	Native budworm (<i>Helicoverpa punctigera</i>)	NSW, ACT, Vic, SA only	80 or 120 mL/ha	4 weeks (Harvest)	Do NOT apply more than a total of 240 mL/ha per season to any one lupin crop. Apply when damaging pest numbers first appear on the crop and repeat if necessary. Use the higher rate if larvae longer than 10 mm are present. Best results will be obtained by spraying at egg hatch.
		WA only	48 or 80 mL/ha		Spraying should be timed to precede the first visible damage to the pods. Use the higher rate when the infestation is severe, or when residual activity is required.
	Cutworm (<i>Agrotis</i> spp.)	NSW, ACT, Vic, Tas, SA, WA only	30 mL/ha	Check emerging and establishing crops in the late afternoon and evening for caterpillars crawling on the soil surface and feeding on seedlings. Spray in late afternoon and evening.	
	Common armyworm (<i>Mythimna convecta</i>), southern armyworm (<i>Persectania ewingii</i>)	ACT, NSW only	96 mL/ha	Spray in the cool of the day (late afternoon) when larvae are most active.	
	Redlegged earth mite (<i>Halotydeus destructor</i>)	NSW, Vic, Tas, SA, WA only	40 mL/ha	4 weeks (Harvest)	Pre-emergence: Apply by ground rig only. Treat infested paddocks after sowing but prior to crop emergence when soil is moist. Monitor redlegged earth mite numbers and re-treat if necessary.
	Redlegged earth mite, blue oat mite (<i>Penthaleus major</i>)		20 mL/ha		Apply when mite numbers reach damaging levels. Do NOT apply as a pre- emergence treatment.

CROP	INSECT PESTS	STATE	RATE	WHP	CRITICAL COMMENTS
Maize	Corn earworm (<i>Helicoverpa armigera</i>)	Qld, NSW, ACT, Vic, NT only	120 or 160 mL/ha	7 days (Harvest)	Thoroughly and regularly check the crop. Apply from early silking according to pest incidence. Use the higher rate if larvae longer than 10 mm are present. In Qld, NSW and NT, preferably apply to eggs or apply to larvae only if they are less than 5 mm long.
	Native budworm (<i>Helicoverpa punctigera</i>)	All States			Thoroughly and regularly check the crop. Apply when infestation reaches an economically damaging level and repeat if necessary. Best results will be obtained by applying at egg hatch. Use the higher rate if larvae longer than 10 mm are present.
Mung beans, navy beans	Native budworm (<i>Helicoverpa punctigera</i>)	Qld, NSW, ACT, NT only	120 or 160 mL/ha	7 days (Harvest)	Crop checking should be aimed to detect larvae as they hatch. Small larvae are easier to kill than large larvae. Apply when the number of larvae feeding on flowers or pods reach 1 to 2 per metre of row. Repeat as required. Use the higher rate when larvae larger than 10 mm are present or when canopy is dense. Best results will be obtained by spraying at egg hatch.
	Corn earworm (<i>Helicoverpa armigera</i>)				Thoroughly and regularly check the crop. Apply when the infestation reaches an economically damaging level and repeat as required. Preferably apply to eggs. In Qld and NSW, apply to larvae only if they are less than 5 mm long. Use the higher rate when pest pressure is high.
Pastures (legume and grass based pastures)	Wingless grasshopper (<i>Phaulacridium vittatum</i>)	All States	64 mL/ha	3 days (Grazing) 14 days (Cut for stockfeed)	Do NOT apply more than a total 128 mL/ha per season. Apply to infested areas and repeat as necessary. Spraying is most effective on newly emerged hoppers before they begin dispersing. Spray in the warmer parts of the day when hoppers are exposed. Later sprays should be applied before the start of egg laying. Good coverage is essential.
	Brown pasture looper (<i>Ciampa arietaria</i>)	NSW, Vic, Tas, SA, WA only	20 mL/ha		Apply when pest infestation reaches an economically damaging level.
	Blackheaded pasture cockchafer (<i>Aphodius tasmaniae</i>)	NSW, Vic, Tas, SA only	40 mL/ha		Spraying is most effective when larvae are detected and treated early. Suspect paddocks should be dug after the first substantial rain in April/May and inspected to ensure grubs are present in sufficient numbers to warrant treatment. Spraying after June will give poorer results.
	Redlegged earth mite (<i>Halotydeus destructor</i>)	ACT, NSW, SA, Tas, Vic, WA only			Pre-emergence: Apply by ground rig only. Treat infested paddocks after sowing but prior to crop emergence when soil is moist. Monitor redlegged earth mite numbers and re-treat if necessary.
	Redlegged earth mite, blue oat mite (<i>Penthaleus major</i>)	ACT, NSW, SA, Tas, Vic, WA only	20 mL/ha	3 days (Grazing) 14 days (Cut for stockfeed)	Apply when mite numbers reach damaging levels. Autumn/Winter Apply 4 to 7 weeks after the opening rains in late autumn/early winter when RLEM are present (2-3 weeks after egg hatch occurs). CONQUEST ALPHA FORTE 250 SC INSECTICIDE is rainfast after spray deposits have dried on the leaf surface. CONQUEST ALPHA FORTE 250 SC INSECTICIDE can be mixed with herbicides used for winter cleaning of sub clover pastures. Spring If RLEM/BOM numbers increase in the spring, spray when damage is observed and again before diapause egg production begins. CONQUEST ALPHA FORTE 250 SC INSECTICIDE can be mixed with herbicides used for spray topping pastures if timing coincides. Do NOT apply as a pre-emergence treatment.

CROP	INSECT PESTS	STATE	RATE	WHP	CRITICAL COMMENTS
Pome fruit: apples, pears	Apple weevil (<i>Otiorhynchus cribricollis</i>), garden weevil (<i>Phlyctinus callosus</i>)	NSW, Vic, SA, WA only	40 mL/100 L water	14 days (Harvest)	Spray approx. 1-2 litres of solution onto the crotch, trunk and the soil at the base of each tree at peak weevil emergence. This is usually late October – late November for garden weevil, and late November – mid December for apple weevil. Monitor weevil emergence using a single sided cardboard trunk band. Continue monitoring after spraying as a second spray may be needed 3-4 weeks later.
Rice (both aerial and drill sown)	Common armyworm (<i>Mythimna convecta</i>)	NSW only	80 mL/ha	7 days	Do NOT apply more than a total 160 mL/ha per season to any one crop. Inspect crops regularly for the presence of grubs from flowering onwards. Apply when rice damaging pest numbers first appear. Apply by aircraft in 20-30 litres of water/ha, to drained fields only. Spray in the cool of the day (early morning or late afternoon) when larvae are most active. Monitor crop closely and re-treat if necessary. Poor control may occur in crops that have lodged. See application section for correct water rates.
Sorghum	Corn earworm (<i>Helicoverpa armigera</i>), native budworm (<i>Helicoverpa punctigera</i>)	Qld, NSW, ACT, NT only	120 or 160 mL/ha	7 days (Harvest)	Crop checking should commence when the head emerges from the boot and continue at daily intervals until the end of flowering for midge and at weekly intervals until maturity for <i>Helicoverpa armigera</i> . Do NOT apply to tight headed varieties. Apply when there are 2 or more actively feeding larvae per head, or when numbers are sufficient to cause economic damage. Use the higher rate if longer residual control is required. Preferably apply to eggs. Apply to <i>H.armigera</i> larvae only if they are less than 5 mm long. Repeat as required.
	Sorghum midge (<i>Contarinia sorghicola</i>)		40 or 80 mL/ha		Apply when numbers reach 1 to 2 per head, between head emergence and the end of flowering. Repeat as required. Use the higher rate for increased residual protection.
Soybeans	Native budworm (<i>Helicoverpa punctigera</i>)	Qld, NSW, ACT, NT only	120 or 160 mL/ha	7 days (Harvest)	Thoroughly and regularly check the crop. Apply when the number of larvae feeding on flowers plus pods reaches 1 to 2 per metre of row. Repeat as required. Use the higher rate if larvae longer than 10 mm are present. Best results will be obtained by applying at egg hatch.
	Corn earworm (<i>Helicoverpa armigera</i>)				Thoroughly and regularly check the crop. Apply when numbers are sufficient to cause economic damage. Preferably apply to eggs. In Qld and NSW, apply to larvae only if they are less than 5 mm long. Repeat as required. Use the higher rate when pest pressure is high.
Stone fruit: apricots, nectarines, peaches, plums	Apple weevil (<i>Otiorhynchus cribricollis</i>), garden weevil (<i>Phlyctinus callosus</i>)	WA only	40 mL/100 L water	14 days (Harvest)	Spray approx. 1-2 litres of solution onto the crotch, trunk and soil at the base of each tree at peak weevil emergence. This is usually late October – late November for garden weevil, and late November – mid December for apple weevil. Monitor weevil emergence using a single sided cardboard trunk band. Continue monitoring after spraying as a second spray 3-4 weeks later may be needed.

CROP	INSECT PESTS	STATE	RATE	WHP	CRITICAL COMMENTS
Sunflowers	Native budworm (<i>Helicoverpa punctigera</i>)	Qld, NSW, Vic, ACT, NT only	120 or 160 mL/ha	21 days (Harvest)	TO PROTECT BEES and ensure adequate pollination, application during flowering should be avoided. If application is necessary at flowering, apply early morning or late afternoon when bees are not actively foraging. Crop checking should be aimed to detect larvae as they hatch. Small larvae are easier to kill than large larvae. Apply when the infestation reaches an average of 2-3 larvae per head or when economic damage is occurring. Repeat as required. Apply before the heads turn downwards to ensure adequate coverage. Use the higher rate if larvae longer than 10 mm are present. Best results will be obtained by applying at egg hatch.
	Corn earworm (<i>Helicoverpa armigera</i>)				Thoroughly and regularly check the crop. Apply when numbers are sufficient to cause economic damage. Preferably apply to eggs. In NSW and Qld, apply to larvae only if they are less than 5 mm long. Repeat as required. Use the higher rate under heavy pest pressure.
	Grey cluster bug (<i>Nysius clevelandensis</i>), Rutherglen bug (<i>Nysius vinitor</i>)				Apply from budding when adult numbers per plant reach 10 to 15 in dryland crops and 20 to 25 in irrigated crops. After flowering apply when adult numbers on the face of heads reach 20 to 25. Repeat as required. The higher rate should be used when numbers are very high.
	Rutherglen bug (<i>Nysius vinitor</i>)	Vic, Tas, WA only	100 mL/ha		Apply from budding when adult numbers per plant reach 10 to 15 in dryland crops and 20 to 25 in irrigated crops. After flowering, apply when adult numbers on the face of heads reach 20 to 25. Repeat as required.

CROP	INSECT PESTS	STATE	RATE	WHP	CRITICAL COMMENTS	
Sweet corn	Native budworm (<i>Helicoverpa punctigera</i>), corn earworm (<i>Helicoverpa armigera</i>)	All States	120 or 160 mL/ha	7 days (Harvest)	Thoroughly and regularly check the crop. The level of cob damage tolerated varies with market requirements. FRESH MARKET CORN: Apply at 5-8 day intervals, accordingly to pest incidence, from tassel emergence until the silks wither. PROCESSING CORN: Apply from early silking according to pest incidence. Larvae in protected feeding sites within the cob are not effectively controlled. Apply before this situation occurs. Best results will be obtained by applying at egg hatch. Use the higher rate if larvae longer than 10 mm are present. To help contain pyrethroid resistance in <i>Helicoverpa armigera</i> in summer crops, do NOT apply to corn earworm longer than 5 mm.	
Tobacco	Native budworm (<i>Helicoverpa punctigera</i>), tobacco budworm (<i>Helicoverpa armigera</i>)	Vic only	12 or 16 mL/100 L	7 days (Harvest)	Apply from just after transplanting on a 7 to 10 day schedule, according to pest incidence. Apply as a medium to fine spray using hollow and/or solid cone nozzles. The spray volume should be gradually increased as the plants grow, from 200 L/ha just after transplanting to 1000 L/ha at maturity. Use the higher rate when larvae longer than 10 mm are present or when egg laying is intense.	
Tomatoes (bush and trellis)	Native budworm (<i>Helicoverpa punctigera</i>)	All States	LOW VOLUME 80, 120 or 160 mL/ha HIGH VOLUME 8, 12, or 20 mL/100 L	1 day (Harvest)	Do NOT apply to trellis tomatoes by aircraft. Apply on a 7 to 10 day schedule while the pests are active. Use the middle rate when pest activity is high and/or when larvae between 10 and 20 mm in length are present. Use the highest rate when larvae longer than 20 mm are present and/or when interruption of the schedule enables a very severe infestation to develop. LOW VOLUME: By ground-rig: apply in 100 to 400 L of water per hectare as a fine spray. By aircraft: apply in minimum of 10 L of water per hectare as a spray of 100 to 150 microns VMD. HIGH VOLUME: Apply as a medium to fine spray. Gradually increase the spray volume as the plants grow, from 200 L/ha just after transplanting establishment to 1000 L/ha at maturity.	
	Tomato grub (<i>Helicoverpa armigera</i>)	Vic, Tas, SA, WA only				
	Cluster caterpillar (<i>Spodoptera litura</i>)	Qld, NSW, ACT, WA, NT only				
	Tomato grub (<i>Helicoverpa armigera</i>)	Qld, NSW, NT only	LOW VOLUME 120 mL/ha HIGH VOLUME 12 mL/ 100 L			Thoroughly check the crop at 2-3 day intervals from transplanting/ emergence. Apply according to pest incidence. Preferably apply to eggs. Apply to larvae only if they are less than 5 mm long. Apply using the methods described for native budworm above.
	Plague thrips (<i>Thrips imaginis</i>)	Qld, NSW, ACT, Vic, Tas, WA, NT only	LOW VOLUME 130 mL/ha HIGH VOLUME 7.2 mL/ 100 L			The crop should be frequently checked when it is flowering for the presence of the pest. Apply when the infestation reaches an economically damaging level, using the application methods described for native budworm above.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORIZED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

ASPARAGUS, BROCCOLI, BRUSSELS SPROUTS, CABBAGES, CAULIFLOWERS, CHINESE CABBAGE, KALE, KOHLRABI, TOMATOES, TURNIPS: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION
LETTUCE: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION
PASTURES: DO NOT GRAZE FOR 3 DAYS AFTER APPLICATION
DO NOT CUT FOR STOCKFEED FOR 14 DAYS AFTER APPLICATION
MAIZE, MUNG BEANS, NAVY BEANS, RICE, SORGHUM, SOYBEANS, SWEET CORN, TOBACCO: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION
WINTER CEREALS: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION
DO NOT GRAZE TREATED STUBBLE FOR 14 DAYS AFTER APPLICATION
LUCERNE: DO NOT GRAZE OR CUT FOR STOCKFEED FOR 14 DAY AFTER APPLICATION
COTTON, LINSEED, POME FRUIT, STONE FRUIT: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION
CANOLA: DO NOT GRAZE OR CUT FOR STOCKFEED FOR 21 DAYS AFTER APPLICATION
DO NOT CUT AND WINDROW FOR HARVEST FOR 21 DAYS AFTER APPLICATION
CHICKPEAS: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION
DO NOT GRAZE OR CUT FOR STOCKFEED FOR 35 DAYS AFTER APPLICATION
FABA BEANS: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION
DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 35 DAYS AFTER APPLICATION
SUNFLOWERS; DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION
FIELD PEAS, LUPINS: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION
LINOLA: DO NOT HARVEST FOR 12 WEEKS AFTER APPLICATION

GENERAL INSTRUCTIONS

CONQUEST ALPHA FORTE 250 SC INSECTICIDE is a suspension concentrate which acts as a contact and residual insecticide. It can be used as a protective treatment when applied at regular intervals or as a knockdown treatment to control existing infestation.

MIXING

Shake or roll the container several times before measuring out product to ensure that the ingredients are well mixed. There is no need for pre-mixing. Add the product to the partly filled spray vat while agitating. Continue to agitate while topping up the tank and during spraying. Reseal partly used container immediately after use. If using as part of a tank mix always add Alpha Forte after the addition of any granular product. Always add Alpha Forte before paraquat based products.

APPLICATION

CONQUEST ALPHA FORTE 250 SC INSECTICIDE can be applied by ground or aircraft. Thorough coverage is essential to ensure adequate control. Always apply with a non-ionic surfactant unless detailed on the label of a tank mix partner. Apply during the cooler parts of the day or night.

Ground application –

For low volume spraying of field crops with ground rigs, use a total volume of 50-200 L/ha except for sweet corn, tomatoes and tobacco where higher volume should be used. Drop arms should be used on ground rigs in row crops taller than 30 cm (0.3 m). The application should be made as a fine spray, preferably using hollow cone nozzles, unless otherwise directed in the Critical Comments.

Aerial application –

Do NOT apply to trellis tomatoes by aircraft. Use a minimum spray volume of 20 L/ha. For spring/early summer application to cereals, linola, canola, rice and to other dense crops, apply in a total spray volume of 30 to 35 L/ha. If possible, spray in a crosswind. Avoid spraying in calm conditions or when wind is light and variable in direction. Apply as a spray of 100-150 microns VMD.

GROUP	3 A	INSECTICIDE
-------	------------	-------------

INSECTICIDE RESISTANCE WARNING

For insecticide resistance management CONQUEST ALPHA FORTE 250 SC INSECTICIDE is a Group 3A insecticide. Some naturally occurring insect biotypes resistant to CONQUEST ALPHA FORTE 250 SC INSECTICIDE and other Group 3A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if CONQUEST ALPHA FORTE 250 SC INSECTICIDE or other Group 3A insecticides are used repeatedly. The effectiveness of CONQUEST ALPHA FORTE 250 SC INSECTICIDE on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Conquest Crop Protection Pty Ltd accepts no liability for any losses that may result from the failure of CONQUEST ALPHA FORTE 250 SC INSECTICIDE to control resistant insects. CONQUEST ALPHA FORTE 250 SC INSECTICIDE may be subject to specific resistance management

strategies. For further information, contact your local supplier, Conquest Crop Protection Pty Ltd representative or local agricultural department agronomist. In NSW and Qld, application of this product to *Helicoverpa armigera* larvae longer than 5 mm may not only be ineffective but it may increase the level of synthetic pyrethroid resistance. This product should NOT be used to treat infestations that were not controlled by an earlier application of it or another synthetic pyrethroid. Infestations not controlled by this product should be treated with an insecticide from another chemical group. Application of this product with an insecticide from another chemical group such as NUDRIN® will assist with the management of synthetic pyrethroid resistant *Helicoverpa armigera*.

PRECAUTION

Do NOT use human flaggers unless they are protected by engineering controls such as vehicles with cabs.

RE-ENTRY PERIOD

Do NOT allow entry into treated areas for 12 hours after application. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist, a washable hat and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF LIVESTOCK

Dangerous to bees. Do NOT spray on any plants in flower while bees are foraging. CONQUEST ALPHA FORTE 250 SC INSECTICIDE is known to have a deterrent effect on foraging bees for a short period of time after spraying. Risk to bees is reduced by spraying in early morning and late evening while bees are not foraging.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish and aquatic invertebrates. Do NOT contaminate fish ponds, dams, drains, rivers or streams with product or used containers. Drift and run-off from treated areas may be hazardous to fish in adjacent sites.

STORAGE AND DISPOSAL

Store in the closed, original container, in a dry, cool, well-ventilated area out of direct sunlight.

Do NOT dispose of unused chemicals on-site.

The method of disposal of the container depends on the container type. Read the "Storage and Disposal" instructions on the label that is attached to the container.

SAFETY DIRECTIONS

Harmful if inhaled. Will irritate the eyes and skin. Facial skin contact may cause temporary facial numbness. Avoid contact with eyes and skin. Do NOT inhale spray mist. When opening the container, preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Telephone 13 11 26 Australia-wide.

MSDS

Additional information is listed in the Material Safety Data Sheet. A copy of the MSDS is available from our website: www.conquestag.com.au

CONDITIONS OF SALE

Conquest Crop Protection Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

Batch No:

DOM:

APVMA Approval No: 65245/1021102