

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Custodia® 320 SC Fungicide

ACTIVE CONSTITUENTS:

200 g/L TEBUCONAZOLE

120 g/L AZOXYSTROBIN

GROUP **3|11** FUNGICIDE

Crops: Almonds and Grapevines

Uses: Management of foliar diseases specified
in the Directions for Use Table



ADAMA

adama.com

CONTENTS: 1 L - 100 L

RESTRAINTS

DO NOT apply if heavy rain or storms that are likely to cause runoff is forecast within 3 days.

SPRAY DRIFT RESTRAINTS

DO NOT apply when wind speed is less than 3 or more than 20 kilometres per hour as measured at the application site.

DO NOT apply during surface temperature inversion conditions at the application site.

Users of this product **MUST make an accurate written record** of the details of each spray application within 24 hours following application, and must **KEEP** this record for at least 2 years. The spray application details that must be recorded are:

1) Date with start and finish times of application; 2) Location address and paddock(s) sprayed; 3) Full name of this product; 4) Amount of product used per hectare and number of hectares applied to; 5) Crop or situation and pest; 6) Wind speed and direction during application; 7) Air temperature and relative humidity during application; 8) Nozzle brand, type, spray angle, nozzle capacity and spray system pressure measured during application; 9) Name and address of person applying this product. (Additional record details may be required by the state or territory where this product is used.)

MANDATORY NO-SPRAY ZONES

Aquatic Environment

DO NOT apply if there are aquatic or wetland areas including aquaculture ponds downwind from the application area and within the **mandatory no-spray zones** shown in the table below.

No-Spray Zones for Protection of the Aquatic Environment - Almonds and Grapevines

FOR GROUND APPLICATION	
Wind Speed Range at Time of Application	Downwind Mandatory No-Spray Zone
From 3 to 20 kilometres per hour	5 metres

DIRECTIONS FOR USE

Table 1. Almonds

DISEASE	RATE	WHP	CRITICAL COMMENTS
Brown rot / Blossom blight (<i>Monilinia laxa</i>) Shot-hole (<i>Wilsonomyces carpophilus</i>) Rust (<i>Tranzschelia discolor</i>)	Dilute spraying: 60 mL/100 L Concentrate spraying: 1 L/ha	H 4 weeks	Application rate and spray volume: Use the 60 mL/100 L rate in up to 1600 litres/ha water. If using higher application volumes, dilute accordingly. Do not exceed a total use rate of 1 L per ha. Brown rot / Blossom blight and Shot-hole: Brown rot/ blossom blight and Shot-hole should be managed from budswell using a season-long spray program including fungicides from different mode of action groups. Apply CUSTODIA® 320 SC from early (10%) bloom. Subsequent applications of CUSTODIA® 320 SC and other mode of action fungicides should be applied on a 7 to 14 day interval, targeting key growth stages including full bloom, petal fall and shuck fall. Post-shuck fall, apply fungicides registered for control of Shot-hole such as CAVALRY® WEATHERGUARD on 14 day spray intervals. Rust: Apply up to two applications of CUSTODIA® 320 SC per season as part of a disease management spray program. Commence applications for rust from late flowering and/or when the first new leaves emerge. Apply CUSTODIA® 320 SC prior to disease development and in rotation with fungicides from other mode of action groups on a 10-14 day spray interval. This use is subject to a CropLife Fungicide Resistance Management Strategy: Apply CUSTODIA® 320 SC as part of a season long spray program incorporating fungicides from other mode of action groups. DO NOT apply more than two sprays of Group 11 fungicides per season including CUSTODIA® 320 SC. DO NOT apply more than four sprays of Group 3 fungicides per season including CUSTODIA® 320 SC. DO NOT apply more than two consecutive applications of products containing Group 3 fungicides.

H = harvest

Table 2. Grapevines.

DISEASE	RATE	WHP	CRITICAL COMMENTS
Powdery Mildew (<i>Uncinular necator</i>)	Dilute spraying: 65 or 100 mL/100 L Concentrate spraying: 1 L/ha	H 4 weeks	Apply CUSTODIA® 320 SC as part of a season-long spray program targeting the following growth stages: 1. When shoots are 20 cm long. 2. Pre-flowering. 3. Flowering. 4. After fruit set. 5. Before bunches close. Use the lower rate prior to flowering (before E-L 19) or under low disease pressure. Apply the higher rate from the start of flowering (from E-L 19) and under conditions of high disease pressure. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. DO NOT apply in volumes less than 250 L/ha. DO NOT apply at concentrations greater than 4X maximum label rate. This use is subject to a CropLife Fungicide Resistance Management Strategy: DO NOT apply more than two sprays of CUSTODIA® 320 SC per season. DO NOT use CUSTODIA® 320 SC curatively.
Downy Mildew (<i>Plasmopara viticola</i>)	Dilute spraying: 100 mL/100 L Concentrate spraying: 1 L/ha		Apply CUSTODIA® 320 SC as part of a seasonal preventative spray program commencing when shoots are approximately 20 cm long. Use CUSTODIA® 320 SC in rotation with other mode of action fungicides on a 7-21 day spray interval. Under conditions of high disease pressure and/or at critical timings i.e. during flowering; ensure that CUSTODIA® 320 SC is applied in rotation with fungicides from a different mode of action group such as AXIOM® PLUS.
Botrytis Bunch Rot (<i>Botrytis cinerea</i>)			When CUSTODIA® 320 SC is used in a seasonal spray programme it will provide control of Botrytis additional to that of specific botryticides. Under conditions of high disease pressure, CUSTODIA® 320 SC should not be used for Botrytis control particularly at critical timings such as 80 to 100% capfall and pre-harvest. This use is subject to a CropLife Fungicide Resistance Management Strategy: Apply CUSTODIA® 320 SC preventatively. DO NOT apply more than two sprays per season of CUSTODIA® 320 SC or any other Group 11 fungicide. If three or fewer Botrytis sprays are applied in a season, use no more than one spray of CUSTODIA® 320 SC during the season. If four or more Botrytis sprays are to be applied in a season, up to two CUSTODIA® 320 SC sprays can be applied per season.

H = harvest

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

ALMONDS AND GRAPEVINES: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

MIXING

Prior to pouring, shake container vigorously, then add the required quantity of CUSTODIA® 320 SC to water in the spray vat while stirring or with agitators in motion. Rinse the empty container with water and add rinsings to spray tank.

APPLICATION

Foliar applications for Botrytis, Powdery and Downy mildew in Grapevines and diseases in Almonds

Dilute Spraying

Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop canopy size being sprayed. Set up and operate the sprayer to achieve even coverage throughout the canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Directions for Use table for each 100 L of water and modify the concentration of CUSTODIA® 320 SC in the spray solutions according to the following guidelines:

1. **Grapevines:** If greater than 1000 L/ha is required to achieve the point of run-off, adjust the amount of product added for each 100 L of water to **ensure that no more than 1.0 L/ha product is applied.**
2. **Almonds:** If greater than 1600 L/ha is required to achieve the point of run-off, adjust the amount of product added for each 100 L of water to **ensure that no more than 1.0 L/ha product is applied.**

The required dilute spray volume will change, and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying

Use a sprayer designed and set up for the concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate can then be calculated in the following way:

EXAMPLE ONLY

1. Dilute spray volume as determined above: For example 1000 L/ha.
2. Your chosen concentrate spray volume: For example 500 L/ha.
3. The concentration factor in this example is: $2x$ (i.e. $1000 L \div 500 L = 2$).
4. If the dilute label rate is 100 mL/100 L, then the concentrate rate becomes 2×100 , that is 200 mL/100 L of concentrate spraying.

The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows. DO NOT use a concentrate rate higher than that specified in the Critical Comments.

For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

FUNGICIDE RESISTANCE WARNING

CUSTODIA® 320 SC is a member of the DMI and quinone outside inhibitors (Qols) group of fungicides. For fungicide resistance management the product is both a Group 3 and a Group 11 fungicide. Some naturally occurring individual fungi resistant to CUSTODIA® 320 SC and other Group 3 and/or Group 11 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product and other Group 3 and/or Group 11 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, ADAMA Australia Pty. Ltd. accepts no liability for any losses that may result from failure of CUSTODIA® 320 SC to control resistant fungi.

GROUP **311** FUNGICIDE

PRECAUTION

Re-entry period

DO NOT enter treated areas until the spray has dried unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

WARNING

- CUSTODIA® 320SC is extremely phytotoxic to certain apple varieties. Extreme care must be used to prevent injury to apple trees.
- AVOID SPRAY DRIFT. DO NOT spray CUSTODIA® 320SC where spray drift may reach apple trees.
- DO NOT spray when conditions favour drift beyond the area intended for application. Conditions that may contribute to drift include thermal inversions, excessive wind speed, certain sprayer nozzle/pressure combinations, small spray droplet size etc.
- DO NOT use spray equipment to spray apple trees that has been previously used to apply CUSTODIA® 320SC. Even trace amounts can cause unacceptable phytotoxicity.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. DO NOT contaminate dams, waterways or drains with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Harmful if swallowed. May irritate the eyes. Avoid contact with eyes. When opening the container, preparing and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. When opening the container and preparing the product for aerial spraying equipment, wear cotton overalls over normal clothing, buttoned to the neck and wrist and elbow-length chemical resistant gloves. When preparing the product and using the prepared product for airblast equipment, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

The above Safety Directions should be combined with the following engineering controls:

- Airblast application MUST BE carried out with enclosed cab equipment.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131 126.

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for CUSTODIA® 320 SC Fungicide is available from adama.com or call Customer Service on 1800 423 262.

CONDITIONS OF SALE: The use of CUSTODIA® 320 SC Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Adama Australia Pty. Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Adama Australia Pty. Ltd. accepts no responsibility for any consequence whatsoever from the use of this product.

® Registered trademark of an ADAMA Group Company

05/2016

NOT A DANGEROUS GOOD ACCORDING TO THE AUSTRALIAN DANGEROUS GOODS (ADG) CODE.