

MAIN PANEL (for 397/175 solvent mix)

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



RLP  
Approved



**Jaguar<sup>®</sup>**  
SELECTIVE HERBICIDE



**ACTIVE CONSTITUENTS:** 250 g/L BROMOXYNIL present as the octanoate  
25 g/L DIFLUFENICAN

**SOLVENTS:** 397 g/L LIQUID HYDROCARBONS  
175 g/L N-METHYL-2-PYRROLIDONE

**GROUP C F HERBICIDE**

\* L

**IMPORTANT: READ THE  
ATTACHED BOOKLET  
BEFORE USE**

For the control of certain broadleaf weeds in winter cereals and pasture as specified in DIRECTIONS FOR USE Table

\* 5 L, 10 L, 20 L, 110 L or 1000 L

(label code)

## JAGUAR SELECTIVE HERBICIDE

### STORAGE AND DISPOSAL (5 L, 10 L and 20 L containers)

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. Do not re-use empty containers for any other purpose.

### STORAGE AND DISPOSAL (110 L returnable containers)

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured. Empty container by pumping through dry-break connection system. Do not attempt to breach the valve system or the filling point, or contaminate the container with water or other products. Ensure that the coupler, pump, meter and hoses are disconnected, triple rinsed with clean water and drained after each use. When empty, or contents no longer required, return the container to the point of purchase. This container remains the property of Bayer CropScience Pty Ltd.

### STORAGE AND DISPOSAL (1000 L containers)

If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured. The container must be vented before discharging contents. To empty, connect a camlock fitted hose to the bottom valve. Remove top cap when discharging for venting purposes. When the container is empty, close all caps and valves and return the container to the point of purchase.

## SAFETY DIRECTIONS

Product is harmful if inhaled or swallowed. Will irritate eyes, nose, throat and skin. Avoid inhaling spray mist. When preparing spray wear elbow length PVC gloves and face-shield. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

## FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (telephone 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

## MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from [www.bayercropscience.com.au](http://www.bayercropscience.com.au).

## EXCLUSION OF LIABILITY

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

APVMA Approval No.: 40383/ 54907

Jaguar® is a Registered Trademark of Bayer.

**IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE**

\*

BAR  
CODE

FOR 24 HOUR SPECIALIST  
ADVICE  
IN EMERGENCY ONLY  
PHONE 1800 033 111

Bayer CropScience Pty Ltd  
ABN 87 000 226 022  
391-393 Tooronga Rd  
East Hawthorn Vic. 3123  
Phone: (03) 9248 6888  
Fax: (03) 9248 6800  
[www.bayercropscience.com.au](http://www.bayercropscience.com.au)  
Technical enquiries: 1800 804 479



Bayer CropScience

Batch Number:  
Date of Manufacture:

(label code)

---

\* drumMuster logo required for 5 L, 10 L and 20 L pack only

**POISON****KEEP OUT OF REACH OF CHILDREN****READ SAFETY DIRECTIONS BEFORE OPENING OR USING****JAGUAR SELECTIVE HERBICIDE**

ACTIVE CONSTITUENTS:	250 g/L BROMOXYNIL present as the octanoate 25 g/L DIFLUFENICAN
SOLVENTS:	397 g/L LIQUID HYDROCARBON 175 g/L N-METHYL-2-PYRROLIDONE

For the control of certain broadleaf weeds in winter cereals and pasture as specified in the DIRECTIONS FOR USE Table.

**STORAGE AND DISPOSAL (5 L, 10 L and 20 L containers)**

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. Do not re-use empty containers for any other purpose.

**STORAGE AND DISPOSAL (110 L returnable containers)**

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured.

Empty container by pumping through dry-break connection system. Do not attempt to breach the valve system or the filling point, or contaminate the container with water or other products.

Ensure that the coupler, pump, meter and hoses are disconnected, triple rinsed with clean water and drained after each use.

When empty, or contents no longer required, return the container to the point of purchase. This container remains the property of Bayer CropScience Pty Ltd.

**STORAGE AND DISPOSAL (1000 L containers)**

If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured. The container must be vented before discharging contents. To empty, connect a camlock fitted hose to the bottom valve. Remove top cap when discharging for venting purposes. When the container is empty, close all caps and valves and return the container to the point of purchase.

**SAFETY DIRECTIONS**

Product is harmful if inhaled or swallowed. Will irritate eyes, nose, throat and skin. Avoid inhaling spray mist. When preparing spray wear elbow length PVC gloves and face-shield. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

**FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre (telephone 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

LABEL BOOKLET - front page of booklet - continued

(397/175 solvent mix)

---

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet, which can be obtained from [www.bayercropscience.com.au](http://www.bayercropscience.com.au).

**EXCLUSION OF LIABILITY**

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

APVMA Approval No. 40383/ 54907

Jaguar® is a Registered Trademark of Bayer.

**IMPORTANT: READ THIS BOOKLET BEFORE USE**

## LABEL BOOKLET (continued)

<b>WEED (Common name)</b>	<b>WEEDS LIST (Scientific name)</b>	<b>PAGE NO.</b>
Amsinckia	<i>Amsinckia</i> spp.	X
Ball mustard	<i>Neslia paniculata</i>	X
Canola (rapeseed)	<i>Brassica napus</i>	X
Capeweed	<i>Arctotheca calendula</i>	X
Chamomile	<i>Matricaria matricarioides</i>	X
Charlock	<i>Sinapis arvensis</i>	X
Chickweed	<i>Stellaria media</i>	X
Cleavers	<i>Galium aparine</i>	X
Climbing buckwheat	<i>Fallopia convolvulus</i>	X
Common cotula (bird's eye)	<i>Cotula australis</i>	X
Common peppergrass	<i>Lepidium africanum</i>	X
Common sowthistle (milk thistle)	<i>Sonchus oleraceus</i>	X
Corn gromwell	<i>Buglossoides arvensis</i>	X
Crassula (stonecrop)	<i>Crassula</i> spp.	X
Deadnettle	<i>Lamium amplexicaule</i>	X
Dense-flower fumitory	<i>Fumaria densiflora</i>	X
Dock	<i>Rumex</i> spp.	X
Doublegee (spiny emex)	<i>Emex australis</i>	X
Fat hen	<i>Chenopodium album</i>	X
Field madder	<i>Sherardia arvensis</i>	X
Fireweed	<i>Senecio</i> spp.	X
Fumitory	<i>Fumaria</i> spp.	X
Hexham scent (King Island melilot)	<i>Melilotus indicus</i>	X
Horehound	<i>Marubium vulgare</i>	X
Lesser swinegrass	<i>Coronopus didymus</i>	X
Long storksbill	<i>Erodium botrys</i>	X
Marshmallow	<i>Malva parviflora</i>	X
Mexican poppy	<i>Argemone ochroleuca</i>	X
Mintweed	<i>Salvia reflexa</i>	X
Mouse-eared chickweed	<i>Cerastium glomeratum</i>	X
New Zealand spinach	<i>Tetragonia tetragonoides</i>	X
Ox-tongue	<i>Picris echioides</i>	X
Paterson's curse (Salvation Jane)	<i>Echium plantagineum</i>	X
Pheasants eye (adonis)	<i>Adonis dentatus</i>	X
Prickly lettuce	<i>Lactuca serriola</i>	X
Purple calandrinia (mountain sorrel)	<i>Calandrinia menziesii</i>	X
Rough poppy	<i>Papaver hybridum</i>	X
Saffron thistle	<i>Carthamus lanatus</i>	X
Scarlet pimpernel	<i>Anagallis arvensis</i>	X
Shepherd's purse	<i>Capsella bursa-pastoris</i>	X
Skeleton weed	<i>Chondrilla juncea</i>	X
Sorrel	<i>Rumex acetosella</i>	X
Speedwell	<i>Veronica</i> spp.	X
Spoon cudweed	<i>Stuartina muelleri</i>	X
Three-horned bedstraw	<i>Galium tricornutum</i>	X
Toad rush	<i>Juncus bufonius</i>	X
Tree hogweed	<i>Polygonum patulum</i>	X
Turnip weed	<i>Rapistrum rugosum</i>	X
Variiegated thistle	<i>Silybum marianum</i>	X
Vetch	<i>Vicia sativa</i>	X
Volunteer field peas	<i>Pisum sativum</i>	X
Volunteer lupins	<i>Lupinus angustifolius</i>	X
Ward's weed	<i>Carrichtera annua</i>	X
Wild mustard	<i>Sisymbrium</i> spp.	X
Wild radish	<i>Raphanus raphanistrum</i>	X
Wild turnip	<i>Brassica tournefortii</i>	X
Wireweed	<i>Polygonum aviculare</i>	X

## LABEL BOOKLET (continued)

**DIRECTIONS FOR USE****Restrictions**

DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

DO NOT apply to crops under stress due to disease or insect damage.

DO NOT apply to frost-affected crops or if frosts are imminent.

DO NOT apply if heavy rain is expected within 4 hours.

DO NOT apply with crop oils (cereals only).

CROP	WEEDS CONTROLLED	WEED STAGE	RATE/HA	STATE	CRITICAL COMMENTS
<b>Wheat, barley, triticale, cereal rye</b> (including undersown with clover and/or lucerne), and these cover crops in vineyards  <b>Pasture</b> Clover and/or lucerne-based pasture (newly sown or established) including cover crops in vineyards	Wild radish	Up to 2 leaf stage and not more than 60 mm in diameter and where weed density is less than 50 plants/m <sup>2</sup>	350 mL	WA only	<p><b>CROP STAGE:</b> <i>Cereals</i> 2 leaf to fully tillered (Zadok's Z12-29) Optimum results are achieved when sprayed at 4-8 weeks post-sowing. <b>Warning:</b> Jaguar may cause transient crop yellowing of cereals. (Refer to "Crop Tolerance" section of General Instructions).</p> <p><i>Clover and lucerne</i> Application is recommended prior to the 8th trifoliolate leaf stage. Application can be made from the 1st trifoliolate leaf stage in Qld, NSW, ACT and Vic only. In other States applications prior to the 3 leaf stage may result in crop damage if seedlings are under stress and in sandy soils. DO NOT apply to annual medics. <b>Warning:</b> Jaguar may affect growth and seed set of some varieties of clover and lucerne (Refer to "Crop Tolerance" section of General Instructions).</p> <p><b>COVER CROPS IN VINEYARDS:</b> When using in vineyard situations, apply during vine dormancy only. Contact with vines must be avoided. Particular care should be taken if applied in late autumn or early spring, when vines may not be fully dormant.</p> <p><b>WEED STAGE:</b> Apply from early post-emergence.</p> <p><b>APPLICATION:</b> Apply when weeds are actively growing. Ensure thorough coverage of weeds. Where crop or weed density is high, increase water volume.</p> <p><b>CRITICAL COMMENTS</b> CONTINUED ON NEXT PAGE</p>
	Wild mustard Wild radish	Up to 4 leaf stage and not more than 120 mm in diameter	500 mL	All States	
		Up to 6 leaf stage and not more than 150 mm in diameter	750 mL		
		Up to 8 leaf stage and not more than 180 mm in diameter	1.0 L		
	Canola (rapeseed) charlock turnip weed wild turnip.	Up to 2 leaf stage and not more than 60 mm in diameter	500 mL		
		Up to 4 leaf stage and not more than 120 mm in diameter	750 mL		
	Shepherd's purse	120 mm in diameter	1.0 L		
	Capeweed	Up to 4 leaf stage and not more than 120 mm in diameter	500 mL		
		Up to 6 leaf stage and not more than 150 mm in diameter	750 mL		
		Up to 8 leaf stage and not more than 180 mm in diameter	1.0 L		
	Corn gromwell	Up to 4 leaf stage	500 mL		
		Up to 6 leaf stage	750 mL		
	Climbing buckwheat	Up to 2 leaf stage	500 mL		
		Up to 4 leaf stage	750 mL		
		Up to 6 leaf stage	1.0 L		
Deadnettle, Paterson's curse, (Salvation Jane), rough poppy	Up to 2 leaf stage	500 mL			
	Up to 4 leaf stage	750 mL			
Amsinckia					
Doublegee (spiny emex)	Up to 2 leaf stage	500 mL	Qld, NSW, ACT, Vic, Tas, WA only		
	Up to 4 leaf stage	750 mL	All States		

## LABEL BOOKLET (continued)

## Directions For Use

CROP	WEEDS CONTROLLED	WEED STAGE	RATE/HA	STATE	CRITICAL COMMENTS	
Wheat, barley, triticale, cereal rye (including undersown with clover and/or lucerne), and these cover crops in vineyards  Pasture Clover and/or lucerne-based pasture (newly sown or established) including cover crops in vineyards	Chamomile, common peppergrass, lesser swinecress, purple calandrinia, (mountain sorrel), tree hogweed	Up to 4 leaf stage.	1.1 L	All States	<b>CRITICAL COMMENTS CONTINUED FROM PREVIOUS PAGE</b> In most situations the rate specified for each weed size will give satisfactory control. However, under certain conditions such as: * high crop and weed density, * late season germinations, * abnormal weed growth (including early flowering); higher rates of product (up to the maximum rate of application specified for that weed) may be required.	
	Fat hen, field madder, saffron thistle, variegated thistle		1.0 L			
	Ox-tongue, wireweed	Up to 2 leaf stage				
	Fireweed	Up to 4 leaf stage	500 mL	Qld, NSW, ACT, Vic, SA, WA, NT only		
	Common cotula (bird's eye) pheasants eye (adonis)	Up to 4 leaf stage Greater than 4 leaf stage	560 mL 1.1 L	SA only		
Wheat, barley, triticale, cereal Rye	Fumitory	2-6 leaf stage	350 + 200 mL/ha terbutryn (500 g/L)	WA only	Jaguar will not effectively control: * regrowth of suppressed weeds, * transplanted weeds, * regrowth from rhizomes or roots, * weeds growing under stress from previous herbicide applications. * Radish plants beyond rosette stage	
Wheat, barley, triticale, cereal rye (including undersown with clover and/or lucerne), and these cover crops in vineyards  Pasture Clover and/or lucerne-based pasture (newly sown or established) including cover crops in vineyards	<b>Suppression of the Following Weeds</b>					
	Dense-flower fumitory	Up to 2 leaf stage Up to 4 leaf stage	750 mL 1.0 L	All States	Effective residual activity of this product may be reduced where: * rates lower than 1.0 L/ha are used; * dry conditions prevail; * poor coverage of the soil surface is achieved; * crop is grown in non-wetting sand; * soils have a high content of clay or organic matter.  <b>WILD RADISH:</b> Effective residual activity of this product may be reduced where: * rates lower than 1.0 L/ha are used; * dry conditions prevail; * poor coverage of the soil surface is achieved; * crop is grown in non-wetting sand; * soils have a high content of clay or organic matter.  <b>VOLUNTEER LUPINS:</b> In some situations, the higher rate of 1.0 L/ha may be required to effectively suppress volunteer lupins at the 4 leaf stage.  # Jaguar will suppress seedling dock but will not suppress regrowth from transplanted roots.	
	Chickweed, common sowthistle, (milk thistle), dock#, hexham scent, (King Island melilot), prickly lettuce, scarlet pimpernel, skeleton weed, sorrel, speedwell, three-horned bedstraw, toad rush		500 mL -1.0 L			
	Volunteer lupins					
	Crassula (stonecrop)	Up to 5 leaf stage	500 mL			
	Long storksbill	Up to 4 leaf stage				
	Volunteer field peas	Up to 5 node stage	750 mL			
	Ward's weed	Up to 5 leaf stage	1.0 L			
	Vetch	Up to 2 leaf stage				
	Mouse-eared chickweed					NSW & ACT only
	Mexican poppy					Qld only
	Mintweed, spoon cudweed	Up to 4 leaf stage				NSW & ACT only
	New Zealand spinach	Up to 2 leaf stage	750 mL			Qld only
	Cleavers	Up to 1 whorl stage	1.0 L			SA only
	Ball mustard	Up to 4 leaf stage				
	Horehound	Pre-emergence				
Marshmallow	Up to 2 leaf stage					



## LABEL BOOKLET (continued)

CROP	WEED CONTROLLED	WEED STAGE	RATE/HA	STATE	CRITICAL COMMENTS
Wheat, barley, triticale, cereal rye	Wild radish	Up to the 4 leaf stage and not more than 120 mm in diameter	350 mL plus 200 mL MCPA LVE (500 g/L)	WA only	<p><b>Refer also to all Critical Comments for cereals above.</b></p> <p><b>DO NOT</b> use this tank-mix if cereals are undersown with lucerne or annual medics. <b>DO NOT</b> use this tank-mix in vineyards.</p> <p><u>Crop Stage</u> Jaguar 350 mL + MCPA LVE 200 mL: Apply from 3 leaf to fully tillered (Zadok's Z13 to Z30). Jaguar 500 mL + MCPA LVE 200 mL: Apply from 3 leaf to fully tillered (Zadok's Z13 to Z30). Jaguar 500 mL + MCPA LVE 400 mL: Apply from 5 leaf stage to fully tillered (Zadok's Z15 to Z30).</p> <p>Optimum results are achieved when sprayed at 4-8 weeks post sowing. <b>Warning:</b> Jaguar may cause transient crop yellowing of cereals. (Refer to "Crop Tolerance" section of <b>General Instructions</b>)</p> <p>Observe instructions also on MCPA LVE product label.</p>
		Up to the 6 leaf stage and not more than 150 mm in diameter	500 mL plus 200 mL MCPA LVE (500 g/L)	All States	
		Up to the 8 leaf stage and not more than 180 mm in diameter.	500 mL plus 400 mL MCPA LVE (500 g/L)		

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD****Harvest:****Cereals, Grapes – NOT REQUIRED WHEN USED AS DIRECTED****Grazing:****Pasture, Cereals – DO NOT GRAZE OR CUT FOR STOCK FOOD WITHIN 14 DAYS AFTER APPLICATION.****GENERAL INSTRUCTIONS:**

- This product is a post-emergence contact herbicide, which may provide residual control of wild radish up to 4 weeks after application.
- Apply Jaguar Selective Herbicide immediately after mixing. Do not allow to stand in the spray tank overnight.
- Optimum results will be obtained if good soil moisture exists at and after application and weeds are not stressed.
- Some pre-emergence herbicides, such as atrazine, can cause stress to certain crops resulting in an increase in crop damage when using this product. Crops which are particularly sensitive are lucerne and subterranean clover.

LABEL BOOKLET (continued)**Resistant Weeds Warning****GROUP C F HERBICIDE**

Jaguar Selective Herbicide is a member of the nitrile and nicotinamide groups of herbicides. Jaguar is an inhibitor of photosynthesis at photosystem II and carotenoid biosynthesis. For weed resistance management, Jaguar is a Group C, F herbicide. Some naturally occurring weed biotypes resistant to Jaguar and other Group C, F herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Jaguar or other Group C, F herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that may result from the failure of Jaguar to control resistant weeds.

**Temperature warning**

Do not apply Jaguar if frosts are imminent. Frost causes stress on crops and weeds and could result in increased crop effects and/or decreased weed control. To ensure good results Jaguar should only be applied once the weeds and crop are no longer under stress from the frost conditions.

Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application, as increased crop damage may result.

**CROP TOLERANCE*****Cereals***

After application some transient crop yellowing may occur. This usually appears as yellow or white banding on leaves. Provided the crop is not under stress from pre-emergent herbicide, root disease, insect damage, frost, dry or excessively moist conditions, the development of the crop and subsequent growth will be unaffected.

***Lucerne*****Warning**

The tolerance of lucerne varieties to Jaguar can vary with rate of application, soil type, crop health, stage of growth and degree of moisture and temperature stress. Jaguar may result in transient crop yellowing and suppression of growth with a resultant initial reduction in dry matter. For this reason we recommend application prior to the 8 trifoliate leaf stage. However, under normal growing conditions subsequent growth and seed yield should not be affected. Crop damage may be increased if rates higher than 500 mL/ha are used and in areas where spray overlapping has occurred.

Under normal growing conditions, the following lucerne varieties have shown acceptable levels of foliage tolerance to Jaguar applied at 500 mL/ha: Hunter River, Nova and Dekalb 185. Varieties not listed should be tested before using Jaguar over large areas.

Consult your local Bayer CropScience representative for advice on specific varieties.

***Subterranean clover*****Warning**

The tolerance of subterranean clover varieties to Jaguar can vary with rate of application, soil type, crop health, stage of growth and degree of moisture and temperature stress. Jaguar may result in transient crop yellowing and suppression of growth with an initial reduction in dry matter. For this reason we recommend application prior to the 8 trifoliate leaf stage. However, under normal growing conditions subsequent growth and seed yield should not be affected. Crop damage may be increased if rates higher than 500 mL/ha are used and in areas where spray overlapping has occurred.

Under normal growing conditions, the following varieties have shown acceptable levels of foliage tolerance to Jaguar applied at 500 mL/ha:

Daliak, Dalkeith, Denmark, Goulburn, Karridale, Leura, Mt. Barker, Nungarin, Rosedale, Seaton Park, Trikkala and Woogenellup.

The variety Junee has shown increased sensitivity to Jaguar so care should be taken if this variety is part of the pasture sward.

The effects of Jaguar on subterranean clover seed yield have been tested on the following varieties. Under normal growing conditions they show acceptable levels of tolerance to Jaguar applied at 500 mL/ha. However, higher rates may reduce seed yield under conditions of low weed pressure:

Denmark, Goulburn, Larissa, Nungarin, Seaton Park, Trikkala and Woogenellup.

Varieties not listed should be tested before using Jaguar over large areas. Consult your local Bayer CropScience representative for advice on specific varieties.

### **Other Clovers**

#### **Warning**

The tolerance of clover varieties to Jaguar can vary with rate of application, soil type, crop health, stage of growth and degree of moisture and temperature stress. Jaguar may result in transient crop yellowing and suppression of growth with a resultant initial reduction in dry matter. For this reason we recommend application prior to the 8 trifoliolate leaf stage. However, under normal growing conditions subsequent growth and seed yield should not be affected. Crop damage may be increased if rates higher than 500 mL/ha are used and in areas where spray overlapping has occurred. The effect on seed yield of other clovers has not been determined.

The following varieties of clover have shown increased sensitivity to Jaguar:

Big Bee, Sacromonte (Berseem), Haifa (White), Zulu (Arrowleaf), Kyambro, Lupers and Maral (Persian).

Care should be exercised if these clovers are part of the pasture sward.

Varieties not listed should be tested before using Jaguar over large areas.

Consult your local Bayer CropScience representative for advice on specific varieties.

### **Subsequent Crops**

To reduce effect on subsequent susceptible crops (e.g. canola), ensure thorough cultivation of soil prior to the sowing of these crops.

### **MIXING**

To ensure even mixing, half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly while carrying out spray operations. Reseal part-used container immediately after use.

### **APPLICATION**

#### **Boom Sprayer**

A minimum of 50 L water/ha should be used, however, for optimum results water rates of 70-100 L/ha are recommended. Increase the water volume where weed infestation is heavy or the crop cover is dense. Complete coverage of weeds is essential. Higher water volumes (up to 100 L/ha) will ensure faster activity of the product on the weeds but may increase the symptoms of crop damage.

## LABEL BOOKLET (continued)

The following settings are examples which will ensure excellent coverage of exposed weeds:

Water Rate	50 L/ha	75 L/ha	75 L/ha
Nozzle	Hardi No. 10 or equivalent	Hardi No. 12 or equivalent	Hardi No. 14 or equivalent
Speed	10 km/h	10 km/h	12 km/h
Pressure	240 kPa (2.4 bar)	220 kPa (2.2 bar)	210 kPa (2.1 bar)

**Controlled Droplet Application (CDA)**

Insufficient information is available to recommend the application of this product by CDA.

**Warning**

The rubber components present in some spraying units may be affected by exposure to the solvents in Jaguar. To reduce this risk it is recommended that the spray unit be thoroughly washed with a boom cleaner and fresh water after use.

**Aircraft**

Insufficient information is available to recommend the application of this product by air.

**COMPATIBILITY**

The following herbicide products are physically compatible with Jaguar as two-way mixtures in the spray tank, but should only be used for the crops specified, and only when the crop is also specified on the label of the compatible product:

(See below for list of compatible insecticides.)

Crop	Jaguar	Compatible Product
Wheat, triticale, cereal rye (including undersown)	Up to 750 mL/ha	Hoegrass <sup>®</sup> (barley also), Tristar <sup>®</sup> Advance (barley also), Wildcat <sup>®</sup> 110 EC (wild oats only, high rate)
Wheat, barley, triticale, cereal rye (including undersown)	All rates	Broadstrike <sup>®</sup>
Wheat, barley, triticale, cereal rye (not undersown)	Up to 500 mL/ha	Ally <sup>®</sup> , Glean <sup>®</sup> , MCPA LVE (500 g/L product) (up to 500 mL/ha only)
	All rates	2,4-D amine 500, Eclipse <sup>®</sup> , Cadence <sup>®</sup> (up to 115 g only), Lontrel <sup>®</sup>
Wheat only (not undersown)		Topik <sup>®</sup>
Established lucerne only	Up to 750 mL/ha	Simazine (500 g/L product) (up to 1.25 L/ha only) and simazine (500 g/L)/paraquat (200 g/L) mixture
Newly sown and established lucerne and clover only	Up to 750 mL/ha	Targa <sup>®</sup> , Fusilade <sup>®</sup> , 2,4-DB amine (500 g/L)
	Up to 1.0 L/ha	Broadstrike <sup>®</sup>

When mixing Jaguar with other herbicides, crop yellowing may be enhanced.

When mixing with Hoegrass<sup>®</sup>, Wildcat<sup>®</sup> 110 EC, Puma<sup>®</sup> Progress or Tristar<sup>®</sup> Advance some reduction in the efficacy and speed of action of these products may occur.

When mixing with Targa<sup>®</sup> or Fusilade<sup>®</sup> some reduction in the efficacy and speed of action of these products and Jaguar may occur.

In tank-mixtures with Ally<sup>®</sup> and Glean<sup>®</sup>, rates of Jaguar higher than 500 mL/ha may cause significant crop damage.

If the crop is stressed, the application of the herbicide tank-mixtures may cause yield reduction.

When mixing with Cadence<sup>®</sup> a temporary wilting may be evident in some crops after application.

The mixture of Jaguar and simazine should be applied during winter to lucerne which is not actively growing. This mixture may result in an increased crop effect but this can be reduced if the lucerne is grazed or cut before spraying.

DO NOT mix Jaguar with Verdict<sup>®</sup>.

Growers should seek advice before spraying recently released cereal varieties.

This product may be mixed in the spray tank with one of the following insecticides according to the directions for the insecticide product: Chlorpyrifos (500 g/L product), Decis Options<sup>®</sup>, dimethoate, Dominex<sup>®</sup>100EC, Fastac<sup>®</sup> Duo, Le-mat<sup>®</sup> 290 SL, Talstar<sup>®</sup> and Thiodan<sup>®</sup>.

Use the recommended rates for Jaguar and its tank-mix partner as well as the surfactant recommendation of the tank-mix partner. Read the label of the tank-mix partner before mixing and using the tank mixture. If another herbicide is applied as a tank mix, observe the plantback restrictions on that label.

### **Warning**

DO NOT use crop oils with Jaguar or Jaguar tank mixtures in cereals.

As formulations of other manufacturers' products are beyond the control of Bayer CropScience, all mixtures should be tested prior to mixing commercial quantities.

### **PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Wash sprayer thoroughly after use.

### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

Decis Options<sup>®</sup>, Eclipse<sup>®</sup>, Hoegrass<sup>®</sup>, Jaguar<sup>®</sup>, Puma<sup>®</sup>, Thiodan<sup>®</sup>, Tristar<sup>®</sup> and Wildcat<sup>®</sup> are Registered Trademarks of Bayer.

Bayer CropScience Pty Ltd  
ABN 87 000 226 022  
391-393 Tooronga Rd  
East Hawthorn Vic. 3123  
Phone: (03) 9248 6888  
Fax: (03) 9248 6800  
Website: [www.bayercropscience.com.au](http://www.bayercropscience.com.au)  
Technical enquiries: 1800 804 479



**Bayer CropScience**