

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

## Ospray Nimble®

### HERBICIDE

ACTIVE CONSTITUENTS: 250g/L BROMOXYNIL present as the octanoate  
25g/L DIFLUFENICAN

SOLVENTS: 405 g/L LIQUID HYDROCARBONS  
175 g/L N-METHYL-PYRROLIDONE

GROUP **CIF** HERBICIDES

For the control of certain broadleaf weeds in winter cereals and pasture  
as specified in the Directions for Use table.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT.**



0268wm



Ospray Pty Ltd ABN 23 110 199 169

50 Middleton Road  
Cottonvale QLD 4375 AUSTRALIA

Phone: 1800 624 597

[www.ospray.com.au](http://www.ospray.com.au)

Part of the  CHEMINOVA Group

## Contents 5L

# Ospray Nimble®

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

Keep from contact with fertilisers, insecticides, fungicides and seeds.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## SAFETY DIRECTIONS

Product is harmful if inhaled or swallowed. Will irritate the eyes, nose, throat and skin. Avoid inhaling spray mist. When preparing spray wear elbow length PVC gloves and face shield. If product in eyes wash out immediately with water. After use and before eating drinking or smoking, wash hand arms and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

## MATERIAL SAFETY DATA SHEET

For further information, refer to the Material Safety Data Sheet (MSDS) which is available from the supplier.

## MANUFACTURERS WARRANTY AND EXCLUSION OF LIABILITY

Ospray Pty Ltd has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Ospray Pty Ltd accepts no responsibility for any loss arising from incorrect storage, handling or use.

**IN A TRANSPORT EMERGENCY**

**DIAL 000**

**POLICE OR FIRE BRIGADE**

® - Registered Trademark of Ospray Pty Ltd

APVMA approval No: 62887/5L/0109

Batch No:

Date of Manufacture:



9 335944 003437

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**Ospray**  
**Nimble®**

## HERBICIDE

ACTIVE CONSTITUENTS: 250g/L BROMOXYNIL  
present as the octanoate  
25g/L DIFLUFENICAN

SOLVENTS: 405 g/L LIQUID HYDROCARBONS  
175 g/L N-METHYL-PYRROLIDONE

GROUP **CIF** HERBICIDES

For the control of certain broadleaf weeds in winter cereals and pasture as specified in the Directions for Use table.

**IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT**

APVMA Approval No. :  
62887/0109



Ospray Pty Ltd ABN 23 110 199 169  
50 Middleton Road  
Cottonvale QLD 4375 AUSTRALIA  
Phone: 1800 624 597  
www.ospray.com.au  
Part of the **CHEMINOVA** Group

0268wm

## DIRECTIONS FOR USE

Restrictions

DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

DO NOT apply to crops under stress due to disease or insect damage.

DO NOT apply to frost-affected crops or if frosts are imminent.

DO NOT apply if heavy rain is expected within 4 hours.

DO NOT apply with crop oils (cereals only).

CROP	WEEDS CONTROLLED	WEED STAGE	RATE/HA	STATE	CRITICAL COMMENTS
<b>Wheat, barley, triticale, cereal rye</b> (including undersown with clover and/or Lucerne), and these <b>cover crops in vineyards</b> .	Wild radish	Up to 2 leaf stage and not more than 60 mm in diameter and where weed density is less than 50 plants/m <sup>2</sup>	350 mL	WA only	<b>CROP STAGE:</b> <b>Cereals</b> 2 leaf to fully tillered (Zadok's Z12-29) Optimum results are achieved when sprayed at 4-8 weeds post-sowing. <b>Warning:</b> Nimble may cause transient crop yellowing of cereals. (Refer to <b>"Crop Tolerance"</b> section of <b>General Instructions</b> ). <b>Clover and Lucerne</b> Application is recommended prior to the 8th trifoliate leaf stage. Application can be made from the 1st trifoliate leaf stage in Qld, NSW, ACT and Vic only. In other States applications prior to the 3 leaf stage may result in crop damage if seedlings are under stress and in sandy soils. DO NOT apply to annual medics. <b>Warning:</b> Nimble may affect growth and seed set of some varieties of clover and Lucerne. (Refer to <b>"Crop Tolerance"</b> section of <b>General Instructions</b> ). <b>COVER CROPS IN VINEYARDS:</b> <b>When using in vineyard situations, apply during vine dormancy only.</b> <b>Contact with vines must be avoided. Particular care should be taken if applied in late autumn or early spring, when vines may not be fully dormant.</b> <b>WEED STAGE:</b> Apply from early post-emergence. <b>APPLICATION:</b> Apply when weeds are actively growing. Ensure thorough coverage of weeds. Where crop or weed density is high, increase water volume. In most situations the rate specified for each weed size will give satisfactory control. However, under certain conditions such as: <ul style="list-style-type: none"> <li>high crop and weed density,</li> <li>late season germinations,</li> <li>abnormal weed growth (including early flowering);</li> </ul> higher rates of product (up to the maximum rate of application specified for that weed) may be required.
	Wild mustard Wild radish	Up to 4 leaf stage and not more than 120 mm in diameter	500 mL	All States	
		Up to 6 leaf stage and not more than 150 mm in diameter	750 mL		
		Up to 8 leaf stage and not more than 180 mm in diameter	1.0 L		
	Canola (rapeseed) charlock turnip weed wild turnip Shepherd's purse Capeweed	Up to 2 leaf stage and not more than 60 mm in diameter	500 mL	All States	
		Up to 4 leaf stage and not more than 120 mm in diameter	750 mL		
		Up to 6 leaf stage and not more than 150 mm in diameter	750 mL		
		Up to 8 leaf stage and not more than 180 mm in diameter	1.0 L		
	Corn gromwell	Up to 4 leaf stage	500 mL	All States	
		Up to 6 leaf stage	750 mL		
Climbing buckwheat	Up to 2 leaf stage	500 mL	All States		
	Up to 4 leaf stage	750 mL			
	Up to 6 leaf stage	1.0 L			
Deadnettle, Paterson's curse (Salvation Jane), rough poppy Amsinckia	Up to 2 leaf stage	500 mL	All States		
	Up to 4 leaf stage	750 mL			
Doublegee (spiny emex)	Up to 2 leaf stage	500 mL	Qld, NSW, ACT, Vic, Tas, WA only		
	Up to 4 leaf stage	750 mL	All States		

CROP	WEEDS CONTROLLED	WEED STAGE	RATE/HA	STATE	CRITICAL COMMENTS (Con't)	
<b>Wheat, barley, triticale, cereal rye</b> (including undersown with clover and/or Lucerne), and these <b>cover crops in vineyards</b>	Chamomile, common peppercress, lesser swinecress, purple calandrinia (mountain sorrel), tree hogweed	Up to 4 leaf stage.	1.1 L	All States	Nimble will not effectively control: <ul style="list-style-type: none"> <li>regrowth of suppressed weeds,</li> <li>transplanted weeds,</li> <li>regrowth from rhizomes or roots,</li> <li>weeds growing under stress from previous herbicide applications.</li> <li>Radish plants beyond rosette stage</li> </ul> <b>WILD RADISH:</b> Effective residual activity of this product may be reduced where: <ul style="list-style-type: none"> <li>rates lower than 1.0 L/ha are used;</li> <li>dry conditions prevail;</li> <li>poor coverage of the soil surface is achieved;</li> <li>crop is grown in non-wetting sand;</li> <li>soils have a high content of clay or organic matter.</li> </ul> <b>VOLUNTEER LUPINS:</b> In some situations, the higher rate of 1.0L/ha may be required to effectively suppress volunteer lupins at the 4 leaf stage.	
	Fat hen, field madder, saffron thistle, variegated thistle		1.0 L			
	<b>Pasture</b> Clover and/or Lucerne based pasture (newly sown or established) including <b>cover crops in vineyards</b>	Ox-tongue, wireweed Fireweed	Up to 2 leaf stage Up to 4 leaf stage	500 mL		Qld, NSW, ACT, Vic, SA, WA, NT only SA only
<b>Wheat, barley, triticale, cereal rye</b>	Common cotula (bird's eye) Pheasants eye (Adonis)	Up to 4 leaf stage Greater than 4 leaf stage	560 mL 1.1 L	WA only		
	Fumitory	2-6 leaf stage	350+200 mL/ha terbutryn (500 g/L)			
<b>Wheat, barley, triticale, cereal rye</b> (including undersown with clover and/or Lucerne), and these <b>cover crops in vineyards</b>	<b>Suppression of the following Weeds</b>					
	Dense-flower fumitory	Up to 2 leaf stage	750 mL	All States	#Nimble will suppress seedling dock but will not suppress regrowth from transplanted roots.	
		Up to 4 leaf stage	1.0 L			
	Chickweed, common sowthistle (milk thistle), dock#, hexham scent (King Island melilot), prickly lettuce, scarlet pimpernel, skeleton weed, sorrel, speedwell, three-horned bedstraw, toad rush Volunteer lupins	Up to 4 leaf stage	500mL-1.0L			
			500mL			
			500mL			
Crassula (stonecrop) Long storksbill	Up to 5 leaf stage Up to 4 leaf stage	500mL				
<b>Wheat, barley, triticale, cereal rye</b> (including undersown with clover and/or lucerne), and these <b>cover crops in vineyards</b>	<b>Suppression of the following Weeds</b>					
	Volunteer field peas Ward's weed Vetch Mouse-eared chickweed	Up to 5 node stage Up to 5 leaf stage Up to 2 leaf stage	750 mL 1.0 L	All States	(refer Critical Comments above)	
	Mexican poppy	Up to 4 leaf stage	NSW & ACT only			
			Qld only			
	Mintweed, spoon cudweed New Zealand spinach Cleavers	Up to 4 leaf stage Up to 2 leaf stage Up to 1 whorl stage	750 mL 1.0 L			Qld only SA only
	Ball mustard Horehound Marshmallow	Pre-emergence Up to 2 leaf stage	SA only			
SA only						
<b>Wheat, barley, triticale, cereal rye</b>	Wild radish	Up to the 4 leaf stage and not more than 120 mm in diameter	350mL plus 200mL MCPA LVE (500g/L)	WA only	<b>Refer also to all Critical Comments for cereals above.</b> <b>DO NOT</b> use this tank-mix if cereals are undersown with lucerne or annual medics. <b>DO NOT</b> use this tank-mix in vineyards <b>Crop Stage</b> Nimble 350 mL + MCPA LVE 200 mL: Apply from 3 leaf to fully tillered (Zadok's Z13 to Z30). Nimble 500 mL + MCPA LVE 200 mL: Apply from 3 leaf to fully tillered (Zadok's Z13 to Z30). Nimble 500 mL + MCPA LVE 400 mL: Apply from 5 leaf to fully tillered (Zadok's Z15 to Z30). Optimum results are achieved when sprayed at 4-8 weeks post sowing. <b>Warning:</b> Nimble may cause transient crop yellowing of cereals. (Refer to <b>"Crop Tolerance"</b> section of <b>General Instructions</b> ) Observe instructions also on MCPA LVE product label.	
		Up to the 6 leaf stage and not more than 150 mm in diameter	500mL plus 200mL MCPA LVE (500g/L)	All States		
		Up to the 8 leaf stage and not more than 180 mm in diameter	500mL plus 400mL MCPA LVE (500g/L)	All States		

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD**

**Harvest:**

**Cereals, Grapes – NOT REQUIRED WHEN USED AS DIRECTED**

**Grazing:**

**Pasture, Cereals – DO NOT GRAZE OR CUT FOR STOCK FOOD WITHIN 14 DAYS AFTER APPLICATION**

## GENERAL INSTRUCTIONS

This product is a post-emergence contact herbicide, which may provide residual control of wild radish up to 4 weeks after application.

Apply Nimble Herbicide immediately after mixing. Do not allow to stand in the spray tank overnight.

Optimum results will be obtained if good soil moisture exists at and after application and weeds are not stressed.

Some pre-emergence herbicides, such as atrazine, can cause stress to certain crops resulting in an increase in crop damage when using this product. Crops which are particularly sensitive are lucerne and subterranean clover.

### Resistant Weeds Warning

<b>GROUP</b>	<b>CIF</b>	<b>HERBICIDES</b>
--------------	------------	-------------------

Nimble Herbicide is a member of the nitrile and nicotinilide groups of herbicides. Nimble is an inhibitor of photosynthesis at photosystem II and carotenoid biosynthesis. For weed resistance management, Nimble is a Group **C, F** herbicide. Some naturally occurring weed biotypes resistant to Nimble and other Group **C, F** herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Nimble or other Group **C, F** herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Ospray Pty Ltd accepts no liability for any losses that may result from the failure of Nimble to control resistant weeds.

### Temperature warning

Do not apply Nimble if frosts are imminent. Frost causes stress on crops and weeds and could result in increased crop effects and/or decreased weed control. To ensure good results Nimble should only be applied once the weeds and crop are no longer under stress from the frost conditions.

Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application, as increased crop damage may result.

### CROP TOLERANCE

#### *Cereals*

After application some transient crop yellowing may occur. This usually appears as yellow or white banding on leaves. Provided the crop is not under stress from pre-emergent herbicide, root disease, insect damage, frost, dry or excessively moist conditions, the development of the crop and subsequent growth will be unaffected.

#### *Lucerne*

#### Warning

The tolerance of lucerne varieties to Nimble can vary with rate of application, soil type, crop health, stage of growth and degree of moisture and temperature stress. Nimble may result in transient crop yellowing and suppression of growth with a resultant initial reduction in dry matter. For this reason we recommend application prior to the 8 trifoliolate leaf stage. However, under normal growing conditions subsequent growth and seed yield should not be affected. Crop damage may be increased if rates higher than 500mL/ha are used and in areas where spray overlapping has occurred.

Under normal growing conditions, the following lucerne have shown acceptable levels of foliage tolerance to Nimble applied at 400 mL/ha:

Hunter River, Nova and Dekalb 185.

Varieties not listed should be tested before using Nimble over large areas.

Consult your local Ospray representative for advice on specific varieties.

#### *Subterranean clover*

#### Warning

The tolerance of subterranean clover varieties to Nimble can vary with rate of application, soil type, crop health, stage of growth and degree of moisture and temperature stress. Nimble may result in transient crop yellowing and suppression of growth with an initial reduction in dry matter. For this reason we recommend application prior to the 8 trifoliolate leaf stage. However, under normal growing conditions subsequent growth and seed yield should not be affected. Crop damage may be increased if rates higher than 500 mL/ha are used and in areas where spray overlapping has occurred.

Under normal growing conditions, the following varieties have shown acceptable levels of foliage tolerance to Nimble applied at 500 mL/ha:

Daliak, Dalkeith, Denmark, Goulburn, Karridale, Leura, Mt. Barker, Nungarin, Rosedale, Seaton Park, Trikkala and Woogenellup.

The variety Junee has shown increased sensitivity to Nimble so care should be taken if this variety is part of the pasture sward.

The effects of Nimble on subterranean clover seed yield have been tested on the following varieties. Under normal growing conditions they show acceptable levels of tolerance to Nimble applied at 500 mL/ha. However, higher rates may reduce seed yield under conditions of low weed pressure:

Denmark, Goulburn, Larissa, Nungarin, Seaton Park, Trikkala and Woogenellup.

Varieties not listed should be tested before using Nimble over large areas.

Consult your local Ospray representative for advice on specific varieties.

#### *Other Clovers*

#### Warning

The tolerance of clover varieties to Nimble can vary with rate of application, soil type, crop health, stage of growth and degree of moisture and temperature stress. Nimble may result in transient crop yellowing and suppression of growth with a resultant initial reduction in dry matter. For this reason we recommend application prior to the 8 trifoliolate leaf stage. However, under normal growing conditions subsequent growth and seed yield should not be affected. Crop damage may be increased if rates higher than 500 mL/ha are used and in areas where spray overlapping has occurred. The effect on seed yield of other clovers has not been determined.

The following varieties of clover have shown increased sensitivity to Nimble:

Big Bee, Sacromonte (Berseem), Haifa (White), Zulu (Arrowleaf), Kyambro, Lupers and Maral (Persian).

Care should be exercised if these clovers are part of the pasture sward.

Varieties not listed should be tested before using Nimble over large areas.

Consult your local Ospray representative for advice on specific varieties.

#### *Subsequent Crops*

To reduce effect on subsequent susceptible crops (e.g. canola), ensure thorough cultivation of soil prior to the sowing of these crops.

#### MIXING

To ensure even mixing, half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly while carrying out spray operations. Reseal part-used container immediately after use.

#### APPLICATION

#### *Boom Sprayer*

A minimum of 50L water/ha should be used, however, for optimum results, water rates of 70-100L/ha are recommended. Increase the water volume where weed infestation is heavy or the crop cover is dense. Complete coverage of weeds is essential. Higher water volumes (up to 100L/ha) will ensure faster activity of the product on the weeds but may increase the symptoms of crop damage.

The following settings are examples which will ensure excellent coverage of exposed weeds:

Water Rate	50L/ha	75L/ha	75L/ha
Nozzle	Hardi No. 10 or equivalent	Hardi No. 12 or equivalent	Hardi No. 14 or equivalent
Speed	10 km/h	10 km/h	12 km/h
Pressure	240 kPa (2.4 bar)	220 kPa (2.2 bar)	210 kPa (2.1 bar)

#### *Controlled Droplet Application (CDA)*

Insufficient information is available to recommend the application of this product by CDA.

#### Warning

The rubber components present in some spraying units may be affected by exposure to the solvents in Nimble. To reduce this risk it is recommended that the spray unit be thoroughly washed with a boom cleaner and fresh water after use.

#### *Aircraft*

Insufficient information is available to recommend the application of this product by air.

#### COMPATIBILITY

The following herbicide products are physically compatible with Nimble as two-way mixtures in the spray tank, but should only be used for the crops specified, and only when the crop is also specified on the label of the compatible product:

(See below for list of compatible insecticides)

Crop	Nimble	Compatible Product
Wheat, triticale, cereal rye (including undersown)	Up to 750 mL/ha	Ospray Diclofop methyl (barley also), Tristar Advance (barley also), Wildcat 110 EC (wild oats only, high rate)
Wheat, barley, triticale, cereal rye (including undersown)	All rates	Flumetsulam (800 g/kg)
Wheat, barley, triticale, cereal rye (not undersown)	Up to 500 mL/ha	Metsulfuron, Chlorsulfuron, MCPA LVE (500g/L product) (up to 500 mL/ha only)
	All rates	2,4-D amine 500, Eclipse, Cadence (up to 115g only), Clopyralid (300g/L)
Wheat only (not undersown)		Castrate®
Established lucerne only	Up to 750 mL/ha	Simazine (500g/L product) (up to 1.25L/ha only) and simazine (500g/L)/paraquat (200g/L) mixture
Newly sown and established lucerne and clover only	Up to 750 mL/ha	Ospray Quinella®, Fuzilier®, 2,4-DB amine (500g/L)
	Up to 1.0 L/ha	Flumetsulam (800 g/kg)

When mixing Nimble with other herbicides, crop yellowing may be enhanced.

When mixing with Ospray Diclofop methyl, Wildcat110EC or Tristar Advance some reduction in the efficacy and speed of action of these products may occur.

When mixing with Ospray Quinella® or Fuzilier® some reduction in the efficacy and speed of action of these products and Nimble may occur.

In tank-mixtures with Metsulfuron and Chlorsulfuron, rates of Nimble higher than 500 mL/ha may cause significant crop damage.

If the crop is stressed, the application of the herbicide tank-mixtures may cause yield reduction.

When mixing with Cadence a temporary wilting may be evident in some crops after application.

The mixture of Nimble and simazine should be applied during winter to lucerne which is not actively growing. This mixture may result in an increased crop effect but this can be reduced if the lucerne is grazed or cut before spraying.

DO NOT mix Nimble with Ospray Haloxyfop 520 .

Growers should seek advice before spraying recently released cereal varieties.

This product may be mixed in the spray tank with one of the following insecticides according to the directions for the insecticide product: Cyren® ( 500 g/L chlorpyrifos) , deltamethrin (27.5g/L product), Danadim® (400 g/L dimethoate), Ospray Alpha-cypermethrin (100g/L product), Omethoate (290g/L product), Ospray Bifenthrin and endosulfan.

Use the recommended rates for Nimble and its tank-mix partner as well as the surfactant recommendation of the tank-mix partner. Read the label of the tank-mix partner before mixing and using the tank mixture. If another herbicide is applied as a tank mix, observe the plantback restrictions on that label.

#### Warning

DO NOT use crop oils with Nimble or Nimble tank mixtures in cereals.

As formulations of other manufacturers' products are beyond the control of Ospray, all mixtures should be tested prior to mixing commercial quantities.

#### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Wash sprayer thoroughly after use.

#### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

#### STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area.

Do not store for prolonged periods in direct sun light.

Keep from contact with fertilisers, insecticides, fungicides and seeds.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

#### SAFETY DIRECTIONS

Product is harmful if inhaled or swallowed. Will irritate the eyes, nose, throat and skin. Avoid inhaling spray mist. When preparing spray wear elbow length PVC gloves and face shield. If product in eyes wash out immediately with water. After use and before eating drinking or smoking, wash hand arms and face thoroughly with soap and water.

After each day’s use wash gloves, face shield and contaminated clothing.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

#### MATERIAL SAFETY DATA SHEET

For further information, refer to the Material Safety Data Sheet (MSDS) which is available from the supplier.

#### Manufacturers Warranty and Exclusion of Liability

Ospray Pty Ltd has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Ospray Pty Ltd accepts no responsibility for any loss arising from incorrect storage, handling or use.

® - Registered Trademark of Ospray Pty Ltd