



POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm

Bromicide® 200

Selective Herbicide

ACTIVE CONSTITUENT: 200 g/L BROMOXYNIL present as the n-octanoyl ester

SOLVENT: 616 g/L LIQUID HYDROCARBON

GROUP **C** HERBICIDE

For the control of weeds in wheat, oats, barley, cereal rye and triticale (alone or undersown), grain sorghum, linseed, clover and lucerne pastures, turf, fallow, non-crop, roadsides and rights-of-way as per the Directions For Use table.

READ COMPLETE DIRECTIONS FOR USE BEFORE USING THIS PRODUCT.

nufarm.com.au

® Bromicide is a registered trademark of Nufarm Australia Limited

APVMA Approval No.:31614/113089

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply by mister machines.

DO NOT apply if crops or weeds are stressed due to excessively dry or moist conditions.

DO NOT apply to diseased or frost affected crops or turf or if frosts are imminent.

DO NOT apply when rain is expected within 3 hours.

DO NOT apply to lucerne crops or clover based pastures or turf in temperatures above 20°C or when temperatures above 20°C may follow for some days after application.

DO NOT apply when windspeed is less than 3 or more than 20km per hour at application site.

DO NOT apply during surface temperature inversion conditions at application site.

DO NOT apply with smaller than COARSE spray quality according to ASAE S572 definition for standard nozzles.

1. CROP USES

SITUATION AND CROP	WEEDS CONTROLLED	WEED STAGE	STATE	RATE	CRITICAL COMMENTS
Wheat, Barley, Cereal rye, Oats and Triticale: including those undersown with Clover, Lucerne or Medic (Vic only); Linseed: Lotus seed or forage crops (Qld only)	Amsinckia (<i>Amsinckia</i> spp.), Black bindweed (Climbing buckwheat) (<i>Fallopia convolvulus</i>), Capeweed (<i>Arctotheca calendula</i>), Chamomile (<i>Matricaria matricarioides</i>), Common peppergrass (<i>Lepidium africanum</i>), Corn gromwell (Ironweed, Sheepweed) (<i>Buglossoides arvensis</i>), Fat hen (<i>Chenopodium album</i>), Field madder (<i>Sherardia arvensis</i>), Lesser swinecress (<i>Coronopus didymus</i>), Mountain sorrel (<i>Oxalis acetosella</i>), Purple calandrinia (<i>Calandrinia menziesii</i>), Saffron thistle (<i>Carthamus lanatus</i>), Shepherd's purse (<i>Capsella bursa-pastoris</i>), Three-horned bedstraw (<i>Gallium tricornutum</i>), Tree hogweed (<i>Polygonum patulum</i>), Variegated thistle (<i>Silybum marianum</i>)	Up to the 4 leaf stage or when plants are no more than 35mm in diameter	All States	1.4L/ha	CROP STAGE: Wheat, Barley, Cereal, Rye, Oats and Triticale: 3 leaf to full tillered (Z13-30). Linseed: 50 to 150mm high. Undersown Lucerne and Clover: After seedlings have one trifoliolate leaf or before there is 150mm regrowth in established stands. Lotus: Apply post-emergence at least 4 weeks after sowing (crop seedlings with a minimum of 2-6 true leaves). APPLICATION: Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Good coverage of the spray on the weed is essential. Apply only in the Autumn and Winter (mid-April to end of August). For optimum control, apply only when the weeds are actively growing and before the weeds are shielded by the crop.
		Up to the 6 leaf stage or when plants are no more than 50mm in diameter		2L/ha	

(Cont over)

SITUATION AND CROP	WEEDS CONTROLLED	WEED STAGE	STATE	RATE	CRITICAL COMMENTS
Wheat, Barley, Cereal rye, Oats and Triticale: including those undersown with Clover, Lucerne or Medic (Vic only): Linseed: Lotus seed or forage crops (Qld only)	Mexican poppy (<i>Argemone ochroleuca</i>), Three cornered Jack (Doublegee, Spiny emex) (<i>Emex australis</i>), Turnip weed (Giant mustard) (<i>Rapistrum rugosum</i>), Wild radish (<i>Raphanus raphanistrum</i>), Wild mustard (<i>Sisymbrium</i> spp.), Wild turnip (<i>Brassica tournefortii</i>), Wireweed [#] (<i>Polygonum aviculare</i>), Paterson's curse [#] (Salvation Jane) (<i>Echium plantagineum</i>)	Up to the 4 leaf stage or when plants are no more than 50mm in diameter	All States	2L/ha	CAUTION: Undersown Lucerne and Clover: This product will cause slight leaf burning of undersown lucerne and clover. DO NOT apply to Persian clover or Berseem clover. Linseed: DO NOT spray linseed by aerial application. Lotus: Some initial scorching of leaves may occur in some cases. Medics: Application to some cultivars of Medics may result in unacceptable crop damage in some situations. [#] Effective control of Paterson's curse and Fumitories will only be obtained under good growing conditions. ^{##} Where Wireweed is growing under stress, it has been found to be less susceptible.
	Fumitories [#] (<i>Fumaria</i> spp.)		Qld, NSW, Vic, SA, WA, ACT only	2L/ha	
	Pheasant's eye (<i>Adonis microcarpa</i>), Birds eye (Carrot Weed) (<i>Cotula australis</i>)	Up to the 4 true leaf stage Greater than the 4 true leaf stage	SA only	700mL/ha 1.4L/ha	
	Fireweed (<i>Senecio</i> spp.)	Seedlings Early flowering stage	Qld, NSW, ACT only	1.4L/ha 2.8L/ha	
Wheat, Barley, Cereal rye, Oats and Triticale: including those undersown with Clover, Lucerne or Medics: Clover Seed Crops	Amsinckia (<i>Amsinckia</i> spp.), Capeweed (<i>Arctotheca calendula</i>), Chamomile (<i>Matricaria matricarioides</i>), Charlock (<i>Sinapis arvensis</i>), Corn gromwell (Ironweed, Sheepweed) (<i>Buglossoides arvensis</i>), Fat hen (<i>Chenopodium album</i>), Field madder (<i>Sherardia arvensis</i>), Hexham scent (King Island Mellilot) (<i>Melilotus indicus</i>), Horned poppy (<i>Glaucium flavum</i>), Indian hedge mustard (<i>Sisymbrium orientale</i>), Lesser swinecress (<i>Coronopus didymus</i>), Mexican poppy (<i>Argemone ochroleuca</i>), Mintweed (<i>Salvia reflexa</i>), Purple calandrinia (<i>Calandrinia menziesii</i>), Paterson's curse (Salvation Jane) (<i>Echium plantagineum</i>), Rough poppy (<i>Papaver hybridum</i>), Saffron thistle (<i>Carthamus lanatus</i>), Shepherd's purse (<i>Capsella bursa-pastoris</i>), Slender thistle (<i>Carduus tenuiflorus</i>), Turnip weed (Giant mustard) (<i>Rapistrum rugosum</i>), Wild radish (<i>Raphanus raphanistrum</i>)	Up to the 5 leaf stage or when plants are no more than 50mm in diameter	WA only	750mL/ha PLUS 440mL Nufarm Amicide [®] 625/ha OR PLUS 700mL of Nufarm MCPA 500/ha	CROP STAGE: Cereals: 3 leaf to fully tillered. (Z13-30). Undersown and seed Clover crops: Apply after clover seedlings have at least three trifoliate leaves APPLICATION: Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Apply only in the Autumn and Winter (mid-April to end of August). Apply only when weeds are actively growing and before the weeds are shielded by the crop. CAUTION: Some crop damage and/or yield loss may occur in Olympic and Shortim wheat cultivars. DO NOT apply to Persian clover or Berseem clover. A slight burning of clover may occur. DO NOT use 2,4-D or MCPA tank mix if wheat, oats, barley or triticale are undersown with lucerne, clover or medics unless some damage is acceptable. This product may be tank mixed with Hoegrass/Nugrass [®] for control of Annual rye grass and wild oats in wheat and barley only. DO NOT use 2,4-D or MCPA tank mix if using a tank mix with Hoegrass/Nugrass [®]
	Common peppergrass (<i>Lepidium africanum</i>), Tree hogweed (<i>Polygonum patulum</i>)		Vic, SA only	1.4/ha PLUS 440mL Nufarm Amicide [®] 625/ha OR PLUS 700mL of Nufarm MCPA 500/ha	
	Black bindweed (Climbing buckwheat) (<i>Fallopia convolvulus</i>)	Up to the 8 leaf stage.	Vic, SA only		

SITUATION AND CROP	WEEDS CONTROLLED	WEED STAGE	STATE	RATE	CRITICAL COMMENTS
Wheat, Oats, Barley, Triticale alone or with undersown lucerne and clover	Common sowthistle (<i>Sonchus oleraceus</i>)	Up to 4 leaves	Qld, Northern NSW	1.4L/ha	<p>APPLICATION: Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Good coverage of the spray on the weed is essential. Apply only in the Autumn and Winter (mid-April to end of August). For optimum control, apply only when the weeds are actively growing and before the weeds are shielded by the crop.</p> <p>CAUTION:</p> <p>Lucerne and Clover: This product will cause slight leaf burning of undersown lucerne and clover. DO NOT apply to Persian clover or Berseem clover. Strictly for use on small weeds early in crop before shading out.</p> <p>CROP STAGE:</p> <p>Wheat, Barley, Oats and Triticale: 3 leaf to mid tillering (Z13-23). Use a minimum spray volume of 70L/ha. If Group B resistance is suspected, or weeds have grown through a previous application of chlorsulfuron apply 2.1L/ha.</p>
		4-8 leaves		2.1L/ha	
Wheat, Barley, Oats and Triticale	Three cornered Jack (Double gee, Spiny emex) (<i>Emex australis</i>), Wireweed (<i>Polygonum aviculare</i>), Variegated thistle (<i>Silybum marianum</i>)	Up to the 4 leaf stage or when plants are no more than 50mm in diameter	Qld, NSW, Tas, SA, ACT only	1.4/ha PLUS 440mL Nufarm Amicide® 625/ha OR PLUS 700mL of Nufarm MCPA 500/ha	<p>CROP STAGE: 5 leaf to fully tillered (Z15-30): Apply under good growing conditions.</p> <p>BOOM SPRAYING: Use a minimum of 220L spray per hectare on Black bindweed when crop is greater than 300mm high.</p> <p>CAUTION: Some crop damage and/or yield loss may occur in Olympic and Shortim wheat cultivars. DO NOT mix with 2,4-D or MCPA tank mix if wheat, oats, barley or triticale are undersown with lucerne, clover or medics unless some damage is acceptable. DO NOT apply to medics (WA only). This product may be tank mixed with Hoegrass/Nugrass® for control of Annual rye grass and wild oats in Wheat and Barley only. DO NOT use 2,4-D or MCPA tank mix if using a tank mix with Hoegrass®/Nugrass®.</p> <p>APPLICATION: Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application.</p>
	Fumitories (<i>Fumaria</i> spp.)		Qld, NSW, ACT, Vic only		
	Wild mustard (<i>Sisymbrium</i> spp.), Wild turnip (<i>Brassica tournefortii</i>)		Qld, Vic only		
	Cleavers	2-4 stems + 1-3 whorls of leaves/stem	Qld, Vic, NSW, ACT, Tas only		
	Three cornered Jack (Double gee, Spiny emex) (<i>Emex australis</i>), Wireweed (<i>Polygonum aviculare</i>)	4-5 leaf stage	WA only		
	Fumitories (<i>Fumaria</i> spp.)		Qld, Vic, NSW, ACT, SA, Tas only		
	Black bindweed (Climbing buckwheat) (<i>Fallopia convolvulus</i>), Chamomile (<i>Matricaria matricarioides</i>), Fat hen (<i>Chenopodium album</i>), Field madder (<i>Sherardia arvensis</i>), Lesser swinecress (<i>Coronopus didymus</i>), Mountain sorrel (<i>Oxalis acetosella</i>), Purple calandrinia (<i>Calandrinia menziesii</i>), Paterson's curse (Salvation Jane) (<i>Echium plantagineum</i>), Shepherd's purse (<i>Capsella bursa-pastoris</i>), Three-horned bedstraw (<i>Gallium tricornutum</i>), Variegated thistle (<i>Silybum marianum</i>)	5-8 leaf stage	Qld, NSW, Vic, SA, ACT only		

SITUATION AND CROP	WEEDS CONTROLLED	WEED STAGE	STATE	RATE	CRITICAL COMMENTS
Grain sorghum	Cowvine or Peachvine (<i>Ipomoea lonchophylla</i>), Bellvine (<i>I. plebeia</i>)	Up to 4 leaves	Qld, NSW only	1.5L/ha	<p>CROP STAGE: 4-12 leaf stage. This product may cause transient leaf burn of some sorghum varieties.</p> <p>APPLICATION: Beyond the 6-8 leaf stage of the crop, dropped nozzles should be used to minimize chemical being sprayed into the whorl and on upper leaves of the crop and to reduce shading of weeds. Use on small weeds early in the crop life to prevent shading out of the weeds. Spray when weeds are actively growing and not when drought stressed. Use a minimum spray volume of 70L/ha.</p>
	Cowvine or Peachvine (<i>Ipomoea lonchophylla</i>)	4-8 leaves		2L/ha	
	Cowvine or Peachvine (<i>Ipomoea lonchophylla</i>), Bellvine (<i>I. plebeia</i>), Morning glory (<i>I. Purpurea</i>), Bladder Ketmia (<i>Hibiscus trionum</i>), Annual ground cherry (<i>Physalis angulata</i>), Fierce thornapple (<i>Datura ferox</i>), Anoda weed (<i>Anoda cristata</i>)	Up to 4 leaves 4-8 leaves	Qld, NSW only	1.5L/ha PLUS 1.1kg/ha Atradex	<p>CROP STAGE: 4-12 leaf stage. This product may cause transient leaf burn of some sorghum varieties.</p> <p>APPLICATION: Beyond the 6-8 leaf stage of the crop, dropped nozzles should be used to minimize chemical being sprayed into the whorl and on upper leaves of the crop and to reduce shading of weeds. Use on small weeds early in the crop life to prevent shading out of the weeds. Spray when weeds are actively growing and not when drought stressed. Use a minimum spray volume of 70L/ha.</p> <p>Add a non-ionic surfactant. DO NOT use on sandy soils.</p>

2. PASTURES AND OTHER USES

SITUATION AND CROP	WEEDS CONTROLLED	WEED STAGE	STATE	RATE			CRITICAL COMMENTS
				Boom sprayer	High Volume	Knapsak	
Pastures - Grass, Lucerne and Clover based	Pheasant's eye (<i>Adonis microcarpa</i>), Birds eye (Carrot Weed) (<i>Cotula australis</i>)	Up to 4 leaves More than 4 leaves	SA only	750 mL/ha 1.5L/ha	75mL per 100L	15mL/100m ² per 15L	PASTURE GROWTH STAGE: Grass pastures: 3 leaf to full tillered (Z13-30). Lucerne and Clover pastures: After seedlings have one trifoliate leaf or before there is 150mm regrowth in established stands. CAUTION: Lucerne and Clover pastures: This product will cause slight leaf burning of lucerne and clover. DO NOT apply to Persian clover or Berseem clover. Application to some cultivars of Medics may result in unacceptable crop damage in some situations. DO NOT apply to polymorpha medics. Avoid application to clover-based pastures in temperatures over 20°C or when temperatures above 20°C may follow for some days after application; as seedling mortality may occur and established plants may be damaged. APPLICATION: Apply during Autumn/Winter, when weeds are young and actively growing. Not effective on mature plants. For volunteer cotton control, including Roundup Ready varieties a minimum spray volume of 80L/ha should be used for boom sprayer application. Bromicide 200 alone or with glyphosate will not provide satisfactory control of ratoon or "stub" cotton.
	Fireweed (<i>Senecio madagascariensis</i>)	Seedlings	Qld, NSW, ACT only	1.5L/ha	75mL per 100L	15mL/100m ² per 15L	
	Amsinckia (<i>Amsinckia</i> spp.), Bellvine (<i>I. plebeia</i>), Black bindweed (Climbing buckwheat) (<i>Fallopia convolvulus</i>), Capeweed (<i>Arctotheca calendula</i>), Chamomile (<i>Matricaria matricarioides</i>), Common peppergrass (<i>Lepidium africanum</i>), Common sowthistle (<i>Sonchus oleraceus</i>), Corn gromwell (Ironweed, Sheepweed) (<i>Buglossoides arvensis</i>), Cowvine or Peachvine (<i>Ipomoea lonchophylla</i>), Fat hen (<i>Chenopodium album</i>), Field madder (<i>Sherardia arvensis</i>), Lesser swinecress (<i>Coronopus didymus</i>), Mountain sorrel (<i>Oxalis acetosella</i>), Purple calandrinia (<i>Calandrinia menziesii</i>), Saffron thistle (<i>Carthamus lanatus</i>), Shepherd's purse (<i>Capsella bursa-pastoris</i>), Three cornered Jack (Double gee, Spiny emex) (<i>Emex australis</i>), Three-horned bedstraw (<i>Gallium tricornutum</i>), Tree hogweed (<i>Polygonum patulum</i>), Variegated thistle (<i>Silybum marianum</i>), Wild radish (<i>Raphanus raphanistrum</i>)	Up to the 4 leaf stage or when plants are no more than 35mm in diameter	All States				

SITUATION AND CROP	WEEDS CONTROLLED	WEED STAGE	STATE	RATE			CRITICAL COMMENTS	
				Boom sprayer	High Volume	Knapsak		
Fallow, Non Crop, Roadsides and Rights of Way	African daisy (<i>Senecio pterophorus</i>)	Up to and including the flowering stage	SA only	6L/ 500L water per ha	300mL per 100L	60mL/100m ² per 15L	The plants, including the stems, must be thoroughly wet to the point of run-off. Apply when the plants are actively growing. Nufarm Activator® should be added at 125mL per 100L of spray. A follow-up treatment may be necessary for large, well established plants. Ensure an overall spray coverage on weed seedlings.	
	Boneseed/Bitou-bush (<i>Chrysanthemoides monilifera</i>)	Seedlings only	Vic, Tas only		160 mL/100L			
	Cowvine or Peachvine (<i>Ipomoea lonchophylla</i>)	Up to 6 leaves	Qld, NSW, ACT only	1.5L/ha	75mL per 100L	15mL/100m ² per 15L		
		Up to 8 leaves		2.1L/ha	100mL per 100L	20mL /100m ² per 15L		
	Climbing buckwheat (<i>Fallopia convolvulus</i>)	Cotyledon to mature, vining plants	Qld, NSW, WA and NT only	1.5L/ha or 1-1.5L/ha plus glyphosate	75mL per 100L or 75mL per 100L plus glyphosate	15mL /100m ² per 15L or 15mL/100m ² per 15L plus glyphosate		Complete coverage is essential, plants which are shaded by stubble or other weeds will not be controlled. Weeds that are drought stressed and sprayed under hot conditions may not be adequately controlled. The addition of glyphosate will provide broader spectrum weed control where needed. Refer to the Compatibility section for appropriate glyphosate products. The rate of glyphosate used should be robust for the conditions and sufficient to control the hardest-to-kill weed in the situation. For general fallow weed control application volumes above 50L/ha are recommended.
Volunteer conventional cotton (<i>Gossypium hirsutum</i>), Volunteer Roundup Ready® cotton (<i>Gossypium hirsutum</i>)	Cotyledon to 6 leaves	Qld, NSW, WA and NT only						
Turf: Established Couch (including Queensland blue), Buffalo, Bent, Paspalum, Kikuyu, Fescue, and Rye lawns	Bindy-eye (<i>Calotis hispidula</i>), Jo-jo (Onehunga) (<i>Soliva pterosperma</i>), Cudweed (<i>Gnaphalium</i> spp.)	Seedlings only	All states	6L/ 500L water per ha	300mL per 100L	60mL/100m ² per 15L	Apply evenly over the area to be treated. A slight browning of grasses, especially Paspalum, may occur in hot conditions.	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

HARVEST

ALL CROPS: NOT REQUIRED WHEN USED AS DIRECTED.

GRAZING/STOCKFOOD

PASTURES: DO NOT GRAZE TREATED PASTURES OR CUT FOR STOCKFOOD FOR 14 DAYS AFTER APPLICATION

ALL OTHER CROPS AND SITUATIONS, EXCEPT PASTURES: DO NOT GRAZE TREATED CROPS/AREAS OR CUT FOR STOCKFOOD FOR 8 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Before opening, carefully read Directions for Use, Precautionary and Protection Statements, Storage and Disposal, Safety Directions and First Aid Instructions. Use only at recommended growth stages of crops and weeds. DO NOT use on crops other than recommended.

MIXING INFORMATION

To ensure even mixing, half fill the spray tank with clean water and add the required amount of this product. Agitate thoroughly, then add the remainder of water. Agitate again before spraying commences.

BOOM SPRAYER APPLICATION

USE ONLY low boom equipment set up to provide good coverage of weeds within the crop canopy. USE ONLY coarse spray quality according to ASAE S572 definition for standard nozzles. Apply 50-200L (WA: 50-100L) of water per hectare. Calibrate the machine before use to ensure that the correct application rate is obtained. Check height of boom above the crop to ensure that the spray is evenly distributed. A minimum of 160L and preferably 200L of water per hectare is recommended where the weed infestation is heavy or the crop cover is dense.

HIGH VOLUME (HAND GUN) AND KNAPSACK APPLICATION

The plants, including the stems, must be thoroughly wet to the point of run-off. Apply when the plants are actively growing. A follow-up treatment may be necessary. Ensure an overall spray coverage on weed seedlings.

AERIAL APPLICATION

Apply a minimum of 22L of water per hectare. Markers should always move upwind before aircraft passes overhead. Application by microneaire equipment may give variable results.

SPRAY APPLICATIONS AND DRIFT RISK ASSESSMENT



DO NOT apply when wind speed is less than 3 or more than 20 kilometres per hour (ground application) as measured at the application site.

DO NOT apply when wind speed is less than 3 or more than 15 kilometres per hour (aerial application) as measured at the application site.

For aerial application it is recommended where possible for this product to be applied by an aerial applicator business that holds current accreditation for the Aerial Improvement Management System – "AIMS", issued by the Aerial Agricultural Association of Australia Ltd.

Checklist:

- Have you cleaned/decontaminated your boom sprayer?
- Have you contacted your neighbour prior to spraying?
- Is your sprayer set-up correctly for the particular application?
- Check
 - boom calibration
 - at nozzle - nozzle choice
 - low drift/what spray quality
 - coarse or larger spray quality?
 - boom height - speed of intended application
 - water volume
- You must check, determine and record the weather conditions immediately prior to, and immediately after the spray application is made.
- Record
 - Temperatures
 - Relative Humidity
 - Delta T
 - Wind speed
 - Is there a temperature inversion?
- Night Spraying - Extra care is required to ensure that inversion conditions are not present. Use smoke generator to determine wind direction and presence of inversion conditions.

For further information refer to nufarm.com.au/spraywise



spraywisedecisions.com.au is an online weather forecasting program and is recommended for use when planning your pesticide application

EQUIPMENT MAINTENANCE

After use, wash equipment thoroughly and rinse several times prior to re-use. Wash sprayer thoroughly with clean water after use. If using tank mixes with Amicide® 625 or MCPA 500 apply with equipment that will not be used in 2,4-D and MCPA susceptible crops, particularly cotton and vines.

COMPATIBILITY

This product may be mixed in the spray tank with one of the following herbicides or insecticides in wheat and barley only. Nufarm Nugrass®, Omethoate, Chlorpyrifos, Dimethoate, Nufarm Amicide® 625 Herbicide (2,4-D Amine), Nufarm MCPA 500 Herbicide, Agritone® 750, Glean/Lusta® Ally/Associate®, Brodal®, Glyphosate products (Nufarm Glyphosate CT, Weedmaster® DUO, Weedmaster® DST, Weedmaster® ARGO®, Nufarm Kamba® 500, Paragon®. Bromicide® 200 is compatible with Comet® 200/Starane® 200 when needed for control of mixed weed populations in grain sorghum. As formulations of other manufacturers products are beyond the control of Nufarm, all mixtures should be tested prior to mixing commercial quantities.

RESISTANT WEEDS WARNING

GROUP	C	HERBICIDE
-------	----------	-----------

Bromicide 200 Selective Herbicide (Bromicide 200) is a member of the nitriles group of herbicides. Bromicide 200 has the inhibitors of photosynthesis at photosystem II mode of action. For weed resistance management Bromicide 200 is a Group C herbicide. Some naturally-occurring weed biotypes resistant to Bromicide 200 and other Group C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group C herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from the failure of Bromicide 200 to control resistant weeds.

PRECAUTION

RE-ENTRY PERIOD: DO NOT allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Drift Warning: DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

Note: Maximum daily temperatures at or above 20°C at application, or on days following treatment may result in seedling mortality and established plants may be damaged.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Non Returnable containers

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

Refillable containers (1000L only)

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Envirodrum® 110L Mini Bulk Returnable Container

Store the original sealed Envirodrum in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any other foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Nufarm Australia Limited.

SAFETY DIRECTIONS

Product is poisonous if inhaled or swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray wear elbow-length PVC gloves and face-shield. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone 13 11 26). If swallowed, DO NOT induce vomiting. Give a glass of water.

ADDITIONAL STATEMENTS (WHS REGULATIONS 2011)

Suspected of damaging fertility or the unborn child. Suspected of causing cancer. Causes serious eye irritation. Causes skin irritation. May cause an allergic skin reaction. May be fatal if swallowed and enters airways. May cause drowsiness or dizziness. Repeated exposure may cause skin dryness or cracking. Do not eat, drink or smoke when using this product. Use only outdoors or in a well-ventilated area. Wear protective gloves, clothing, eye and face protection. Contaminated work clothing should not be allowed out of the workplace. IF SWALLOWED: Rinse mouth. IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention. IF ON SKIN: Wash with plenty of soap and water. If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash before reuse.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained from your supplier or from the Nufarm website – www.nufarm.com.au

In case of emergency: Phone 1800 033 498. Ask for shift supervisor. Toll free 24 hours.

CONDITIONS OF SALE

"Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded.

Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute."

® Bromicide, Nugrass, Amicide, Agritone, Associate, Weedmaster, ARGO, Comet, Lusta and Kamba are registered trademarks of, or used under licence by Nufarm Australia Limited.

* Other registered trademarks

**POISON**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm

Bromicide[®] 200

Selective Herbicide

ACTIVE CONSTITUENTS: 200 g/L BROMOXYNIL present as the n-octanoyl ester
SOLVENT: 616 g/L LIQUID HYDROCARBONGROUP **C** HERBICIDE

For the control of weeds in wheat, oats, barley, cereal rye and triticale (alone or undersown), grain sorghum, linseed, clover and lucerne pastures, turf, fallow, non-crop, roadsides and rights-of-way as per the Directions For Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

® Bromicide is a registered trademark of Nufarm Australia Limited

nufarm.com.au

Nufarm Australia Limited
ACN 004 377 780
103-105 Pipe Road
Laverton North Victoria 3026
Tel: (03) 9282 1000
Fax: (03) 9282 1001

Contents: 1L -1000L

APVMA Approval No.: 31614/113089

STORAGE AND DISPOSAL**Non Returnable containers**

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

Refillable containers (1000L only)

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Envirodrum[®] 110L Mini Bulk Returnable Container

Store the original sealed Envirodrum in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any other foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Nufarm Australia Limited.

SAFETY DIRECTIONS

Product is poisonous if inhaled or swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray wear elbow-length PVC gloves and face-shield. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed DO NOT induce vomiting. Give a glass of water.

ADDITIONAL STATEMENTS (WHS REGULATIONS 2011)

Suspected of damaging fertility or the unborn child. Suspected of causing cancer. Causes serious eye irritation. Causes skin irritation. May cause an allergic skin reaction. May be fatal if swallowed and enters airways. May cause drowsiness or dizziness. Repeated exposure may cause skin dryness or cracking. Do not eat, drink or smoke when using this product. Use only outdoors or in a well-ventilated area. Wear protective gloves, clothing, eye and face protection. Contaminated work clothing should not be allowed out of the workplace. IF SWALLOWED: Rinse mouth. IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention. IF ON SKIN: Wash with plenty of soap and water. If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash before reuse.

SAFETY DATA SHEET


For further information refer to the Safety Data Sheet (SDS), which can be obtained from your supplier or from the Nufarm website – www.nufarm.com.au

In case of emergency: Phone 1800 033 498. Ask for shift supervisor. Toll free 24 hours.

CONDITIONS OF SALE

"Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute."

BN / DOM / Barcode / drumMUSTER logo / Envirodrum /



ENVIRONMENTALLY HAZARDOUS SUBSTANCES, LIQUID, N.O.S. (CONTAINS BROMOXYNIL)
UN No: 3082
PACKING GROUP: III
HAZCHEM: 2 Z
IN AN EMERGENCY DIAL: 000 POLICE OR FIRE BRIGADE
IN AN EMERGENCY, SPECIALIST ADVICE NUFARM AUSTRALIA LIMITED 1800 033 498

"NOTE: MANUFACTURED IN AUSTRALIA – DG CLASS 9 MARKING ONLY REQUIRED FOR SHIPMENT TO TASMANIA – THIS IS DONE BY NUFARM LOGISTICS"