

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

## Genfarm Bromoxynil 200 Selective Herbicide

**ACTIVE CONSTITUENT: 200 g/L BROMOXYNIL**  
Present as the n-octanoyl ester  
**SOLVENT: 662 g/L LIQUID HYDROCARBON**

GROUP	<b>C</b>	HERBICIDE
-------	----------	-----------

For the control of weeds in wheat, oats, barley, cereal rye and triticale (alone or undersown), linseed, clover and Lucerne pastures, turf, fallow, non-crop, roadsides and rights-of-way as per the Directions For Use table.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE.**

Contents: 20 Litres, 110 Litres, 120 litres

**Genfarm Crop Protection Pty Ltd**  
Suite 2, Level 3, 64 Talavera Road,  
Macquarie Park, NSW, 2113  
Tel: (02) 9889 5400

## GENERAL INSTRUCTIONS

Before opening, carefully read Directions for Use, Precautionary and Protection Statements, Storage and Disposal, Safety Directions and First Aid Instructions. Use only at recommended growth stages of crops and weeds. Do not use on crops other than recommended.

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## SAFETY DIRECTIONS

Product is poisonous if inhaled or swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray wear elbow-length PVC gloves and face-shield. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed do NOT induce vomiting. Give a glass of water.

## MATERIAL SAFETY DATA SHEET

For further information, refer to the Material Safety Data Sheet (MSDS) which is available from the supplier.

## CONDITIONS OF SALE

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Genfarm Crop Protection Pty Ltd. accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd. will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

APVMA Approval No.: 62204/20L/1107, 62204/110L/1107, 62204/120L/1107

Batch No:

Date of Manufacture:

## Bar codes

20L 933 537 100 224 9

110L 933 537 100 225 6

120L 933 537 100 226 3



In a Transport Emergency Dial <b>000</b> Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY <b>1800 033 111</b> ALL HOURS AUSTRALIA WIDE
--	---

# **POISON**

**KEEP OUT OF REACH OF CHILDREN**

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

## **Genfarm Bromoxynil 200 Selective Herbicide**

**ACTIVE CONSTITUENT: 200 g/L BROMOXYNIL**

**Present as the n-octanoyl ester**

**SOLVENT: 662 g/L LIQUID HYDROCARBON**

For the control of weeds in wheat, oats, barley, cereal rye and triticale (alone or undersown), linseed, clover and Lucerne pastures, turf, fallow, non-crop, roadsides and rights-of-way as per the Directions For Use table.

**IMPORTANT: READ THIS LEAFLET BEFORE USE**

**APVMA APPROVAL NO: 62204/1107**

**Genfarm Crop Protection Pty Ltd**

**Suite 2, Level 3, 64 Talavera Road,**

**Macquarie Park, NSW, 2113**

**Tel: (02) 9889 5400**

## DIRECTIONS FOR USE

### RESTRAINTS

DO NOT apply by mister machines.

DO NOT apply if crops or weeds are stressed due to excessively dry or moist conditions.

DO NOT apply to diseased or frost affected crops or turf or if frosts are imminent.

DO NOT apply when rain is expected within 3 hours.

DO NOT apply to Lucerne crops or clover based pastures or turf in temperatures above 20°C or when temperatures above 20°C may follow for some days after application.

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate	Critical Comments
<b>Wheat, Barley, Cereal Rye, Oats and Triticale:</b> Including those undersown with <b>Clover, Lucerne or Medic</b> (Vic only); <b>Linseed; Clover or Lucerne Pastures; Lotus seed or forage crops</b> (Qld only)	Amsinckia ( <i>Amsinckia</i> spp.) Black Bindweed (Climbing buckwheat) ( <i>Fallopia convolvulus</i> ) Capeweed ( <i>Arctotheca calendula</i> ) Chamomile ( <i>Matricaria matricarioides</i> ) Common Peppergrass ( <i>Lepidium africanum</i> ) Corn Gromwell (Ironweed, Sheepweed) ( <i>Buglossoides arvensis</i> ) Fat Hen ( <i>Chenopodium album</i> ) Field Madder ( <i>Sherardia arvensis</i> ) Lesser Swinecress ( <i>Coronopus didymus</i> ) Mountain sorrel ( <i>Oxalis acetosella</i> ) Purpose Calandrinia ( <i>Calandrinia menziesii</i> ) Saffron Thistle ( <i>Carthamus lanatus</i> ) Shepherd's Purse ( <i>Capsella bursa-pastoris</i> ) Three-horned bedstraw ( <i>Gallium tricornutum</i> ) Tree Hogweed ( <i>Polygonum patulum</i> ) Variegated Thistle ( <i>Silybum marianum</i> )	Up to the 4 leaf stage or when plants are no more than 35mm in diameter	All States	1.4L/ha	<b>CROP STAGE:</b> <b>Wheat, Barley, Cereal, Rye, Oats and Triticale:</b> 3 leaf to full tillered (Zadoks Scale Z13-Z30). <b>Linseed:</b> 50 to 150mm high. <b>Lucerne and Clover:</b> After seedlings have one trifoliate leaf or before there is 150mm regrowth in established stands. <b>Lotus:</b> Apply post-emergence at least 4 weeks after sowing (crop seedlings with a minimum of 2-6 true leaves). <b>APPLICATION:</b> Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Good coverage of the spray on the weed is essential. Apply only in the Autumn and Winter (mid-April to end of August). For optimum control, apply only when the weeds are actively growing and before the weeds are shielded by the crop. <b>CAUTION:</b> <b>Lucerne and Clover:</b> This product will cause slight leaf burning of undersown Lucerne and Clover. DO NOT apply to Persian clover or Berseem clover. <b>Linseed:</b> DO NOT spray linseed by aerial application. <b>Lotus:</b> Some initial scorching of leaves may occur in some cases. <b>Medics:</b> Application to some cultivars of Medics may result in unacceptable crop damage in some situations. #Effective control of Paterson's Curse and Fumitories will only be obtained under good growing conditions. ##Where Wireweed is growing under stress, it has been found to be less susceptible.
		Up to the 6 leaf stage or when plants are no more than 50mm in diameter		2.0L/ha	
	Mexican Poppy ( <i>Argemone ochroleuca</i> ) Threecornered Jack (Doublegee, Spiny Emex) ( <i>Emex australis</i> ) Turnip Weed (Giant Mustard) ( <i>Rapistrum rugosum</i> ) Wild Radish ( <i>Raphanus raphanistrum</i> ) Wild Mustard ( <i>Sisymbrium</i> spp.) Wild Turnip ( <i>Brassica tournefortii</i> ) Wireweed <sup>##</sup> ( <i>Polygonum aviculare</i> ) Paterson's Curse <sup>#</sup> ( <i>Echium plantagineum</i> ) (Salvation Jane)	Up to the 4 leaf stage or when plants are no more than 50mm in diameter	Qld, NSW, Vic, SA, WA, ACT only		
	Pheasant's Eye ( <i>Adonis microcarpa</i> ) Birds Eye (Carrot Weed) ( <i>Cotula australis</i> )	Up to the 4 true leaf stage	SA only	700mL/ha	DO NOT use where undersown medics occur.
		Greater than the 4 true leaf stage		1.4L/ha	

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate	Critical Comments
<b>Wheat, Barley, Cereal Rye, Oats and Triticale:</b> including those undersown with <b>Clover, Lucerne or Medic</b> (Vic only); <b>Linseed; Clover or Lucerne Pastures; Lotus seed or forage crops</b> (Qld only) (cont'd)	Fireweed ( <i>Senecio</i> spp.)	Seedlings	Qld, NSW, ACT only	1.4L/ha	Apply during the Autumn-Winter period when weeds are young and actively growing. Not effective on mature plants.
		Early flowering stage		2.8L/ha	
<b>Wheat, Barley, Cereal Rye, Oats and Triticale:</b> including those undersown with Clover, Lucerne or Medics; <b>Clover Seed Crops</b>	Amsinckia ( <i>Amsinckia</i> spp.) Capeweed ( <i>Arctotheca calendula</i> ) Chamomile ( <i>Matricaria matricarioides</i> ) Charlock ( <i>Sinapis arvenses</i> ) Corn Gromwell (Ironweed, Sheepweed) ( <i>Buglossoides arvensis</i> ) Fat Hen ( <i>Chenopodium album</i> ) Field Madder ( <i>Sherardia arvensis</i> ) Hexham Scent (King Island Melilot) ( <i>Melilotus indicus</i> ) Horned Poppy ( <i>Glaucium flavum</i> ) Indian Hedge Mustard ( <i>Sisymbrium orientale</i> ) Lesser Swinecress ( <i>Coronopus didymus</i> ) Mexican Poppy ( <i>Argemone ochroleuca</i> ) Mintweed ( <i>Salvia reflexa</i> ) Purple Calandrinia ( <i>Calandrinia menziesii</i> ) Paterson's Curse (Salvation Jane) ( <i>Echium plantagineum</i> ) Rough Poppy ( <i>Papaver hybridum</i> ) Saffron Thistle ( <i>Carthamus lanatus</i> ) Shepherd's Purse ( <i>Capsella bursa-pastoris</i> ) Slender Thistle ( <i>Carduus tenuiflorus</i> ) Turnip Weed (Giant Mustard) ( <i>Rapistrum rugosum</i> ) Wild Radish ( <i>Raphanus raphanistrum</i> )	Up to the 5 leaf stage or when plants are no more than 50mm in diameter	WA only	750mL/ha PLUS 440mL Amine 625/ha OR PLUS 700mL of MCPA 500/ha	<b>CROP STAGE:</b> <b>Cereals:</b> 3 leaf to fully tillered. (Zadoks Scale Z13-30). <b>Clover:</b> Apply after clover seedlings have at least three trifoliolate leaves <b>APPLICATION:</b> Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Apply only in the Autumn and Winter (mid-April to end of August). Apply only when weeds are actively growing and before the weeds are shielded by the crop. <b>CAUTION:</b> Some crop damage and/or yield loss may occur in Olympic and Shortim wheat cultivars. DO NOT apply to Persian clover of Berseem clover. A slight burning of clover may occur. DO NOT use 2,4-D or MCPA tank mix if wheat, oats, barley or triticale are undersown with Lucerne, clover or medics unless some damage is acceptable. This product may be tank mixed with Diclofop methyl for control of Annual Rye Grass and Wild Oats in Wheat and Barley only. DO NOT use 2,4-D or MCPA tank mix if using a tank mix with Diclofop methyl.
			Vic, SA only	1.4/ha PLUS 440mL Amine 625/ha OR PLUS 700mL of MCPA 500/ha	
			Vic only		
			Vic, SA only		
	Common Peppergrass ( <i>Lepidium africanum</i> ) Tree Hogweed ( <i>Polygonum patulum</i> )				
	Black Bindweed (Climbing Buckwheat) ( <i>Fallopia convolvulus</i> )	Up to the 8 leaf stage.	Vic, SA only		
<b>Wheat, Oats, Barley, Triticale alone or with undersown Lucerne and clover</b>	Common Sowthistle ( <i>Sonchus oleraceus</i> )				<b>APPLICATION:</b> Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Good coverage of the spray on the weed is essential. Apply only in the Autumn and Winter (mid-April to end of August). For optimum control, apply only when the weeds are actively growing and before the weeds are shielded by the crop.

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate/ha	Critical Comments
Wheat, Oats, Barley, Triticale alone or with undersown Lucerne and clover (cont'd)	Common Sowthistle ( <i>Sonchus oleraceus</i> ) (cont'd)				<b>CAUTION:</b> <b>Lucerne and Clover:</b> This product will cause slight leaf burning of undersown Lucerne and Clover. DO NOT apply to Persian clover or Berseem clover.
		Up to 4 leaves	Qld, Northern NSW	1.4 L/ha	Strictly for use on small weeds early in crop before shading out. <b>CROP STAGE:</b> <b>Wheat, Barley, Oats and Triticale:</b> 3 leaf to mid tillering (Z13-Z23). Use a minimum spray volume of 70 L/ha. If Group B resistance is suspected, or weeds have grown through a previous application of chlorsulfuron apply 2.1 L/ha
		4 to 8 leaves		2.1 L/ha	<b>CROP STAGE:</b> <b>Wheat, Barley, Oats and Triticale:</b> 3 leaf to full tillering (Z13-Z30). This product will cause slight leaf burning of undersown Lucerne and clover.
Wheat, Barley, Oats and Triticale	Threecornered Jack (Doublegee, Spiny Emex) ( <i>Emex australis</i> ) Wireweed ( <i>Polygonum aviculare</i> ) Variegated Thistle ( <i>Silybum marianum</i> ) Fumitories ( <i>Fumaria</i> spp.)	Up to the 4 leaf stage or when plants are no more than 50mm in diameter	Qld, NSW, Tas, SA, ACT only	1.4L/ha PLUS 440mL 2,4-D Amine 625 /ha OR PLUS 700mL of MCPA 500/ha	<b>CROP STAGE:</b> 5 leaf to fully tillered (Zadoks scale Z15030): Apply under good growing conditions. <b>BOOM SPRAYING:</b> Use a minimum of 200L spray per hectare on Black Bindweed when crop is greater than 300mm high. <b>CAUTION:</b> Some crop damage and/or yield loss may occur in Olympic and Shortim wheat cultivars. DO NOT mix with 2,4-D or MCPA tank mix if wheat, oats, barley or triticale are undersown with Lucerne, clover or medics unless some damage is acceptable. DO NOT apply to medics (WA only). This product may be tank mixed with Diclofop methyl for control of Annual Rye Grass and Wild Oats in Wheat and Barley only. DO NOT use 2,4-D or MCPA tank mix if using tank mix with Diclofop methyl. <b>APPLICATION:</b> Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application.
	Qld, NSW, ACT, Vic only				
	Qld, Vic only				
	Wild Mustard ( <i>Sisymbrium</i> spp.) Wild Turnip ( <i>Brassica tournefortii</i> )				
	Cleavers	2-4 stems + 1-3whorls of leaves/stem	Qld, Vic, NSW, ACT, Tas only		
	Threecornered Jack (Doublegee, Spiny Emex) ( <i>Emex australis</i> ) Wireweed ( <i>Polygonum aviculare</i> ) Fumitories ( <i>Fumaria</i> spp.)	4-5 leaf stage	WA only	2.1L/ha PLUS 400mL 2,4-D Amine 625 /ha OR PLUS 700mL of MCPA 500/ha	
	Black Bindweed (Climbing Buckwheat) ( <i>Fallopia convolvulus</i> ) Chamomile ( <i>Matricaria matricarioides</i> ) Fat Hen ( <i>Chenopodium album</i> ) Field Madder ( <i>Sherardia arvensis</i> ) Lesser Swinecress ( <i>Coronopus didymus</i> ) Mountain sorrel ( <i>Oxalis acetosella</i> ) Purple Calandrinia ( <i>Calandrinia menziesii</i> ) Paterson's Curse (Salvation Jane) ( <i>Echium plantagineum</i> ) Shepherd's Purse ( <i>Capsella bursa-pastoris</i> ) Three-horned bedstraw ( <i>Gallium tricornutum</i> ) Variegated Thistle ( <i>Silybum marianum</i> )	5 to 8 leaf stage	All States		

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate	Critical Comments
<b>Fallow &amp; Non Crop</b>	Cowvine or Peachvine ( <i>Ipomoea lonchophylla</i> )	Up to 6 leaves	Qld, NSW, ACT only	1.4 L/ha	For all fallow and non crop situations Complete coverage is essential, plants which are shaded by stubble or other weeds will not be controlled. Weeds that are drought stressed and sprayed under hot conditions may not be adequately controlled. The addition of glyphosate will provide broader spectrum weed control where needed. Refer to the Compatibility section for appropriate glyphosate products. The rate of glyphosate used should be robust for the conditions and sufficient to control the hardest-to-kill weed in the situation. For general fallow weed control application volumes above 50 L/ha are recommended.
		Up to 8 leaves		2.1 L/ha	
	Climbing buckwheat ( <i>Fallopia convolvulus</i> )	Cotyledon to mature, vining plants		1.5L/ha alone or 1-1.5L/ha plus glyphosate	
	Volunteer conventional cotton ( <i>Gossypium hirsutum</i> )	Cotyledon to 6 leaves		1.5L/ha alone or 1-1.5L/ha plus glyphosate	
	Volunteer Roundup Ready cotton ( <i>Gossypium hirsutum</i> )	Cotyledon to 6 leaves		1.5L/ha alone or plus glyphosate	
<b>Pastures</b>	Adonis (Pheasants Eye) Birds Eye (Carrot Weed)	Up to 4 leaves	SA only	700 mL/ha	Apply in May-June. Medics may be severely scorched at the low rate and killed at the high rate. Do not apply to polymorpha medics. Avoid application to clover-based pastures in temperatures over 20°C or when temperatures above 20°C may follow for some days after application; as seedling mortality may occur and established plants may be damaged.
		More than 4 leaves		1.4 L/ha	
	Double Gee, Wild Radish	Up to 4 leaves	WA only	1.5 L/ha	
	Matricaria	Up to 3 leaves		750 mL/ha	
	Fireweed	Seedlings	Qld, NSW, ACT only	1.4 L/ha	
Early flowering		2.8 L/ha			
<b>Pastures, Roadsides and Rights-of-Way</b>	African Daisy ( <i>Senecio pterophorus</i> )	Up to and including the flowering stage	SA only	300mL per 100L water	The plants, including the stems, must be thoroughly wet to the point of run-off. Apply when the plants are actively growing. Spraymate Activator should be added at 125mL per 100 litres of spray. A follow-up treatment may be necessary for large, well-established plants. Ensure an overall spray coverage on weed seedlings.
	Boneseed/Bitou-bush ( <i>Chrysanthemoides monilifera</i> )	Seedlings only	Vic, Tas only	160mL per 100L water	

Situation & Crop	Weeds Controlled	State	Rate			Critical Comments
<b>Turf:</b> Established Couch (including Queensland Blue), Buffalo, Bent, Paspalum, Kikuyu, Fescue and Rye lawns	Bindy-eye ( <i>Calotis hispidula</i> ) Jo-jo (Onehunga) ( <i>Soliva pterosperma</i> ) Cudweed ( <i>Gnaphalium</i> spp.)	All States	6.0L/500L water per ha	<b>High Volume:</b> 300mL per 100L	60mL/100m <sup>2</sup> per 15L knapsack	Apply evenly over the area to be treated. A slight browning of grasses, especially Paspalum, may occur in hot conditions.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD: DO NOT GRAZE TREATED CROPS OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.**

**CROP HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.**

## GENERAL INSTRUCTIONS:

Before opening, carefully read Directions for Use, Precautionary and Protection Statements, Storage and Disposal, Safety Directions and First Aid instructions. Use only at recommended growth stages of crops and weeds. Do not use on crops other than recommended.

## MIXING INFORMATION

To ensure even mixing, half fill the spray tank with clean water and add the required amount of this product. Agitate thoroughly, then add the remainder of water. Agitate again before spraying commences.

## BOOM SPRAYER

Apply 50-200 litres (WA: 50 to 100 litres) of water per hectare. Calibrate the machine before use to ensure that the correct application rate is obtained. Check height of boom above the crop to ensure that the spray is evenly distributed. A minimum of 160 litres and preferably 200 litres of water per hectare is recommended where the weed infestation is heavy or the crop cover is dense.

DO NOT use rain-drop or air-induction nozzles as coverage of weeds may not be adequate.

## AERIAL APPLICATION

Apply a minimum of 22 litres of water per hectare. Markers should always move upwind before aircraft passes overhead. Application by microneaire equipment may give variable results.

## EQUIPMENT MAINTENANCE

After use, wash equipment thoroughly and rise several times prior to re-use. Wash sprayer thoroughly with clean water after use. If using tank mixes with Amicide 625 or MCPA 500 apply with equipment that will not be used in 2,4-D and MCPA susceptible crops, particularly cotton and vines.

## COMPATIBILITY

This product may be mixed in the spray tank with one of the following herbicides or insecticides in wheat and barley only.

Genfarm Diclofop methyl, Omethoate, Genfarm Chlorpyrifos 500, dimethoate, endosulfan, 2,4-D Amine 625, MCPA 500 Herbicide, Genfarm Huron Herbicide, Genfarm Muron 600 Herbicide, Genfarm Destroyer 500 Herbicide (diflufenican), Glyphosate, Dicamba 500.

As formulations of other manufacturers products are beyond the control of Genfarm Crop Protection Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

## RESISTANT WEEDS WARNING

GROUP	C	HERBICIDE
-------	---	-----------

Genfarm Bromoxynil 200 Selective Herbicide is a member of the nitriles group of herbicides. Genfarm Bromoxynil 200 Selective Herbicide has the inhibitors of photosynthesis at photosystem II mode of action. For weed resistance management Genfarm Bromoxynil 200 Selective Herbicide is a Group C herbicide.

Some naturally-occurring weed biotypes resistant to Genfarm Bromoxynil 200 Selective Herbicide and other Group C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group C herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Genfarm Crop Protection Pty. Ltd. Accepts no liability for any losses that may result from the failure of Genfarm Bromoxynil 200 Selective Herbicide to control resistant weeds.

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

**Drift Warning:** DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

**Note:** Maximum daily temperatures at or above 20°C at application, or on days following treatment may result in seedling mortality and established plants may be damaged.

## PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.



## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## SAFETY DIRECTIONS

Product is poisonous if inhaled or swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray wear elbow-length PVC gloves and face-shield. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water.

## MATERIAL SAFETY DATA SHEET

For further information, refer to the Material Safety Data Sheet (MSDS) which is available from the supplier.

## CONDITIONS OF SALE

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Genfarm Crop Protection Pty Ltd. accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd. will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

In a Transport Emergency Dial <b>000</b> Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY <b>1800 033 111</b> ALL HOURS AUSTRALIA WIDE
--	---