

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

United Farmers

CLOPYRALID

Herbicide

ACTIVE CONSTITUENT: 300g/L Clopyralid

GROUP	I	HERBICIDE
-------	---	-----------

For the control of a wide range of broadleaf weeds in wheat, barley, tritacle, oats, pastures, canola, fallow land, forests and industrial situations as specified in the Directions for Use table.

Note: This information has been adapted from the UFCC Clopyralid Herbicide label. Please consult the label on the pack before use.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING THIS PRODUCT.



UNITED FARMERS
CO-OPERATIVE

ABN 14 014 213 492
2 BIRKSGATE ROAD ROUS HEAD, PMB 10
NORTH FREMANTLE, W.A. 6159
Phone 08 9430 2222 Fax 08 9430 2233

STORAGE AND DISPOSAL:

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT store near feedstuffs, fertilisers or seed.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SMALL SPILL MANAGEMENT

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal. (See STORAGE AND DISPOSAL section). If necessary, wash the spill area with an alkali detergent and water and absorb as above, the wash liquid for disposal.

SAFETY DIRECTIONS:

May irritate the eyes and skin. Avoid contact with the eyes and skin. DO NOT inhale spray mist. When preparing the spray, wear elbow-length PVC gloves and face shield. After each day's use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre (Telephone Australia 13 11 26).

MATERIAL SAFETY DATA SHEET:

Additional information is listed in the Material Safety Data Sheet.

CONDITIONS OF SALE

The use of United Farmers Clopyralid Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by United Farmers Cooperative Limited regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and United Farmers Cooperative Limited accepts no responsibility for any consequence whatsoever resulting from the use of this product.

UNITED FARMERS COOPERATIVE LIMITED.
ABN 14 014 213 492
2 BIRKSGATE ROAD, ROUS HEAD, PMB 10
NORTH FREMANTLE WA 6159
Phone (08) 9430 2222 Fax (08) 9430 2233

DIRECTIONS FOR USE

IT IS ESSENTIAL to select a rate appropriate to weed size. Best results will be obtained when weeds are actively growing at treatment.

Restraints:

DO NOT apply to weeds which may be stressed (inactive growth) due to prolonged periods of extreme heat or cold, moisture stress (water logging or drought) or previous herbicide treatment as reduced levels of control may result.

DO NOT sow susceptible crops (see PROTECTION OF CROPS) for nine months following any application up to 300 mL/ha or for twelve months following an application in excess of 300 mL/ha and up to 500 mL/ha. Field peas should not be sown the season following an application of 500 mL/ha. Where rates in excess of 500 mL/ha have been used, susceptible crops including field peas, should not be sown for at least two years.

DO NOT spray if rain is likely within 3 hours.

CROP	CROP STAGE	WEED	WEED STAGE	RATE mL/ha	STATE	CRITICAL COMMENTS			
Barley Oats Triticale Wheat	Pre-sowing	Capeweed	Up to 8 leaf and maximum 10 cm diameter	150mL plus knockdown herbicide	WA only	Pre-sowing: This rate should only be used in tank mixture with formulations of paraquat/diquat or glyphosate.			
				150mL plus diuron at 300 mL/ha		Post sowing pre-emergent to 3 leaf: This rate should only be used in tank mixture with diuron for control of transplants.			
			Cotyledons to 6 leaf and maximum 5 cm diameter	150mL		Early post-emergent: Weeds should be growing actively and not larger than 5 cm diameter.			
	Post-sowing pre-emergence through to 3 leaf	Early post-emergence (2 leaf to jointing)	Capeweed Soldier thistle	Up to 10 cm diameter (4 to 8 leaf)	300mL	NSW, Vic, SA, Tas and WA only	Weeds should be young and actively growing. Weeds will become stunted and not be competitive soon after application although final results may not show for some weeks. Faba beans and lupins will only be suppressed. Use 75 mL/ha rate only in combination with MCPA LVE.		
					250mL				
					150mL				
					75mL plus 700 mL/ha MCPA LVE				
	4 to 5 leaf onwards	Volunteer chickpeas, lentils and safflower	Up to 6 leaf	Up to 4 leaf	150mL				
Volunteer faba beans and lupins								Maximum 10 cm high or 6 nodes	75mL plus 700 mL/ha MCPA LVE

CROP	CROP STAGE	WEED	WEED STAGE	RATE mL/ha	STATE	CRITICAL COMMENTS
Barley Oats Triticale Wheat	4 to 5 leaf onwards	Volunteer medics and seedling lucerne	Up to 8 leaf	150mL	NSW, Vic, SA, Tas and WA only	Weeds should be young and actively growing. Weeds will become stunted and will not be competitive soon after application although final results may not show for some weeks. Faba beans and lupins will only be suppressed. Use 75 mL/ha rate only in combination with MCPA LVE.
		Volunteer sub-clover	Up to 6 leaf			
		Volunteer vetch	Runners up to 10 cm maximum 16 leaf	100 mL		
	4 to 5 leaf onwards	Prickly lettuce	4 to 6 leaf and maximum 8 cm diameter	150mL plus 700 mL/ha MCPA LVE	NSW, Vic, SA, Tas and WA only	
		Thistles including: Nodding Saffron Scotch Slender Spear Stemless Variegated	Rosettes up to 10 cm maximum diameter	50mL plus 1.0 L/ha MCPA amine (500 g/L) or 50 plus 700 mL/ha MCPA LVE	NSW, Vic, SA, Tas, WA and Qld only	
	5 leaf to late tillering	Skeleton weed	5 to 15 cm rosettes	500 plus 1.0 L/ha MCPA amine 500 g/L	NSW, Vic and SA only	
Canola	2 to 8 leaf	Capeweed Saffron thistle Skeleton weed Soldier thistle	Up to 10 cm diameter (4 to 8 leaf)	300	NSW, Vic, SA, Tas, WA and Qld only	Weeds should be young and actively growing. Weeds will become stunted and will not be competitive soon after application although final results may not show for some weeks. Skeleton weed will only be controlled until harvest. Faba beans and lupins will only be suppressed. For the control of annual grasses, United Farmers Clopyralid Herbicide may be tank mixed with Verdict* Herbicide.
		Volunteer chickpeas, lentils and safflower	Up to 6 leaf	250		
		Volunteer faba beans and lupins	Up to 4 leaf	250		
		Volunteer field peas	Maximum 10 cm high or 6 nodes	150		
		Volunteer medics and seedling lucerne	Up to 8 leaf	150		
		Volunteer sub-clover	Up to 6 leaf	150		
		Volunteer vetch	Runners up to 10 cm maximum 16 leaf	100		

CROP	CROP STAGE	WEED	WEED STAGE	RATE mL/ha	STATE	CRITICAL COMMENTS
Pastures and fallow land	Post-emergence	Hardhead thistle (creeping knapweed, Russian knapweed)	Actively growing plants	Hand gun: 500 mL/100 L of water. Boom spray: 2 or 4 L/ha	Vic only	<p>NOTE: DO NOT USE ON LUCERNE. CLOVERS AND MEDICS WILL BE ELIMINATED FOR AT LEAST ONE YEAR.</p> <p>Victoria only: Use the lower rate only on light soils (sand and sandy loam) where a slightly lower degree of control is acceptable. Use the higher rate on all soil types where complete control is required. Addition of a wetting agent at label rates is recommended for treatment of hardhead thistle. Spray between September and April on actively growing plants for effective control. Thorough coverage is essential. Apply in 200 to 250 L of water/ha.</p> <p>BOOM SPRAYING: Use the higher rates of United Farmers Clopyralid Herbicide plus MCPA on multicrowned plants or rosettes larger than 30 cm in diameter. Spraying may be done at any time during active growth, usually in early winter or spring. Avoid spraying during the dormant winter period or at any time when thistles are not actively growing. <i>Do not spray flowering thistles.</i></p> <p>PRE-SPRAY MANAGEMENT: The pasture should be slightly grazed prior to spraying to reduce clover and grass cover and expose the smaller thistles to the spray. The grazed pasture should be left seven days to allow thistles to freshen prior to treatment.</p> <p>POST-TREATMENT MANAGEMENT: Response of thistles to treatment with the United Farmers Clopyralid Herbicide plus MCPA mixture will be slow compared to the standard treatments with 2,4-D or MCPA. If possible delay grazing of sprayed thistles for 14 days after treatment.</p> <p>CLOVER DAMAGE: The United Farmers Clopyralid Herbicide plus MCPA mixture can be very damaging to subterranean clover. The lower rate is no more damaging than label rates of 2,4-D or MCPA. The higher rate of the United Farmers Clopyralid Herbicide plus MCPA mixture will reduce the clover component of the pasture for about two months. Clover recovery will be quicker during periods of active growth.</p> <p>HANDGUN (Spot spray): Treat from rosette stage to early flowering. Thorough spraying is necessary.</p> <p>DRENCH GUN: Apply 10 mL to rosette crown. To multicrown plants, apply 10 mL to each crown.</p>
				Hand gun: 500 mL/100 L of water. Boom spray: 4 L/ha	Qld only	
		Thistles including: Nodding Variegated Scotch Spear Slender Saffron	Rosette stage prior to stem elongation. Treat rosette stage prior to stem elongation.	50 or 70mL/ha plus 1 to 1.5 L MCPA amine (500 g/L)/ ha Drench gun: 50 mL /1 L of water Hand gun: 250 mL/100 L of water	NSW, Vic, Tas, SA and Qld only	

CROP	CROP STAGE	WEED	WEED STAGE	RATE mL/ha	STATE	CRITICAL COMMENTS
Pastures and fallow land	Post-emergence	Nodding thistle	Rosettes up to 20 cm diameter	100mL	NSW only	Apply the spray from September to October. Apply by boom spray only. DO NOT apply to thistles over 20 cm in diameter. When thistles are over 20 cm in diameter use United Farmers Clopyralid Herbicide plus MCPA (referred to above). Clover Damage: Damage to white clover will be no greater than damage with MCPA alone and less than damage from United Farmers Clopyralid Herbicide plus MCPA mixtures. Damage to sub-clover may be greater than with MCPA or 2,4-D alone. DO NOT use for spot treatment.
		Californian thistle	From early buds to flowering (December to February)	Hand gun: 250 mL/100 L of water Boom spray: 2 L/ha	Vic and Tas only	Addition of a wetting agent at label rates is recommended. Retreatment of regrowth in the year following treatment will usually be necessary to achieve a high level of control. NOTE: Clovers and medics will be eliminated for at least one year.
Pasture, Forests, Rights-of-Way Industrial situations	Post-emergence	Groundsel bush	Young seedlings to mature plants	Hand gun: 330 or 500 mL/100 L of water	Qld and NSW only	Spray foliage when growth is active. Use the lower rate on young seedlings and the higher rate on plants more than 2 m tall or when growth is slow.
Pasture and <i>Pinus radiata</i> plantations		Silver wattle	Active growth spring to summer	Hand gun: 500 mL/100 L of water Boom and aerial spray: 5 L/ha (weeds 30 cm to 2 m) 7.0 L/ha (weeds 2 to 4 m) 8.5 L/ha (weeds 4 to 8 m)	NSW, Vic and Tas only	For effective control apply when bushes are growing actively. Large trees will not show complete necrosis. HANDGUN: Means high volume NOT low volume knapsack. Spray to give full coverage of leaves and stems. Add organosilicone surfactant at 200 mL/100 L for optimum results. See GENERAL INSTRUCTIONS for handgun spraying. BOOM AND AERIAL SPRAYING: For boom spraying apply in 150 to 200 L of water/ha. For aerial treatment apply in a minimum of 50 L/ha of water containing 25 to 50% by volume of anti-evaporant oil such as Ulvapron. Mix United Farmers Clopyralid Herbicide and water first and then add Ulvapron. Maintain continuous agitation. Clovers and other legumes will be eliminated for at least one year.
Forests		Cape ivy	Any growth stage	Hand gun: 3.3 L/ha	Vic and Tas only	Apply by hand held weed wiper or C.D.A. at dilutions with water at 1:3. Application may be made at any time of the year

CROP	CROP STAGE	WEED	WEED STAGE	RATE mL/ha	STATE	CRITICAL COMMENTS
						provided foliage is dry at the time. Avoid spraying non-target plants.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORIZED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

PASTURES:

DO NOT GRAZE OR CUT TREATED PASTURES FOR STOCK FEED FOR 7 DAYS AFTER APPLICATION OF RATES OF 4 L/HA OR LESS.

CEREALS:

DO NOT GRAZE OR CUT TREATED CEREALS FOR STOCK FEED FOR 4 WEEKS AFTER APPLICATION IF RATE IS IN EXCESS OF 300 mL/HA.

DO NOT APPLY LATER THAN 10 WEEKS BEFORE HARVEST.

CANOLA:

DO NOT HARVEST, GRAZE OR CUT FOR STOCK FEED EARLIER THAN TWELVE WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

MIXING:

Half fill the spray tank with water and add the required quantity of United Farmers Clopyralid Herbicide and complete filling. Agitate continuously to ensure thorough mixing before and during application. Only mix sufficient chemical for each day's work.

Tank mixtures: Wettable powder or dry flowable formulations should be added to the spray tank first, followed by suspension concentrates (flowables), aqueous concentrates (UFCC Clopyralid Herbicide) and the emulsifiable concentrate formulations (e.g. MCPA LVE.).

COMPATIBILITY:

United Farmers Clopyralid Herbicide is compatible with the following:

BROADLEAF HERBICIDES: Starane*, Metsulfuron Methyl, bromoxynil, Chlorsulfuron, Diuron, Glyphosate, MCPA amine, MCPA LVE, Paraquat, Spray.Seed*, terbutryn, 2,4-D Amine.

GRASS HERBICIDES IN CEREAL CROPS: Diclofop methyl, Grasp*, Puma* (Grasp and Puma for wild oat control only).

GRASS HERBICIDES ON BROADLEAF CROPS: Haloxyfop.

* Not United Farmers trademarks.

APPLICATION

BOOM SPRAYING CROP and PASTURES:

Apply United Farmers Clopyralid Herbicide in sufficient water to obtain good coverage. It should be applied by an accurately calibrated ground rig or aircraft, delivering 200 to 300 micron droplets and not less than 50 L/ha water volume for boom sprayers or not less than 20 L/ha for aerial applications.

Hardhead thistle - Use a spray volume of 200 to 250 L/ha of water.

Silver wattle - Use a spray volume of 150 to 200 L/ha of water by ground boomspray and a minimum spray volume of 50 L/ha by aircraft.

HIGH VOLUME HAND GUN:

Apply the recommended mix to give full coverage of leaves and stems through a No. 6-8 tip at 700 to 1500 kPa. Spray volume for effective coverage of dense two metre high silver wattle should be 30 to 40 litres of spray per 100 m² (10 m x 10 m) of infestation. For larger areas an equivalent would be 3000 to 4000 litres per infested hectare.

CLEANING SPRAY EQUIPMENT:

Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and water courses.

PARTIAL CLEANING (before spraying other labelled or tolerant crops):

After using United Farmers Clopyralid Herbicide, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose. Quarter fill the tank with clean water and circulate through the pump, line, hoses and nozzles. Drain and repeat procedure twice.

COMPLETE CLEANING (before spraying susceptible crops):

After using United Farmers Clopyralid Herbicide, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose. Quarter fill the tank with clean water and circulate as above, then drain.

Quarter fill the tank again and add a liquid alkali detergent (e.g. Surf, Omo, Drive) at 500 mL/100 L water and circulate throughout the system for at least fifteen minutes.

Drain, remove filters and nozzles and clean separately. Rinse inside the tank thoroughly using a pressure hose and flush system with clean water.

RESISTANT WEEDS WARNING

United Farmers Clopyralid Herbicide is a member of the Pyridines group of herbicides. The product has the disrupters of plant cell growth mode of action.

Group	I	Herbicide
--------------	----------	------------------

For weed resistance management, the product is a Group I herbicide.

Some naturally occurring weed biotypes resistant to the product and other disrupters of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by the product or other disrupters of plant cell growth herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, United Farmers Cooperative Limited accepts no liability for any losses that may result from the failure of the product to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Do not apply under weather conditions, or from spraying equipment that may cause spray drift onto nearby susceptible plants/crops, cropping lands or pastures.

Susceptible crops include chickpeas, cotton, faba beans, field peas, fruit trees, lentils, lupins, lucerne, medics, ornamentals, potatoes, safflower, sub-clover, tomatoes, vegetables, vines, wattles and white clover.

Do not apply United Farmers Clopyralid Herbicide to crops or pastures, which are to be used for the production of compost or mulches for use with susceptible crops or plants. The use of straw, hay or other plant material treated with United Farmers Clopyralid Herbicide for composting or mulching susceptible crop may damage these crops.

Plant back periods: Residues in the straw of treated crops can affect subsequent susceptible crops. Susceptible crops listed above should not be sown for: Nine months following any application up to 300 mL/ha. Twelve months following an application in excess of 300 mL/ha and up to 500 mL/ha.

Note: Field peas are particularly susceptible and should not be sown the season following an application of 500 mL/ha. Where rates in excess of 500 mL/ha have been used, susceptible crops, including field peas, should not be sown for at least two years.

PROTECTION OF LIVESTOCK

DO NOT graze or cut treated crops for stock food except as specified under WITHHOLDING PERIODS.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

United Farmers Clopyralid Herbicide has low toxicity to fish, birds, honey bees, livestock, earthworms and aquatic organisms.

DO NOT contaminate streams, rivers or waterways with chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

DO NOT store near feedstuffs, fertilisers or seed.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit

specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SMALL SPILL MANAGEMENT

Wear protective equipment (See SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal. (See STORAGE AND DISPOSAL section). If necessary, wash the spill area with an alkali detergent and water and absorb as above, the wash liquid for disposal.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin.

DO NOT inhale the spray mist.

When preparing the spray, wear elbow-length PVC gloves and a face shield.

After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

After each day's use, wash face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. (Telephone Australia 13 1126)

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

CONDITIONS OF SALE

The use of United Farmers Clopyralid Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by United Farmers Cooperative regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and United Farmers Cooperative accepts no responsibility for any consequence whatsoever resulting from the use of this product.

UNITED FARMERS COOPERATIVE LIMITED.
ACN 14 014 213 492
2 Birksgate Road, PMB 10
ROUS HEAD NORTH FREMANTLE WA 6105
Phone (08) 9430 2222 Fax: (08) 9430 2233

<p>IN EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE</p>	<p>FOR SPECIALIST ADVICE IN AN EMERGENCY CALL 1800 705 766 (24 HOURS)</p>
--	--