

100% on A3

YOUR PROOF

job#: 172491
 name: 02 Clodinafop 1L Base
 client: Genfarm
 date: 21/01/2010

This proof is **NOT** a true representation of the final colours that will be printed

OK to proceed
 Further proof required
 sign: _____
 date: _____

NO responsibility will be taken for errors not noted on this signed proof. PLEASE EXAMINE THIS PROOF VERY CAREFULLY AND RETURN SIGNED PROOF TO US FOR JOB TO PROCEED

PHILSTIC® **Philstic Labels Ltd**
 Phone +64 9 630 5822
 Fax +64 9 630 0957

 Dieline

 Pantone Violet

 PMS 348

 PMS 654

 Black

 Top Area (130mm x 130mm)

 Book area (128mm x 100mm)

POISON
 KEEP OUT OF REACH OF CHILDREN
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Genfarm
CLODINA FOP 240EC
 SELECTIVE HERBICIDE

ACTIVE CONSTITUENTS: 240g/L CLODINA FOP-PROPARGYL
 60g/L CLOQUINTOCET-MEXYL
 SOLVENTS: 507 g/L LIQUID HYDROCARBONS
 100 g/L N-METHYL-PYRROLIDONE

GROUP A HERBICIDE

For the control of Wild Oats, Paradoxa Grass (Annual Phalaris), Canary Grass and Annual Ryegrass in Wheat. Highly active on Wild Oats and can be mixed with many broadleaf herbicides as per the Directions for Use Table.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE



CONTENTS: 1 LITRES



PHILSTIC MULTIFOLD

Genfarm

CLODINA FOP 240EC

SELECTIVE HERBICIDE

STORAGE AND DISPOSAL
 Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS
 Harmful if swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid inhaling spray mist. Avoid contact with eyes and skin. When opening the container, preparing spray and using the product wear:
 • cotton overalls buttoned to the neck and wrist (or equivalent clothing);
 • a washable hat;
 • elbow-length butyl rubber gloves; and
 • face shield or goggles.
 If product gets on skin, immediately wash area with soap and water. If product gets in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID
 If poisoning occurs contact a doctor or Poisons Information Centre. Phone 131 126. If swallowed do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open. Flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET
 For further information, refer to the Material Safety Data Sheet (MSDS) which is available from the supplier.

CONDITIONS OF SALE
 The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Genfarm Crop Protection Pty Ltd. accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd. will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

IN A TRANSPORT EMERGENCY
DIAL 000
 POLICE OR FIRE BRIGADE

FOR SPECIALIST ADVICE
IN AN EMERGENCY DIAL
1800 033 111
 24 HOURS AUSTRALIA WIDE

APVMA Approval No: 61274/1L/0110
 Genfarm Crop Protection Pty Ltd
 Suite 2, Level 3, 64 Talavera Road,
 Macquarie Park NSW 2113
 Tel: (02) 9889 5400
 Product Support Tel: 1800 44 88 92



255mm wide

127mm deep

PEEL HERE

Genfarm

CLODINAPOP 240EC

SELECTIVE HERBICIDE

STORAGE AND DISPOSAL
Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.
Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

MATERIAL SAFETY DATA SHEET
For further information, refer to the Material Safety Data Sheet (MSDS) which is available from the supplier.

CONDITIONS OF SALE
The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Genfarm Crop Protection Pty Ltd. accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd. will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

SAFETY DIRECTIONS
Harmful if swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid inhaling spray mist. Avoid contact with eyes and skin. When opening the container, preparing spray and using the product wear:
• cotton overalls buttoned to the neck and wrist (or equivalent clothing);
• a washable hat;
• elbow-length butyl rubber gloves; and
• face shield or goggles.
If product gets on skin, immediately wash area with soap and water. If product gets in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID
If poisoning occurs contact a doctor or Poisons Information Centre. Phone 131 126. If swallowed do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open. Flood with water for at least 15 minutes and see a doctor.

APVMA Approval No: 61274/1L/0110
Genfarm Crop Protection Pty Ltd
Suite 2, Level 3, 64 Talavera Road,
Macquarie Park NSW 2113
Tel: (02) 9889 5400
Product Support Tel: 1800 44 88 92

**IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE**

**FOR SPECIALIST ADVICE
IN AN EMERGENCY DIAL
1800 033 111
24 HOURS AUSTRALIA WIDE**

Batch No: _____
Date of Manufacture: _____


9 335371 001396



130mm deep

130mm wide

100% on A3

YOUR PROOF






job#: 172491
name: 02 Clodinafop 1L Top
client: Genfarm
date: 21/01/2010

This proof is **NOT** a true representation of the final colours that will be printed

OK to proceed
 Further proof required
sign: _____
date: _____

NO responsibility will be taken for errors not noted on this signed proof.
PLEASE EXAMINE THIS PROOF VERY CAREFULLY AND RETURN SIGNED PROOF TO US FOR JOB TO PROCEED

PHILSTIC® **Philstic Labels Ltd**
Phone +64 9 630 5822
Fax +64 9 630 0957

-  Dieline
-  Pantone Violet
-  PMS 348
-  PMS 654
-  Black

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
GENFARM CLODINAFOP 240EC
SELECTIVE HERBICIDE

ACTIVE CONSTITUENTS: 240g/L CLODINAFOP-PROPARGYL
60g/L CLOQUINTOCET-MAXYL
SOLVENTS: 507g/L g/L LIQUID HYDROCARBONS
100g/L N-METHYL-PYRROLIDONE

GROUP A HERBICIDE

For the control of Wild Oats, Paradoxa Grass (Annual Phalaris), Canary Grass and Annual Ryegrass in Wheat. Highly active on Wild Oats and can be mixed with many broadleaf herbicides as per the Directions for Use Table.

THIS BOOKLET IS PART OF THE LABEL

APVMA APPROVAL NO: 61274/1leaf/0110

**Genfarm Crop Protection Pty Ltd
Suite 2, Level 3, 64 Talavera Road,
Macquarie Park, NSW, 2113
Tel: (02) 9889 5400
Product Support Tel: 1800 44 88 92**

DIRECTIONS FOR USE

Restrains: DO NOT apply if rainfall is expected within 2 hours.
 DO NOT apply to weeds or crops which are under stress. For example, due to very dry, very wet, frosty or diseased conditions, nutrient deficiency or high insect pressure.
 DO NOT apply to weeds or crops immediately after a prolonged period of frosts.
 DO NOT apply under poor growing conditions.

CROP	WEEDS CONTROLLED	STATE	RATE PER HECTARE
Wheat 2 leaf to late jointing (Zadocks 12 – 37)	Wild Oats	Nthn NSW, Qld only	65 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water
			85 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water
			85 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water
			125 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water

WEED GROWTH STAGE	CRITICAL COMMENTS
1 to 5 leaf	NOT TO BE MIXED WITH BROADLEAF HERBICIDES AT THESE RATES. Apply at 85 mL/ha when weeds are not actively growing.
2 leaf to mid tillering.	Where low water volumes are used. DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.
2 to 5 leaf.	MAY BE TANK MIXED WITH COMPATIBLE HERBICIDES. Apply in mixtures for broadleaf weed control only when weeds are actively growing. When mixing GENFARM CLODINAFOP 240EC with certain broadleaf weed herbicides some reduction in grass weed control may occur.
2 leaf to mid tillering.	Unless otherwise specified always use D-C-Trate, Uptake or Hasten when tank mixing broadleaf herbicides. Apply at 125 mL/ha when weeds are not actively growing. Where low water volumes are used. DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.

CROP	WEEDS CONTROLLED	STATE	RATE PER HECTARE
Wheat 2 leaf to late jointing (Zadocks 12 – 37)	Wild Oats	Sthn NSW, VIC, SA, WA only	65 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water. Or 100 mL of a 100% non-ionic surfactant per 100 L of water
			85 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water. Or 100 mL of a 100% non-ionic surfactant per 100 L of water.
	Paradoxa Grass (<i>Phalaris paradoxa</i>) Canary Grass (<i>Phalaris minor</i>)	QLD, NSW, VIC, SA only	85 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.
			125 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.
			160 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.

WEED GROWTH STAGE	CRITICAL COMMENTS
2 to 5 leaf.	MAY BE TANK MIXED WITH COMPATIBLE HERBICIDES. Apply in mixtures for broadleaf weed control only when weeds are actively growing. When mixing GENFARM CLODINAFOP 240EC with certain broadleaf weed herbicides some reduction in grass weed control may occur.
2 leaf to mid tillering.	Refer to compatibility chart for specific rates for tank mixes. Unless otherwise specified always use D-C-Trate, Uptake or Hasten when tank mixing broadleaf herbicides. Apply at 85 mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.
2 to 3 leaf	NOT TO BE MIXED WITH BROADLEAF HERBICIDES AT BELOW 160 mL/ha.
2 to 5 leaf	Apply in mixtures for broadleaf weed control only when weeds are actively growing. When mixing GENFARM CLODINAFOP 240EC with certain broadleaf weed herbicides some reduction in grass weed control may occur.
2 leaf to early tillering	Apply at 160 mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.

CROP	WEEDS CONTROLLED	STATE	RATE PER HECTARE
Wheat 2 leaf to late jointing (Zadocks 12 – 37)	Annual Ryegrass	NSW, VIC, SA, WA only	160 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.
			210 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS

WITHHOLDING PERIOD:

Harvested grain:

WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.

Animal grazing/feeding:

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

GENFARM CLODINAFOP 240 EC Selective Herbicide should be applied only when weeds are actively growing.

Resistant Weeds Warning



GENFARM CLODINAFOP 240EC Selective Herbicide is a member of the aryloxyphenoxypropionate (“fop”) group of herbicides. GENFARM CLODINAFOP 240EC has the inhibition of fat (lipid) synthesis, (or inhibitors of acetyl CoA carboxylase) mode of action. For weed resistance management GENFARM CLODINAFOP 240EC is a Group A herbicide.

WEED GROWTH STAGE	CRITICAL COMMENTS
2 to 4 leaf	<p>MAY BE TANK MIXED WITH COMPATIBLE HERBICIDES.</p> <p>Apply in mixtures for broadleaf weed control only when weeds are actively growing. When mixing GENFARM CLODINAFOP 240EC with certain broadleaf weed herbicides some reduction in grass weed control may occur.</p> <p>Apply at 210 mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes).</p> <p>Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.</p>
2 leaf to early tillering	

LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

Some naturally-occurring weed biotypes resistant to GENFARM CLODINAFOP 240EC and other Group A herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by GENFARM CLODINAFOP 240EC or other Group A herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Genfarm Crop Protection Pty Ltd accepts no liability for any losses that may result from the failure of GENFARM CLODINAFOP 240EC to control resistant weeds.

Weed biotypes resistant to diclofop-methyl and other “grass specific” herbicides can also often be resistant to GENFARM CLODINAFOP 240EC. Before using GENFARM CLODINAFOP 240EC on a population resistant to or suspected to be resistant to “grass specific” herbicides, have a resistance test conducted to ensure that it is susceptible to GENFARM CLODINAFOP 240EC.

Large numbers of healthy surviving weeds can be an indication that resistance is developing. Efforts should be made to prevent seed set of the surviving weeds. DO NOT make more than one application of a Group A herbicide with the inhibition of fat (lipid) synthesis (or inhibitors of acetyl CoA carboxylase) mode-of-action to a crop in the same season.

If the user suspects that the target weed population is resistant to herbicides with this mode of action, GENFARM CLODINAFOP 240EC or other Group A herbicides should not be used.

Strategies to minimize the risk of herbicide resistance are available. The above recommendations should be incorporated into an Integrated Weed Management (IWM) Program. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, or local Genfarm Crop Protection representative for details.

Mixing

Fill the spray tank to one quarter full. Then add GENFARM CLODINAFOP 240EC while adding water to make up the final spray volume. Pour GENFARM CLODINAFOP 240EC into the stream of incoming water. Good agitation at the time GENFARM CLODINAFOP 240EC is added to the tank will ensure good mixing. Add the required amount of D-C-Trate*, Hasten*, or Uptake* oil just before the tank is full of water and with the agitators in motion.

Mixing order

Some products may react with other products if they are not mixed in the correct order. The general mixing order of products should be –

- I. water conditioners or buffers
- II. water dispersible granules (WG)
- III. wettable powders (WP)
- IV. flowable or suspension concentrates (SC)
- V. emulsifiable concentrates (EC) (GENFARM CLODINAFOP 240EC)
- VI. water based or soluble concentrates
- VII. oils and wetters – eg Hasten, BS 1000

It is important to ensure that each individual component of the tank mix is fully dissolved and in solution before the next product is added to the tank mix, otherwise mixing problems may occur.

Minimum Oil Volumes

The minimum amount of oil to be applied with GENFARM CLODINAFOP 240EC will depend on the type of oil used. When applying GENFARM CLODINAFOP 240EC using low water volumes, DO NOT use less than the minimum oil volumes specified below:

- For oils normally applied at 1% v/v (eg D-C-Trate), DO NOT apply less than 500 mL oil/ha.
- For oils normally applied at 0.5% v/v (eg Uptake, Hasten), DO NOT apply less than 250 mL oil/ha.

Cleaning Spray Equipment

After using GENFARM CLODINAFOP 240EC, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose, drain the tank and clean any filters in the tank, pump, line and nozzles.

To rinse: After cleaning the tank as above, quarter fill the tank with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

To decontaminate: Before spraying cereals (except wheat), maize, sorghum or other sensitive crops, wash the tank and rinse the system as above. Drain and repeat the rinsing procedure. Then quarter fill the tank and add a liquid alkali detergent (eg SURF*, OMO*, DRIVE*) at 0.5 L/100 L of water and circulate throughout the system for at least fifteen minutes. Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain. Dispose of all water used for cleaning in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots.

Application

Ground Application: Boom spray with flat fan nozzles. Ensure good spray coverage is obtained. Apply using 50 to 110 L water per hectare. The ideal droplet size is 200 to 300 microns VMD.

Aircraft Application: For best results use 20 to 30 L water per hectare and spray at 2 m to 3 m above the crop. Ideal droplet size is 250 to 350 microns VMD. For rotary atomizers (micronairs) this can be achieved using blade angles of 55° to 65°. Where booms and nozzles are attached, flat fan nozzles with a spray angle of 65° to 90° should be used.

Avoid applying GENFARM CLODINAFOP 240EC if wind speeds are greater than 5 m/s. If spraying in calm conditions use larger droplets and reduce flying height.

Compatability

When GENFARM CLODINAFOP 240EC is applied with compatible herbicides always use a recommended oil. Only use a 100% non-ionic surfactant when mixing with diflufenican based products.

Grass Herbicides: GENFARM CLODINAFOP 240EC is compatible with diclofop methyl, eg Genfarm Diclofop-Methyl 375 Selective Herbicide.

Broadleaf Herbicides: Some broadleaf weed herbicides are incompatible with GENFARM CLODINAFOP 240EC because they reduce the effectiveness of GENFARM CLODINAFOP 240EC or because in mixture they may cause crop yellowing. For the latest information on compatibility of GENFARM CLODINAFOP 240EC with broadleaf herbicides refer to your reseller or Genfarm Crop Protection Pty Ltd.

Incompatible:

The following herbicides are known to be incompatible with GENFARM CLODINAFOP 240EC:

Diuron, Triasulfuron, Barrel* (bromoxynil + dicamba + MCPA) and all 2,4-D formulations.

Herbicides which are incompatible with GENFARM CLODINAFOP 240EC must be applied separately. In this situation, apply GENFARM CLODINAFOP 240EC first and then allow at least 10 days between its application and application of the broadleaf weed herbicide.

Insecticides: GENFARM CLODINAFOP 240EC is compatible with methidathion.

PRECAUTIONS

Re-entry Period

DO NOT enter treated areas without protective clothing until the spray has dried.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply on or near shrubs, trees, lawns or crops other than wheat. DO NOT drain or flush equipment on or near desirable trees or other plants, where their roots may extend, or in situations where by movement of soil or by seepage absorption of the herbicide may occur.

DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto nearby susceptible plants, adjacent crops, crop lands or pastures.

DO NOT allow spray to drift onto adjacent fallow land.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT apply to irrigation drains or channels.

DO NOT contaminate dams, streams, creeks and rivers with the chemical, spray mixture, used containers or equipment washings.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid inhaling spray mist. Avoid contact with eyes and skin.

When opening the container, preparing spray and using the product wear:

- cotton overalls buttoned to the neck and wrist (or equivalent clothing);
- a washable hat;
- elbow-length butyl rubber gloves; and
- face shield or goggles.

If product gets on skin, immediately wash area with soap and water. If product gets in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone 131 126.

If swallowed do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open. Flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET

For further information, refer to the Material Safety Data Sheet (MSDS) which is available from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Genfarm Crop Protection Pty Ltd. accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd. will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product

Genfarm Crop Protection Pty Ltd
Suite 2, Level 3, 64 Talavera Road,
Macquarie Park,
NSW, 2113
Tel: (02) 9889 5400

THIS PAGE DOES NOT PRINT

Pages @ 100% on A3

100% on A3

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Genfarm

CLODINAPOP 240EC

SELECTIVE HERBICIDE

ACTIVE CONSTITUENTS: 240 g/L CLODINAPOP-PROPARGYL
60 g/L CLOQUINTOCET-MEXYL
SOLVENTS: 507 g/L LIQUID HYDROCARBONS
100 g/L N-METHYL-PYRROLIDONE


GROUP A HERBICIDE

For the control of Wild Oats, Paradoxa Grass (Annual Phalaris), Canary Grass and Annual Ryegrass in Wheat. Highly active on Wild Oats and can be mixed with many broadleaf herbicides as per the Directions for Use Table.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE

Genfarm
CONTENTS: 5 LITRES

APVMA APPROVED LABEL



140mm deep

140mm wide

YOUR PROOF

job#: 172491
name: 00 Clodinafop 240EC 5L Front
client: Genfarm
date: 21/01/2010

This proof is **NOT** a true representation of the final colours that will be printed

OK to proceed
 Further proof required
sign: _____
date: _____

NO responsibility will be taken for errors not noted on this signed proof. PLEASE EXAMINE THIS PROOF VERY CAREFULLY AND RETURN SIGNED PROOF TO US FOR JOB TO PROCEED

PHILSTIC Philstic Labels Ltd
Phone +64 9 630 5822
Fax +64 9 630 0957

 Dieline

 Pantone Violet

 PMS 348

 PMS 654

 Black

100% on A3

PHILSTIC MULTIFOLD

Genfarm

CLODINAPOP 240EC

SELECTIVE HERBICIDE

STORAGE AND DISPOSAL
Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.
Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS
Harmful if swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid inhaling spray mist. Avoid contact with eyes and skin. When opening the container, preparing spray and using the product wear:

- cotton overalls buttoned to the neck and wrist (or equivalent clothing);
- a washable hat;
- elbow-length butyl rubber gloves; and
- face shield or goggles.

If product gets on skin, immediately wash area with soap and water. If product gets in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

APVMA Approval No: 61274/5L/0110
Genfarm Crop Protection Pty Ltd
Suite 2, Level 3, 64 Talavera Road,
Macquarie Park NSW 2113
Tel: (02) 9889 5400
Product Support Tel: 1800 44 88 92

FIRST AID
If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131 126. If swallowed do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open. Flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET
For further information, refer to the Material Safety Data Sheet (MSDS) which is available from the supplier.

CONDITIONS OF SALE
The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Genfarm Crop Protection Pty Ltd. accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd. will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE

FOR SPECIALIST ADVICE
IN AN EMERGENCY DIAL
1800 033 111
24 HOURS AUSTRALIA WIDE





137mm deep

139mm wide

YOUR PROOF

job#: 172491
 name: 01 Clodinafop 240EC 5L Base
 client: Genfarm
 date: 21/01/2010

This proof is **NOT** a true representation of the final colours that will be printed

OK to proceed
 Further proof required
 sign: _____
 date: _____

NO responsibility will be taken for errors not noted on this signed proof.

PLEASE EXAMINE THIS PROOF VERY CAREFULLY AND RETURN SIGNED PROOF TO US FOR JOB TO PROCEED

PHILSTIC Philstic Labels Ltd
 Phone +64 9 630 5822
 Fax +64 9 630 0957

 Book area (138mm x 110mm)

 Dieline

 Black

100% on A3

PEEL HERE

Genfarm


CLODINAPOP 240EC

SELECTIVE HERBICIDE

STORAGE AND DISPOSAL
Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS
Harmful if swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid inhaling spray mist. Avoid contact with eyes and skin. When opening the container, preparing spray and using the product wear:
• cotton overalls buttoned to the neck and wrist (or equivalent clothing);
• a washable hat;
• elbow-length butyl rubber gloves; and
• face shield or goggles.
If product gets on skin, immediately wash area with soap and water. If product gets in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

APVMA Approval No: 61274/5L/0110
Genfarm Crop Protection Pty Ltd
Suite 2, Level 3, 64 Talavera Road,
Macquarie Park NSW 2113
Tel: (02) 9889 5400
Product Support Tel: 1800 44 88 92



FIRST AID
If poisoning occurs contact a doctor or Poisons Information Centre. Phone 131 126. If swallowed do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open. Flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET
For further information, refer to the Material Safety Data Sheet (MSDS) which is available from the supplier.

CONDITIONS OF SALE
The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Genfarm Crop Protection Pty Ltd. accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd. will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE

FOR SPECIALIST ADVICE
IN AN EMERGENCY DIAL
1800 033 111
24 HOURS AUSTRALIA WIDE

Batch No:

Date of Manufacture:



9 335371 001402



140mm deep

140mm wide

YOUR PROOF

job#: 172491
 name: 02 Clodinafop 240EC 5L Front
 client: Genfarm
 date: 21/01/2010

This proof is **NOT** a true representation of the final colours that will be printed

OK to proceed
 Further proof required
 sign: _____
 date: _____

NO responsibility will be taken for errors not noted on this signed proof. PLEASE EXAMINE THIS PROOF VERY CAREFULLY AND RETURN SIGNED PROOF TO US FOR JOB TO PROCEED

PHILSTIC® **Philstic Labels Ltd**
 Phone +64 9 630 5822
 Fax +64 9 630 0957

- Dieline
- Pantone Violet
- PMS 348
- PMS 654
- Black

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
GENFARM CLODINAFOP 240EC
SELECTIVE HERBICIDE

ACTIVE CONSTITUENTS: 240g/L CLODINAFOP-PROPARGYL
60g/L CLOQUINTOCET-MAXYL
SOLVENTS: 507g/L g/L LIQUID HYDROCARBONS
100g/L N-METHYL-PYRROLIDONE

GROUP A HERBICIDE

For the control of Wild Oats, Paradoxa Grass (Annual Phalaris), Canary Grass and Annual Ryegrass in Wheat. Highly active on Wild Oats and can be mixed with many broadleaf herbicides as per the Directions for Use Table.

THIS BOOKLET IS PART OF THE LABEL

APVMA APPROVAL NO: 61274/5leaf/0110

**Genfarm Crop Protection Pty Ltd
Suite 2, Level 3, 64 Talavera Road,
Macquarie Park, NSW, 2113
Tel: (02) 9889 5400
Product Support Tel: 1800 44 88 92**

DIRECTIONS FOR USE

Restrains: DO NOT apply if rainfall is expected within 2 hours.

DO NOT apply to weeds or crops which are under stress. For example, due to very dry, very wet, frosty or diseased conditions, nutrient deficiency or high insect pressure.

DO NOT apply to weeds or crops immediately after a prolonged period of frosts.

DO NOT apply under poor growing conditions.

CROP	WEEDS CONTROLLED	STATE	RATE PER HECTARE
Wheat 2 leaf to late jointing (Zadocks 12 – 37)	Wild Oats	Nthn NSW, Qld only	65 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water
			85 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water
			85 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water
			125 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water

WEED GROWTH STAGE	CRITICAL COMMENTS
1 to 5 leaf	NOT TO BE MIXED WITH BROADLEAF HERBICIDES AT THESE RATES. Apply at 85 mL/ha when weeds are not actively growing. Where low water volumes are used. DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.
2 leaf to mid tillering.	
2 to 5 leaf.	MAY BE TANK MIXED WITH COMPATIBLE HERBICIDES. Apply in mixtures for broadleaf weed control only when weeds are actively growing. When mixing GENFARM CLODINAPOP 240EC with certain broadleaf weed herbicides some reduction in grass weed control may occur. Unless otherwise specified always use D-C-Trate, Uptake or Hasten when tank mixing broadleaf herbicides. Apply at 125 mL/ha when weeds are not actively growing. Where low water volumes are used. DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.
2 leaf to mid tillering.	

CROP	WEEDS CONTROLLED	STATE	RATE PER HECTARE
Wheat 2 leaf to late jointing (Zadocks 12 – 37)	Wild Oats	Sthn NSW, VIC, SA, WA only	65 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water. Or 100 mL of a 100% non-ionic surfactant per 100 L of water
			85 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water. Or 100 mL of a 100% non-ionic surfactant per 100 L of water.
	Paradoxa Grass (<i>Phalaris paradoxa</i>) Canary Grass (<i>Phalaris minor</i>)	QLD, NSW, VIC, SA only	85 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.
			125 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.
			160 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.

WEED GROWTH STAGE	CRITICAL COMMENTS
2 to 5 leaf.	MAY BE TANK MIXED WITH COMPATIBLE HERBICIDES. Apply in mixtures for broadleaf weed control only when weeds are actively growing. When mixing GENFARM CLODINAFOP 240EC with certain broadleaf weed herbicides some reduction in grass weed control may occur.
2 leaf to mid tillering.	Refer to compatibility chart for specific rates for tank mixes. Unless otherwise specified always use D-C-Trate, Uptake or Hasten when tank mixing broadleaf herbicides. Apply at 85 mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.
2 to 3 leaf	NOT TO BE MIXED WITH BROADLEAF HERBICIDES AT BELOW 160 mL/ha. Apply in mixtures for broadleaf weed control only when weeds are actively growing. When mixing GENFARM CLODINAFOP 240EC with certain broadleaf weed herbicides some reduction in grass weed control may occur.
2 to 5 leaf	Apply at 160 mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.
2 leaf to early tillering	Apply at 160 mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.

CROP	WEEDS CONTROLLED	STATE	RATE PER HECTARE
Wheat 2 leaf to late jointing (Zadocks 12 – 37)	Annual Ryegrass	NSW, VIC, SA, WA only	160 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.
			210 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL

WITHHOLDING PERIOD:

Harvested grain: WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.

Animal grazing/feeding: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

GENFARM CLODINAFOP 240 EC Selective Herbicide should be applied only when weeds are actively growing.

Resistant Weeds Warning



GENFARM CLODINAFOP 240EC Selective Herbicide is a member of the aryloxyphenoxypropionate (“fop”) group of herbicides. GENFARM CLODINAFOP 240EC has the inhibition of fat (lipid) synthesis, (or inhibitors of acetyl CoA carboxylase) mode of action. For weed resistance management GENFARM CLODINAFOP 240EC is a Group A herbicide. Some naturally-occurring weed biotypes resistant to GENFARM CLODINAFOP 240EC and other Group A herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these

WEED GROWTH STAGE	CRITICAL COMMENTS
2 to 4 leaf	<p>MAY BE TANK MIXED WITH COMPATIBLE HERBICIDES. Apply in mixtures for broadleaf weed control only when weeds are actively growing. When mixing GENFARM CLODINAFOP 240EC with certain broadleaf weed herbicides some reduction in grass weed control may occur. Apply at 210 mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.</p>
2 leaf to early tillering	

UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

herbicides are used repeatedly. These resistant weeds will not be controlled by GENFARM CLODINAFOP 240EC or other Group A herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Genfarm Crop Protection Pty Ltd accepts no liability for any losses that may result from the failure of GENFARM CLODINAFOP 240EC to control resistant weeds.

Weed biotypes resistant to diclofop-methyl and other “grass specific” herbicides can also often be resistant to GENFARM CLODINAFOP 240EC. Before using GENFARM CLODINAFOP 240EC on a population resistant to or suspected to be resistant to “grass specific” herbicides, have a resistance test conducted to ensure that it is susceptible to GENFARM CLODINAFOP 240EC.

Large numbers of healthy surviving weeds can be an indication that resistance is developing. Efforts should be made to prevent seed set of the surviving weeds.

DO NOT make more than one application of a Group A herbicide with the inhibition of fat (lipid) synthesis (or inhibitors of acetyl CoA carboxylase) mode-of-action to a crop in the same season.

If the user suspects that the target weed population is resistant to herbicides with this mode of action, GENFARM CLODINAFOP 240EC or other Group A herbicides should not be used. Strategies to minimize the risk of herbicide resistance are available. The above recommendations should be incorporated into an Integrated Weed Management (IWM) Program. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, or local Genfarm Crop Protection representative for details.

Mixing

Fill the spray tank to one quarter full. Then add GENFARM CLODINAFOP 240EC while adding water to make up the final spray volume. Pour GENFARM CLODINAFOP 240EC into the stream of incoming water. Good agitation at the time GENFARM CLODINAFOP 240EC is added to the tank will ensure good mixing. Add the required amount of D-C-Trate*, Hasten*, or Uptake* oil just before the tank is full of water and with the agitators in motion.

Mixing order

Some products may react with other products if they are not mixed in the correct order. The general mixing order of products should be –

- I. water conditioners or buffers
- II. water dispersable granules (WG)
- III. wettable powders (WP)
- IV. flowable or suspension concentrates (SC)
- V. emulsifiable concentrates (EC) (GENFARM CLODINAFOP 240EC)
- VI. water based or soluble concentrates
- VII. oils and wetters – eg Hasten, BS 1000

It is important to ensure that each individual component of the tank mix is fully dissolved and in solution before the next product is added to the tank mix, otherwise mixing problems may occur.

Minimum Oil Volumes

The minimum amount of oil to be applied with GENFARM CLODINAFOP 240EC will depend on the type of oil used. When applying GENFARM CLODINAFOP 240EC using low water volumes, DO NOT use less than the minimum oil volumes specified below:

- For oils normally applied at 1% v/v (eg D-C-Trate), DO NOT apply less than 500 mL oil/ha.
- For oils normally applied at 0.5% v/v (eg Uptake, Hasten), DO NOT apply less than 250 mL oil/ha.

Cleaning Spray Equipment

After using GENFARM CLODINAFOP 240EC, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose, drain the tank and clean any filters in the tank, pump, line and nozzles.

To rinse: After cleaning the tank as above, quarter fill the tank with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

To decontaminate: Before spraying cereals (except wheat), maize, sorghum or other sensitive crops, wash the tank and rinse the system as above. Drain and repeat the rinsing procedure. Then quarter fill the tank and add a liquid alkali detergent (eg SURF*, OMO*,

10

DRIVE*) at 0.5 L/100 L of water and circulate throughout the system for at least fifteen minutes. Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain. Dispose of all water used for cleaning in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots.

Application

Ground Application: Boom spray with flat fan nozzles. Ensure good spray coverage is obtained. Apply using 50 to 110 L water per hectare. The ideal droplet size is 200 to 300 microns VMD.

Aircraft Application: For best results use 20 to 30 L water per hectare and spray at 2 m to 3 m above the crop. Ideal droplet size is 250 to 350 microns VMD. For rotary atomizers (micronairs) this can be achieved using blade angles of 55° to 65°. Where booms and nozzles are attached, flat fan nozzles with a spray angle of 65° to 90° should be used. Avoid applying GENFARM CLODINAFOP 240EC if wind speeds are greater than 5 m/s. If spraying in calm conditions use larger droplets and reduce flying height.

Compatibility

When GENFARM CLODINAFOP 240EC is applied with compatible herbicides always use a recommended oil. Only use a 100% non-ionic surfactant when mixing with diflufenican based products.

Grass Herbicides: GENFARM CLODINAFOP 240EC is compatible with diclofop methyl, eg Genfarm Diclofop-Methyl 375 Selective Herbicide.

Broadleaf Herbicides: Some broadleaf weed herbicides are incompatible with GENFARM CLODINAFOP 240EC because they reduce the effectiveness of GENFARM CLODINAFOP 240EC or because in mixture they may cause crop yellowing. For the latest information on compatibility of GENFARM CLODINAFOP 240EC with broadleaf herbicides refer to your reseller or Genfarm Crop Protection Pty Ltd.

Incompatible:

The following herbicides are known to be incompatible with GENFARM CLODINAFOP 240EC: Diuron, Triasulfuron, Barrel* (bromoxynil + dicamba + MCPA) and all 2,4-D formulations. Herbicides which are incompatible with GENFARM CLODINAFOP 240EC must be applied separately. In this situation, apply GENFARM CLODINAFOP 240EC first and then allow at least 10 days between its application and application of the broadleaf weed herbicide.

Insecticides: GENFARM CLODINAFOP 240EC is compatible with methidathion.

PRECAUTIONS

Re-entry Period

DO NOT enter treated areas without protective clothing until the spray has dried.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply on or near shrubs, trees, lawns or crops other than wheat. DO NOT drain or flush equipment on or near desirable trees or other plants, where their roots may extend, or in situations where by movement of soil or by seepage absorption of the herbicide may occur. DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto nearby susceptible plants, adjacent crops, crop lands or pastures.

DO NOT allow spray to drift onto adjacent fallow land.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT apply to irrigation drains or channels.

DO NOT contaminate dams, streams, creeks and rivers with the chemical, spray mixture, used containers or equipment washings.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank.

DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid inhaling spray mist. Avoid contact with eyes and skin. When opening the container, preparing spray and using the product wear:

- cotton overalls buttoned to the neck and wrist (or equivalent clothing);
- a washable hat;
- elbow-length butyl rubber gloves; and
- face shield or goggles.

If product gets on skin, immediately wash area with soap and water. If product gets in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone 131 126.

If swallowed do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open.

Flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET

For further information, refer to the Material Safety Data Sheet (MSDS) which is available from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Genfarm Crop Protection Pty Ltd. accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd. will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product

Genfarm Crop Protection Pty Ltd
Suite 2, Level 3, 64 Talavera Road,
Macquarie Park,
NSW, 2113
Tel: (02) 9889 5400

THIS PAGE DOES NOT PRINT

Pages @ 100% on A3