



# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# Kamba<sup>®</sup> 500

Selective Herbicide

**ACTIVE CONSTITUENT: 500 g/L DICAMBA present as the dimethylamine salt**

**GROUP 1 HERBICIDE**

For the control of certain broadleaf weeds in winter cereals, pastures, conservation tillage, sugar cane, turf, rice and non-crop areas

**READ COMPLETE DIRECTIONS FOR USE BEFORE USING THIS PRODUCT.**

© Kamba is a registered trademark of Nufarm Australia Limited

APVMA Approval No.: 51080/51800

nufarm.com.au

## RESTRAINTS

DO NOT spray when rain seems likely to occur within 4 hours.

DO NOT spray when weeds are wet with dew or rain or under stress from drought, low soil fertility, extreme cold or water logging.

DO NOT spray outside recommended crop growth as crop damage may result.

DO NOT apply to crops or pastures with clover, lucerne or medics.

CROP	WEEDS CONTROLLED	STATE	RATE	CRITICAL COMMENTS
Wheat, Oats, Barley, Triticale, Commercial Rye From 5 leaf to early tillering (Z15-22)	Doublegee (Threecornered Jack, Spiny emex) Wireweed (Hogweed), Seedling docks, Tares, Hexham scent (Melilotus) Wild turnip ( <i>Brassica tournefortii</i> ), Soldier thistle, Mustards, Wild radish, Capeweed, Common ice-plant, Charlock, Ward's weed, Vetch, Turnip weed ( <i>Rapistrum rugosum</i> )	All States	160mL/ha plus 700mL/ha of MCPA 500	Spray when most weeds have germinated and are in the 2-4 leaf stage. In some instances suppression only of Wireweed (Hogweed) and Capeweed will occur.
Wheat, Oats, Barley, Triticale, Commercial Rye From 5 leaf early tillering stage until the fully tillered stage and before jointing occurs (Z15, 21-30)	Doublegee (Threecornered Jack, Spiny emex), Wireweed, Vetch, Variegated thistle, Tares, Sunflower, Hexham scent (Melilotus), Docks	SA only	280mL/ha	Spray when most weeds have germinated and are still in the young rosette stage. When weeds DO NOT form a rosette, spray when they have not more than 8 true leaves. In some instances suppression only of Wireweed (Hogweed) and Climbing buckwheat (Black bindweed) will occur.
	Doublegee (Threecornered Jack, Spiny emex), Climbing buckwheat (Black bindweed), Clover, Hexham scent (Melilotus), Docks	WA only		
	New Zealand spinach, Climbing buckwheat (Black bindweed), Doublegee (Threecornered Jack, Spiny emex), Wireweed (Hogweed), Hexham scent (Melilotus), Docks, Clover, Fat hen, Variegated thistle, Scotch (Cotton or Woolly) thistle, Sunflowers, Vetch, Tree hogweed	Qld, NSW, Vic only		
	Wild turnip ( <i>Brassica tournefortii</i> ), Radish, Mustards, Tares, Charlock, Capeweed, Sorrel, Saffron thistle, Chickweed, Clover, Common ice-plant, Hoary cress, Ward's weed, Turnip weed ( <i>Rapistrum rugosum</i> )	SA only	280mL/ha plus 1L/ha of MCPA 500 or 560mL/ha of Amicide <sup>®</sup> 625	
	Common ice-plant	WA only	320mL/ha plus 640mL/ha of Amicide <sup>®</sup> 625	
	Chickweed, Sorrel, Wireweed	WA only	400mL/ha or 280mL/ha plus 640mL/ha of Amicide <sup>®</sup> 625 or 280mL/ha plus 1.2L/ha of MCPA 500	DO NOT use Kamba <sup>®</sup> 500 plus Amicide <sup>®</sup> 625 in Oats in SA. If any of these weeds are found in these cereal crops as well as the above weeds, the addition of MCPA 500 OR Amicide <sup>®</sup> 625 is essential. Spray when most weeds have germinated and are still in the young rosette stage. In some instances, suppression only of Wireweed (Hogweed), Capeweed and Hoary cress will occur. A mixture with Amicide <sup>®</sup> 625 must not be used in Tasmania.

CROP	WEEDS CONTROLLED	STATE	RATE	CRITICAL COMMENTS
------	------------------	-------	------	-------------------

(continued)	Mintweed	NSW, Qld only	280mL/ha plus 1.2L/ha of MCPA	(continued)
	Wild turnip ( <i>Brassica tournefortii</i> ), Radish, Mustards, Tares, Charlock, Capeweed, Sorrel, Saffron thistle, Chickweed, Turnip weed ( <i>Rapistrum rugosum</i> )	Qld, Vic NSW, Tas only	500 or 280mL/ha plus 560mL/ha of Amicide® 625	
Wheat, Barley, Triticale, Commercial Rye From early tillering (4-5 fully emerged main stem leaves, plus one or more tillers) until fully tillered and before the start of jointing (Z21-30)	Climbing buckwheat (Black bindweed), Deadnettle, Turnip weed ( <i>Rapistrum rugosum</i> ), Wireweed, Wild turnip ( <i>Brassica tournefortii</i> )	NSW, Qld only	200-280mL/ha plus 5-7g/ha Ally*/Associate®	Spray when weeds have no more than 8 true leaves. Use higher rate on larger weeds. Tank mixes of Kamba® 500 and Ally*/Associate® may result in reduced control of Wireweed in some situations.
Barley, Triticale, Commercial Rye From 5 leaf until fully tillered and before jointing occurs (Z21-30)	Volunteer peas	SA only	60mL/ha plus 1L/ha Kamba® M	Apply when most peas are 5-7cm high.
Wheat, Oats From 5 leaf until fully tillered and before jointing occurs (Z15-30)			120mL/ha plus 1L/ha Kamba® M	
Rice Pre - Sowing Post - Sowing either before permanent water or until rice is at the mid tillering stage	Docks	NSW, NT only	400mL/ha (seedling docks) 560mL/ha (mature docks)	<b>Pre-sowing:</b> Apply before the end of August and DO NOT cultivate or graze for 14 days after spraying. <b>Post-sowing:</b> Preferably apply before the application of permanent water. If permanent water has been applied Kamba 500 can be applied up to the mid-tillering stage, provided the docks are exposed. Use 100L/ha for boom application.
Pinus Radiata Plantations At least 10 months prior to planting pines	Sorrel (in pastures to be planted to pines)	All States	1.1L/ha or 560mL/ha plus 1L/ha Amicide® 625	Apply in September/October period before the sorrel flowers. Add Nufarm Surfactant 600 or another non-ionic wetting agent to the mixture.
	Naturally regenerated pines	SA, Vic, NSW only	2-8L/ha	Use the high rate where total control is essential. Apply in Spring.
Sugarcane Post-emergence	Bathurst burr, Bellvine, Blackberry nightshade, Calopo, Caltrop, Cleome, Cobbler's pegs, Common sow thistle, Convolvulus, Fat hen, Green amaranth, Khaki weed, Noogoora burr, Parthenium weed, Perennial gooseberry, Pigweed, Pink burr, Redroot amaranth, Sensitive plant, Sida-retusa, Tar vine, Thickhead, Thornapple	NSW, Qld, WA, NT only	560mL/ha	Spray when most weeds have germinated and are in the young rosette stage or when they have not more than 8 true leaves. <b>Method of Application:</b> By boom spray as an overall directed spray, applying 100-250L of water per hectare or by aircraft applying 45L per hectare.
	Amaranthus, Black (Giant) pigweed, Caltrop, Mintweed, Noogoora burr, Pigweed, Thornapple ( <i>Datura</i> )	NSW, Qld, NT only	280mL/ha plus 1.8L/ha Nu-trazine Liquid or 320mL/ha plus 1.5L/ha Nu-trazine Liquid	Apply when weeds have 3-5 true leaves. DO NOT apply in hot dry conditions. Apply by boom spray as an overall spray applying 50-250L of water per hectare. Use the higher rate Kamba 500 where Caltrop is present.
Potatoes	Clover, Docks, Wireweed (Hogweed), Thistles, Chickweed, Climbing buckwheat (Black bindweed), Black nightshade, Fat hen	Tas only	500mL-1.2L/ha	Apply after haulm senescence in sufficient water to give even coverage. May be mixed with Amitrole T or Amicide® 625 to improve spectrum of weeds controlled. Refer to respective labels of these herbicides for application rates. Add suitable surfactant for improved coverage.  NOTE: DO NOT apply when potato haulms are still green.

SITUATION	WEEDS CONTROLLED	STATE	RATE			CRITICAL COMMENTS
			Per ha	High Vol Per 100L	Per 15L K'Sack	
Grass Pastures and Perennial Grass Seed Crops	Cotton (Woolly or True scotch) thistle, Variegated thistle, Star thistle, Black or Spear thistle, Illyrian thistle, Doublegee (Threecornered Jack, Spiny emex), Parthenium weed, Smart-weed (Water pepper), Hexham scent (Melilotus), Stemless thistle, Caltrop (Yellow vine), Horehound and Rumex spp, including Curled and Broadleaf docks	All States	280-560mL for seedlings	40mL for seedlings	10mL for seedlings	When applying by boom spray use 110-280L of mixture per hectare. Add Nufarm Surfactant 600 to spray mix.
	560mL-1.1L for young mature plants		76mL for young mature plants	18mL for young mature plants		
	Common ice-plant, Hoary cress, Sorrel and Saffron thistle		280mL plus 640mL of Amicide®625	20mL plus 50mL of Amicide®625	6mL plus 12mL of Amicide®625	
Non-Crop Situation	Annual ragweed, Blue heliotrope, Burr ragweed, Galvanised burr, Ground cherry, Ox-Eye daisy, Perennial ragweed, Russian knapweed, Skeleton weed, Wild (Field or Crow) garlic		8.8L	600mL	130mL	Use a minimum of 1500L/ha water carrier. Add Nufarm Surfactant 600. Spray Russian knapweed at flowering. Spray Angled onion weed, Bladder campion, Ox-Eye daisy and Paterson's curse before flowering. Spray Wild garlic during Spring before bulbs start to form and respray later if dormant bulbs shoot. Spray prior to flowering.  Blackberry nightshade at seedling stage. For Creeping knotweed apply in early Summer to actively growing plants. For Slender ice-plant and Khaki weed spray when actively growing at young rosette stage. For Rubber vine apply in Autumn during active growth.
	California (Perennial or Canada) thistle, Field bindweed, Paterson's curse, Prairie ground cherry, Ragwort, Stinking mayweed (Foetid Chamomile)		4L	280mL	60mL	
	Afghan thistle, Artichoke thistle, Star thistle		1.6L	100mL	24mL	
	Horehound, Angled onion weed (Threecornered garlic), Bladder campion, Cotton (Woolly or True scotch) thistle, Docks, Sorrel, Stemless thistle, Variegated (or spotted) thistle		1.2L	80mL	18mL	
	Illyrian thistle, Parthenium weed, Smart-weed		600mL	40mL	10mL	
	Poverty weed	SA only		2L	440mL	
	Mist flower	Qld, NT only	12L	800mL	180mL	
	Blackberry nightshade	WA only	320-560mL		5-9mL	
	Creeping knotweed		1.2-2.4L		18-36mL	
	Slender ice-plant		320mL+ 640mL Amicide®625		6mL	
	Khaki weed		600mL		9mL	
	Rubber vine		1.4L		22mL	
Grass Pastures and Non-Crop	Rubber vine	Qld, NT only	2-4L plus 2.5L of Estercide® 800	140-280mL plus 175mL of Estercide® 800	20-40mL plus 25mL of Estercide® 800	Apply during April or May. For application by misting machines, add 600mL Kamba 500 to 10L of water.
Turf	Catsear, Cudweeds, Pennyworth, Lamb's tongue, Fleabanes, Fumitory, Wireweed, Chickweed, Cotula, Sorrel, Dock, Clovers, Capeweed, Toad rush, Peppergrass, Dandelion, Medics, Scarlet pimpernel, Creeping Oxalis, Pearlwort, Trefoil, Yarrow, Bindy-Eye, Jo-Jo (Onehunga)	All States	1.2L plus 3.2L of Amicide®625	80mL plus 200mL of Amicide®625	18mL plus 50mL of Amicide®625	Use a minimum of 1000L/ha water carrier. DO NOT spray on Buffalo grass, Bent grass, Lippia or Strawberry clover. Spray when weeds are young and actively growing.
	Fat hen, Red flowered mallow, Purple - Top, Stagger weed, Swine cress	NSW, NT, SA, Qld, WA, Tas only				
Woody Species	Moreton Bay ash (Carbeen), Blue gum, Bitter bark, Brisbane black wattle	All States	By partial frill or tree injection technique mix 1 part Kamba 500 with 12.5 parts of water and apply 2mL of the mixture to cuts in the softwood at not more than 6cm spacings around the circumference of each tree.			
	Mimosa ( <i>Mimosa pigra</i> )	NT only	2-2.4L	400mL	60mL	Apply to actively growing plants. Preferably apply during wet season. Retreatment may be necessary to control seeding regrowth (for application see General Instructions).

**ADDITIONAL RESTRAINTS FOR CONSERVATION TILLAGE**

- When grass and broadleaf weeds are present, use a mixture of Kamba and Roundup® CT or Glyphosate CT.
- DO NOT disturb treated weeds by cultivation, sowing or grazing for 1 day after treatment of annual weeds and 7 days for perennial weeds.
- DO NOT treat weeds heavily covered with dust or silt.
- DO NOT apply if rainfall is likely within 6 hours of application.
- Observe plant back periods listed under General Instructions.
- When applying to dry soil surfaces, at least 15mm rainfall is required before the plant back period begins.

CROP	WEEDS CONTROLLED	STATE	RATE	CRITICAL COMMENTS
Seed Bed Salvage, Conservation tillage, Direct Drilling For weed control in fallow, stubble or pasture prior to sowing with a full disturbance implement	Amaranthus, (Red shank), Native amaranth, Sow thistle, Volunteer cereals, Turnip weed, Wireweed (Hogweed), Medics, Variegated thistle, Wild oats, Wild mustard, Docks, Paterson's curse	Qld, NSW, Vic, SA, WA, NT only	160-240mL/ha plus glyphosate 450g/L at 400- 600mL/ha or glyphosate 360g/L at 500-700mL/ha	If excessively wet conditions occur shortly after application or sowing some retardation of crop establishment may occur in certain broadleaf crops.  Spray when most weeds have germinated and are still in the young rosette or pre-tillering stage. Use the lower rate when weeds are prior to tillering or seedlings. Increase to the higher rate as weeds reach tillering or are 10-15cm diameter/high. In grazed situations if heavy grazing has occurred allow regrowth to 6-8cm high before spraying. Visible symptoms of weed control may not be apparent for 10-21 days.
	Barnyard grass, Liverseed grass	NSW only		
	Apple of Peru, Mintweed, Noogoora burr, Yellow vine (Caltrop), Climbing buckwheat (Black bindweed), Barnyard grass, Canary grass (Annual Phalaris), Thomapple, Wild or Prickly lettuce	Qld, NSW, NT only		
	Annual ryegrass, Volunteer field peas, Tree hogweed, Capeweed	NSW, Vic, SA, WA only		
	Barnyard grass, Liverseed grass	Qld, NT only	160-240mL/ha plus glyphosate 450g/L at 800mL-1.2L/ha	
	Sorrel	WA, SA, Vic, NSW only	160-240mL/ha plus 650-800mL/ha glyphosate 450g/L or 800mL-1L/ha glyphosate 360g/L	
Direct Drilling, conservation tillage, no-till fallow, minimum tillage, and all between cropping applications	Clover	All States	200mL/ha	Add to Roundup®/Glyphosate or Spray*Seed* as required (mixtures of Spray*Seed* plus Kamba 500 must not be used in Tas). Follow label recommended rates.
	Doublegee (Threecornered Jack, Spiny Emex), Wireweed, Vetch, Docks, Tares, Sunflower, Hexham scent (Melilotus), Variegated thistle	SA only	280mL/ha	Spray when most weeds have germinated and are still in the young rosette stage, when they have not more than 8 true leaves.
	Doublegee (Threecornered Jack, Spiny Emex), Climbing buckwheat (Black bindweed), Hexham scent (Melilotus), Docks	WA only		
	New Zealand spinach	Qld, NSW, NT only		
	Climbing buckwheat (Black bindweed), Doublegee (Threecornered Jack, Spiny Emex), Wireweed (Hogweed), Hexham scent (Melilotus), Docks, Fat hen, Variegated thistle, Scotch (Cotton or Woolly) thistle, Sunflowers, Vetch, Tree hogweed	Qld, NSW, Tas, Vic, NT only		
	Chickweed, Sorrel, Wireweed	WA only	400mL/ha	
	Common ice-plant, Hoary cress, Sorrel, Saffron thistle	All States	280mL/ha plus 640mL/ha Amicide® 625	
	Caltrop (Yellow vine), Datura, Bathurst and Noogoora burrs, Parthenium weed, Amaranthus, Mintweed, Annual gooseberry, Blackberry nightshade, Fat hen, Sunflower, Seedling perennial gooseberry, Seedling khaki weed, Bellvine, Cobbler's pegs, Star thistle, Black or Spear thistle, Illyrian thistle, Doublegee (Threecornered Jack, Spiny Emex), Smart-Weed (Water pepper), Stemless thistle, Horehound and Rumex spp including Curled and Broadleaf docks		320-560mL/ha for seedlings	Apply when weeds have at least 3-5 true leaves and are actively growing. DO NOT treat weeds which are beyond the rosette stage. Use higher rate on large weeds.
	Convolvulus, Sida-retusa, Sensitive Plant, Pink Burr, Cleome, Centro, Thickhead, Tar-Vine	Qld, NSW, NT, Vic, Tas, WA only		

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIOD**

**DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.**

**GENERAL INSTRUCTIONS**

Spray when weeds are young and actively growing, and unless stated otherwise, before the 3-5 true leaf stage.

**MIXING**

Add the required amount of the product directly to the spray tank and mix well.

**APPLICATION**

For Boom Application: Apply by a properly calibrated boom spray using not less than 50L of water/ha unless indicated otherwise in Critical Comments column.

For Aerial Application: Apply at least 25L total spray volume/ha. The aircraft should fly as low as practical under the prevailing conditions to minimise drift.

For High Volume Application: Apply minimum 1500L/ha.

For Knapsack Application: A 15L knapsack treats 150m<sup>2</sup>.

**MIMOSA CONTROL**

**Broadacre application** - Apply 2-2.4L/ha in 60L water/ha. Helicopter application preferred.

**High Volume application** - Apply 400mL in 110L water using ground equipment. Thoroughly wet all leaves and stems of the plant.

**Basal Bark application** - Apply 60mL in 15L water to plants of less than 5cm basal diameter. Spray liberally all around the base and 25-40cm up the stem.

**Cut Stump application** - Apply 60mL in 15L water on plants up to 10cm basal diameter. Cut trees as close to ground level as is possible and spray the freshly cut surface before the sap dries.

**Note:** A temporary wilting may be evident in crops after application. If seasonal conditions are late or dry seek further advice before spraying as damage may result from spraying crops under these conditions. Growers should seek advice before spraying recently released cereal varieties. Certain maize varieties may develop clubfoot of the prop roots.

**RESISTANT WEEDS WARNING**

<b>GROUP</b>	<b>I</b>	<b>HERBICIDE</b>
--------------	----------	------------------

Kamba 500 Herbicide is a member of the Benzoic acid group of herbicides. Kamba 500 has the Disruptor of plant cell growth mode of action. For weed resistance management Kamba 500 is a Group I Herbicide. Some naturally-occurring weed biotypes resistant to Kamba 500 and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Kamba 500 or other Group I Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from the failure of Kamba 500 to control resistant weeds.

**PLANT BACK PERIODS for Conservation Tillage**

Wait for the following periods before planting the following crops into soils sprayed with the product. If applied to a dry soil, at least 15mm of rainfall is required before the Plant Back Period begins.

**Note:** if water-logging occurs shortly after sowing, the product may cause some reduction in crop vigour. Use as directed only in the States indicated.

**NSW, VIC, SA, WA, TAS**

Rate/ha	PLANT BACK PERIOD (DAYS)									
	Wheat	Barley	Oats	Triticale	Rye	Lupins	Clover/ Medics	Mung beans	Sunflower	Safflower
200mL	1	1	1	1	1	7	7	5	1	14
280mL	7	7	7	7	7	14	14	5	7	21
560mL	14	14	14	14	14	21	21	10	14	28

**NSW, VIC, SA, WA, TAS**

Rate/ha	PLANT BACK PERIOD (DAYS)								
	Pigeon peas	Soybeans	Millet	Cotton	Sorghum	Maize	Rapeseed	Field peas	Chick peas
200mL	5	5	1	7	1	1	7	N.D.*	N.D.*
280mL	5	5	3	7	3	3	10	14	21
560mL	10	10	7	14	7	7	14	21	28

\* Not determined

**QLD, NT ONLY**

Rate/ha	CROP	PLANT BACK PERIOD
160-240mL	Wheat, Barley, Oats, Triticale, Commercial Rye, Sorghum, Maize, Millet, Sunflowers	1 day
	Soybeans, Mung beans, Pigeon peas	5 days
	Cotton, Rapeseed	7 days

**Note: For Qld, NT only:** For all crops and rates above 240mL/ha – Plant Back Period 21 days.

**Following Potato haulm destruction in Tas:** For rates and crops not included above, a Plant Back Period of 6 weeks applies generally and 8 weeks for seed crops.

**COMPATIBILITY**

Kamba 500 is compatible with Flowable Igran\*, Roundup® Max, Roundup® CT, Roundup®, Nufarm Glyphosate CT, Weedmaster® Duo, Credit®/Bonus®, Glean\*/Lusta®, Ally\*/Associate®, Amicide® 625, LV Estericide® 600, Estericide® 800, Diuron (900DF and Flowable), Ordram\* (Molinate), LVE MCPA. Also compatible with Nu-trazine (900DF and Flowable) and other Atrazines, but when used in combination with these products some temporary visible effects in the crop growth may be observed.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Spray equipment must be calibrated accurately prior to working on crops. Keep container sealed (airtight) when not in use. To prevent damage, DO NOT permit drift or vapour or spray mist to come into contact with fruit and ornamental trees, vines, vegetables, flowers, tobacco or any other field crop. Minimise spray drift by using low pressure and nozzles which DO NOT give a fine droplet size. DO NOT apply this

product on or near desirable trees or plants or in locations where the chemical may be washed or moved into contact with their roots. DO NOT use in high wind. All spray equipment must be thoroughly washed out after use and must not be used for spraying horticultural crops. DO NOT use at higher rate than recommended.

#### **PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT**

DO NOT use container for any other purpose. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

#### **STORAGE AND DISPOSAL**

##### **(15L)**

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse inner bladder before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean packaging to recycler or designated collection point. If not recycling, deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

##### **(2.5L, 5L, 10L, 20L, 200L)**

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, triple rinse, break, crush or puncture and deliver empty packaging for appropriate disposal at an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Empty containers and product must not be burnt.

##### **Envirodrum 110L Mini Bulk Returnable Container**

Store the original sealed Envirodrum in a cool well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Nufarm Australia Limited.

##### **Refillable containers (1000L)**

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

#### **SAFETY DIRECTIONS**

May irritate the eyes and skin. Avoid contact with eyes and skin. When opening container and preparing spray wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

#### **FIRST AID**

If poisoning occurs, contact a doctor, or Poisons Information Centre (Phone 13 11 26).

#### **MATERIAL SAFETY DATA SHEET**

For further information refer to the Material Safety Data Sheet (MSDS), which can be obtained from your supplier or from the Nufarm website – [www.nufarm.com.au](http://www.nufarm.com.au)

**In case of emergency: Phone 1800 033 498** Ask for shift supervisor. Toll free 24 hours.

#### **CONDITIONS OF SALE**

"Any provisions or rights under the Trade Practices Act 1974 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute."

Nufarm Australia Limited  
ACN 004 377 780  
103-105 Pipe Road  
Laverton North Victoria 3026  
Tel: (03) 9282 1000  
Fax: (03) 9282 1001

© Credit, Bonus, Amicide, Associate, Estercide and Weedmaster are registered trademarks of Nufarm Australia Limited.

© Roundup is a registered trademark of Monsanto Technology LLC used under licence by Nufarm Australia Limited.

\* Other registered trademarks