

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Ospray Dicamba 500 SL

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 500g/L DICAMBA
(present as the DIMETHYLAMINE SALT)

GROUP

HERBICIDE

For the control of certain broadleaf weeds in winter cereals, pastures, conservation tillage, sugar cane, turf, rice, grain sorghum and non-crop areas as specified in the Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING



0205wm



Ospray Pty Ltd ABN 23 110 199 169

50 Middleton Road

Cottonvale QLD 4375 AUSTRALIA

Phone: 1800 624 597

www.ospray.com.au

Part of the  CHEMINOVA Group

Contents 10L

Ospray Dicamba 500 SL

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point (contact *drumMuster* Collection agency or Local Council for collection sites). If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing the spray wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each days use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

MSDS

Additional information is listed in the material safety data sheet (MSDS). The MSDS is available from the supplier.

NOTICE:

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off label permits not endorsed by Ospray Pty Ltd or under abnormal conditions. Ospray Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

**THIS PRODUCT IS NOT CONSIDERED TO BE A DANGEROUS GOOD
UNDER THE AUSTRALIAN CODE FOR THE TRANSPORT OF
DANGEROUS GOODS BY ROAD AND RAIL**

**IN AN TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE**

APVMA Approval No. : 60348/10L/0708

Batch No.

DOM



POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Ospray Dicamba 500 SL

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 500g/L DICAMBA
(present as the DIMETHYLAMINE SALT)

GROUP

HERBICIDE

For the control of certain broadleaf weeds in winter cereals, pastures, conservation tillage, sugar cane, turf, rice, grain sorghum and non-crop areas as specified in the Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING



0205wm



Ospray Pty Ltd ABN 23 110 199 169

50 Middleton Road

Cottonvale QLD 4375 AUSTRALIA

Phone: 1800 624 597

www.ospray.com.au

Part of the  CHEMINOVA Group

Contents 10L

Ospray Dicamba 500 SL

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point (contact *drumMuster* Collection agency or Local Council for collection sites). If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing the spray wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each days use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

MSDS

Additional information is listed in the material safety data sheet (MSDS). The MSDS is available from the supplier.

NOTICE:

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off label permits not endorsed by Ospray Pty Ltd or under abnormal conditions. Ospray Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

**THIS PRODUCT IS NOT CONSIDERED TO BE A DANGEROUS GOOD
UNDER THE AUSTRALIAN CODE FOR THE TRANSPORT OF
DANGEROUS GOODS BY ROAD AND RAIL**

**IN AN TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE**

APVMA Approval No. : 60348/10L/0708

Batch No.

DOM



POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Ospray Dicamba 500 SL

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 500g/L DICAMBA
(present as the DIMETHYLAMINE SALT)

GROUP

HERBICIDE

For the control of certain broadleaf weeds in winter cereals, pastures, conservation tillage, sugar cane, turf, rice, grain sorghum and non-crop areas as specified in the Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING



0205wm



Ospray Pty Ltd ABN 23 110 199 169

50 Middleton Road

Cottonvale QLD 4375 AUSTRALIA

Phone: 1800 624 597

www.ospray.com.au

Part of the  CHEMINOVA Group

Contents 110L

Ospray Dicamba 500 SL

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point (contact *drumMuster* Collection agency or Local Council for collection sites). If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing the spray wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each days use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

MSDS

Additional information is listed in the material safety data sheet (MSDS). The MSDS is available from the supplier.

NOTICE:

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off label permits not endorsed by Ospray Pty Ltd or under abnormal conditions. Ospray Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

**THIS PRODUCT IS NOT CONSIDERED TO BE A DANGEROUS GOOD
UNDER THE AUSTRALIAN CODE FOR THE TRANSPORT OF
DANGEROUS GOODS BY ROAD AND RAIL**

**IN AN TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE**

APVMA Approval No. : 60348/110L/0708

Batch No.

DOM



POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Ospray Dicamba 500 SL

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 500g/L DICAMBA
(present as the DIMETHYLAMINE SALT)

GROUP I HERBICIDE

For the control of certain broadleaf weeds in winter cereals, pastures, conservation tillage, sugar cane, turf, rice, grain sorghum and non-crop areas as specified in the Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING



0205wm



Ospray Pty Ltd ABN 23 110 199 169

50 Middleton Road

Cottonvale QLD 4375 AUSTRALIA

Phone: 1800 624 597

www.ospray.com.au

Part of the  CHEMINOVA Group

Contents 1000L

Ospray Dicamba 500 SL

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point (contact *drumMuster* Collection agency or Local Council for collection sites). If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing the spray wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each days use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

MSDS

Additional information is listed in the material safety data sheet (MSDS). The MSDS is available from the supplier.

NOTICE:

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off label permits not endorsed by Ospray Pty Ltd or under abnormal conditions. Ospray Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

**THIS PRODUCT IS NOT CONSIDERED TO BE A DANGEROUS GOOD
UNDER THE AUSTRALIAN CODE FOR THE TRANSPORT OF
DANGEROUS GOODS BY ROAD AND RAIL**

**IN AN TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE**

APVMA Approval No. : 60348/1000L/0708

Batch No.

DOM



POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Ospray Dicamba 500 SL

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 500 g/L DICAMBA PRESENT AS THE DIMETHYLAMINE SALT

GROUP **I** HERBICIDE

For the control of certain broadleaf weeds in winter cereals, pastures, conservation tillage, sugar cane, turf, rice, grain sorghum and non-crop areas.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

APVMA Approval No. :
60348/0708



Ospray Pty Ltd ABN 23 110 199 169
50 Middleton Road
Cottonvale QLD 4375 AUSTRALIA
Phone: 1800 624 597
www.ospray.com.au
Part of the CHEMINOVA Group

0205wm

DIRECTIONS FOR USE:

Restraints: DO NOT spray when rain seems likely to occur within 4 hours.

DO NOT spray when weeds are wet with dew or rain or under stress from drought, low soil fertility, extreme cold or water logging.

DO NOT spray outside recommended crop growth stages as crop damage may result.

DO NOT apply to crops or pastures with clover, lucerne or medics.

Crop & Situation	Weeds	State	Rate/ha	Critical Comments
Wheat, Oats, Barley, Triticale, Commercial Rye From 5 leaf to early tillering (Zadoks 15 - 22)	Doublegee (Threecornered Jack, Spiny Emex) Wireweed (Hogweed), Seedling Docks, Tares, Hexham Scent (Melilotus) Wild Turnip (<i>Brassica toumefortii</i>), Soldier Thistle, Mustards, Wild Radish, Capeweed, Common Ice-Plant, Charlock, Ward's Weed, Vetch, Turnip Weed (<i>Rapistrum rugosum</i>)	All States	160mL plus 700mL of MCPA Amine 500g/L	Spray when most weeds have germinated and are in the 2 to 4 leaf stage. In some instances suppression only of Wireweed (Hogweed) and Capeweed will occur.
Wheat, Oats, Barley, Triticale, Commercial Rye From 5 leaf early tillering stage until the fully tillered stage and before jointing occurs (Zadoks 15, 21-30)	Doublegee (Threecornered Jack, Spiny Emex), Wireweed, Vetch, Variegated Thistle, Tares, Sunflower, Hexham Scent (Melilotus), Docks	SA only	280mL	Spray when most weeds have germinated and are still in the young rosette stage. When weeds do not form a rosette, spray when they have not more than 8 true leaves. In some instances suppression only of Wireweed (Hogweed) and Climbing Buckwheat (Black Bindweed) will occur.
	Doublegee (Threecornered Jack, Spiny Emex), Climbing Buckwheat (Black Bindweed), Clover, Hexham Scent (Melilotus), Docks	WA only		
	New Zealand Spinach, Climbing Buckwheat (Black Bindweed), Doublegee (Threecornered Jack, Spiny Emex), Wireweed (Hogweed), Hexham Scent (Melilotus), Docks, Clover, Fat-Hen, Variegated Thistle, Scotch (Cotton or Woolly) Thistle, Sunflowers, Vetch, Tree Hogweed	Qld, NSW, ACT, Vic only		
	Wild Turnip (<i>Brassica toumefortii</i>), Radish, Mustards, Tares, Charlock, Capeweed, Sorrel, Saffron Thistle, Chickweed, Clover, Common Ice-Plant, Hoary Cress, Ward's Weed, Turnip Weed (<i>Rapistrum rugosum</i>)	ACT, SA only	280mL plus 1L of MCPA Amine 500g/L or 700mL of 2,4-D Amine 500g/L	DO NOT use DICAMBA 500 SL plus 2,4-D in Oats in SA. If any of these weeds are found in these cereal crops as well as the above weeds, the addition of MCPA OR 2,4-D is essential. Spray when most weeds have germinated and are still in the young rosette stage. In some instances, suppression only of Wireweed (Hogweed), Capeweed and Hoary Cress will occur. DO NOT use DICAMBA 500 SL plus 2,4-D in Oats in SA.
	Common Ice-Plant	WA only	320mL plus 800mL of 2,4-D Amine 500g/L	
Chickweed, Sorrel, Wireweed		WA only	400mL or 280mL plus 800mL of 2,4-D Amine 500g/L or 280mL plus 1.2L of MCPA Amine 500g/L	If any of these weeds are found in these cereal crops as well as the above weeds, the addition of MCPA or 2,4-D is essential. Spray when most weeds have germinated and are still in the young rosette stage. In some instances, suppression only of Wireweed (Hogweed), Capeweed and Hoary Cress will occur. A mixture with 2,4-D Amine must not be used in Tasmania.
Mintweed		NSW, Qld, ACT only	280mL plus 1.2L of MCPA Amine 500g/L	
Wild Turnip (<i>Brassica toumefortii</i>), Radish, Mustards, Tares, Charlock, Capeweed, Sorrel, Saffron Thistle, Chickweed, Turnip Weed (<i>Rapistrum rugosum</i>)		Qld, NSW, Vic, Tas only	500g/L or 280mL plus 700mL of 2,4-D Amine 500g/L	
Wheat, Barley, Triticale, Commercial Rye From early tillering (4 to 5 fully emerged main stem leaves, plus one or more tillers) until fully tillered and before the start of jointing (Zadoks 21 - 30)	Climbing Buckwheat (Black Bindweed), Deadnettle, Turnip Weed (<i>Rapistrum rugosum</i>), Wireweed, Wild Turnip (<i>Brassica toumefortii</i>)	NSW, Qld, ACT only	200 - 280mL plus 5-7g Ally [®]	Spray when weeds have no more than 8 true leaves. Use higher rate on larger weeds. Tank mixes of DICAMBA 500 SL and Ally [®] may result in reduced control of Wireweed in some situations
Barley, Triticale, Commercial Rye From 5 leaf until fully tillered and before jointing occurs (Zadoks 21 - 30)	Volunteer Peas	SA only	60mL- plus 1L BANVEL M or KAMBA M	Apply when most Peas are 5-7cm high.
Wheat, Oats From 5 leaf until fully tillered and before jointing occurs (Zadoks 15 - 30)	Volunteer Peas	SA only	120mL plus BANVEL M or KAMBA M	Apply when most Peas are 5-7cm high.

Crop & Situation (cont)	Weeds	State	Rate/ha	Critical Comments
Grain Sorghum From 3 leaf stage until 6 leaf stage Maize Post emergence until maize is approx. 90cm high or until 15 days prior to tassle emergence	Caltrop (Yellow Vine), Bathurst and Noogoora Burrs, Parthenium Weed, Mintweed, Annual Gooseberry, Blackberry Nightshade, Fat-Hen, Sunflower, Seedling Perennial Gooseberry, Seedling Khaki Weed, Redshank, Dwarf Amaranth, Redroot Amaranth, Green Amaranth, Cobbler's Pegs, Field Bindweed, Thomapples (Datura), Bellvine, Lucerne, Annual Ground Cherry, Wireweed, Climbing Buckwheat (Black Bindweed), Pigweed, Common Sow Thistle, New Zealand Spinach, Doublegee (Threecornered Jack, Spiny Emex)	All States (NOT Central Qld)	320 – 560mL	Apply when weeds have at least 3 to 5 true leaves. Do not treat weeds which are beyond the rosette stage. Method of Application: By boom spray as an overall or directed spray, by applying 100 to 250 litres of water per hectare or by aircraft applying 45 litres per hectare. Use the higher rate on larger weeds and where Amaranthus, Mintweed and Blackberry Nightshade are the major problems or where Annual and Perennial Gooseberry are present Certain Maize varieties may develop clubfoot of the prop roots.
Grain Sorghum From 3 leaf stage until 6 leaf stage	Thornapple (Datura), Mintweed, Amaranthus, Black (Giant) Pigweed Caltrop (Yellow Vine), Bladder Ketmia, Noogoora Burr	NSW, ACT, Qld, NT only	280mL plus 1.8L Atrazine 500g/L or 320mL plus 1.5L Atrazine 500g/L	Apply when weeds have 3 to 5 true leaves. Do not apply in hot dry conditions. Apply by boom spray as an overall spray applying 50 to 250 litres of water per hectare. Use the higher rate of DICAMBA 500 SL where Caltrop or Bladder Ketmia are present.
Rice Pre - Sowing Post - Sowing either before permanent water or until rice is at the mid-tillering stage	Docks	NSW, NT, only	400mL (seedling docks) 560mL (mature docks)	Pre-sowing: Apply before the end of August and do not cultivate or graze for 14 days after planting. Post-sowing: Preferably apply before the application of permanent water. If permanent water has been applied DICAMBA 500 SL can be applied up to the mid-tillering stage, provided the docks are exposed. Use 100L/ha for boom application.
<i>Pinus Radiata</i> Plantations At least 10 months prior to planting pines	Sorrel (in pastures to be planted to pines)	All States	1.1L or 560mL plus 1.4L of 2,4-D Amine 500g/L	Apply in September/October period before the sorrel flowers. Add Wetter 1000 or another non-ionic wetting agent to the mixture.
	Naturally regenerated pines	SA, Vic, NSW, ACT only	2 – 8L	Use the high rate where total control is essential. Apply in Spring.

DIRECTIONS FOR USE (CONTINUED):

Crop	Weeds Controlled	State	Rates/ha	Critical Comments
Sugarcane Post-emergence	Bathurst Burr, Bellvine, Blackberry Nightshade, Calopo, Caltrop, Cleorne, Cobbler's Pegs, Common Sow Thistle, Convolvulus, Fat-Hen, Green Amaranth, Khaki Weed, Noogoora Burr, Parthenium Weed, Perennial Gooseberry, Pigweed, Pink Burr, Redroot Amaranth, Sensitive Plant, Sida-retusa, Tar Vine, Thickhead, Thornapple	NSW, Qld, WA, NT, only	560mL	Spray when most weeds have germinated and are in the young rosette stage or when they have not more than 8 true leaves. Method of Application: By boom spray as an overall directed spray, applying 100 to 250 litres of water per hectare or by aircraft applying 45 litres per hectare.
	Amaranthus, Black (Giant) Pigweed, Caltrop, Mintweed, Noogoora Burr, Pigweed, Thomapple (Datura)	NSW, Qld, NT only	280mL plus 1.8L Atrazine 500g/L or 320mL plus 1.5L Atrazine 500g/L	Apply when weeds have 3 to 5 true leaves. Do not apply in hot dry conditions. Apply by boom spray as an overall spray applying 50 to 250 litres of water per hectare. Use the higher rate DICAMBA 500 SL where Caltrop is present.
Potatoes	Clover, Docks, Wireweed (Hogweed), Thistles, Chickweed, Climbing Buckwheat (Black Bindweed), Blackberry Nightshade, Fat-Hen	Tas only	500mL – 1.2L	Apply after haulm senescence in sufficient water to give even coverage. May be mixed with Amitrole or 2,4-D to improve spectrum of weeds controlled. Refer to respective labels of these herbicides for application rates. Add suitable surfactant for improved coverage. NOTE: Do not apply when potato haulms are still green.

DIRECTIONS FOR USE: (CONTINUED):

Situation	Weeds Controlled	State	Rate			Critical Comments	
			Per ha	High Vol per 100L	Per 15L Knapsack		
Grass Pastures and Perennial Grass Seed Crops	Cotton (Woolly or True Scotch) Thistle, Variegated Thistle, Star Thistle, Black or Spear Thistle, Illyrian Thistle, Doublegee (Threecornered Jack, Spiny Emex), Parthenium Weed, Smart-Weed (Water Pepper), Hexham Scent (Melilotus), Stemless Thistle, Caltrop (Yellow Vine), Horehound and <i>Rumex</i> spp, including Curled and Broadleaf Docks	All States	280-560mL for seedlings	40mL for seedlings	10mL for seedlings	When applying by boom spray use 110 to 280 litres of mixture per hectare.	
			560mL-1.1L for young mature plants	76mL for young mature plants	18mL for young mature plants		
	Common Ice-Plant, Hoary Cress, Sorrel and Saffron Thistle	All States	280mL plus 800mL of 2,4-D Amine 500g/L	20mL plus 60mL of 2,4-D Amine 500g/L	6mL plus 15mL of 2,4-D Amine 500g/L	Add 600g/L non-ionic surfactant to spray mix.	
Non-Crop Situation	Annual Ragweed, Blue Heliotrope, Burr Ragweed, Galvanised Burr, Ground Cherry, Ox-Eye Daisy, Perennial Ragweed, Russian Knapweed, Skeleton Weed, Wild (Field or Crow) Garlic	All States	8.8L	600mL	130mL	Use a minimum of 1500 L/ha water carrier. Add Wetter 1000. Spray Russian Knapweed at flowering. Spray Angled Onion Weed, Bladder Campion, Ox-Eye Daisy and Paterson's Curse before flowering. Spray Wild Garlic during Spring before bulbs start to form and re spray later if dormant bulbs shoot. Spray prior to flowering.	
			4L	280mL	60mL		
			1.6L	100mL	24mL		
			1.2L	80mL	18mL		
	Illyrian Thistle, Parthenium Weed, Smart-Weed			600mL	40mL	10mL	
	Poverty Weed	SA only	-	2L	440mL		
	Mist Flower	Qld, NT only	12L	800mL	180mL		
	Blackberry Nightshade	WA only	320-560mL	-	5 - 9mL	Blackberry Nightshade at seedling stage.	
	Creeping Knotweed	WA only	1.2-2.4L	-	18 - 36mL	For Creeping Knotweed apply in early summer to actively growing plants. For Slender Ice-Plant and Khaki Weed spray when actively growing at young rosette stage. For Rubber Vine apply in autumn during active growth.	
	Slender Ice-Plant	WA only	320mL + 800mL 2,4-D 500g/L	-	6mL		
Khaki Weed	WA only	600mL	-	9mL			
Rubber Vine	WA only	1.4L	-	22mL			

Grass Pastures and Non-Crop	Rubber Vine	Qld, NT only	2-4L plus 2.5L 2,4-D Ester 800g/L	140-280mL plus 175mL of 2,4-D Ester 800g/L	20 - 40mL plus 25mL of 2,4-D Ester 800g/L	Apply during April or May. For application by misting machines, add 600mL DICAMBA 500 SL to 10L of water.
Turf	Catsear, Cudweeds, Pennywort, Lamb's Tongue, Fleabanes, Fumitory, Wireweed, Chickweed, Cotula, Sorrel, Dock, Clovers, Capeweed, Toad Rush, Peppergrass, Dandelion, Medics, Scarlet Pimpernel, Creeping Oxalis, Pearlwort, Trefoil, Yarrow, Bindy-Eye, Jo-Jo (Onehunga)	All States	1.2L plus 4L of 2,4-D Amine 500g/L	80mL - plus 250mL of 2,4-D Amine 500g/L	18mL plus 60mL of 2,4-D Amine 500g/L	Use a minimum of 1000L/ha water carrier. Do not spray on Buffalo Grass, Bent Grass, Lippia or Strawberry Clover. Spray when weeds are young and actively growing.
	Fat-Hen, Red Flowered Mallow, Purple - Top, Stagger Weed, Swine Cress	NSW, ACT, NT, SA, Qld, WA, Tas only				
Woody Species	Moreton Bay Ash (Carbeen), Blue Gum, Bitter Bark, Brisbane Black Wattle	All States	By partial frill or tree injection technique mix 1 part DICAMBA 500 SL with 12.5 parts of water and apply 2mL of the mixture to cuts in the softwood at not more than 6cm spacings around the circumference of each tree.			
	Mimosa (<i>Mimosa pigra</i>)	NT only	2-2.4L	400mL	60mL	Apply to actively growing plants. Preferably apply during wet season. Retreatment may be necessary to control seeding regrowth (for application see General Instructions).

ADDITIONAL RESTRAINTS FOR CONSERVATION TILLAGE

- When grass and broadleaf weeds are present, use a mixture of DICAMBA 500 SL and Roundup CT or Glyphosate 450.

- DO NOT disturb treated weeds by cultivation, sowing or grazing for 1 day after treatment of annual weeds and 7 days for perennial weeds.

- DO NOT treat weeds heavily covered with dust or silt.

- DO NOT apply if rainfall is likely within 6 hours of application.

- Observe plant back periods listed under General Instructions.

- When applying to dry soil surfaces, at least 15mm rainfall is required before the plant back period begins.

Crop	Weeds Controlled	State	Rates/ha	Critical Comments
Seed Bed Salvage, Conservation tillage, Direct Drilling <p>For weed control in fallow, stubble or pasture prior to sowing with a full disturbance implement</p>	Amaranthus (Red Shank), Native Amaranth, Sow Thistle, Volunteer Cereals, Turnip Weed, Wireweed (Hogweed), Medics, Variegated Thistle, Wild Oats, Wild Mustard, Docks, Paterson's Curse	Qld, NSW, ACT, Vic, SA, WA, NT only	160-240mL plus glyphosate 450g/L at 400-600mL or glyphosate 360g/L at 500-700mL	If excessively wet conditions occur shortly after application or sowing some retardation of crop establishment may occur in certain broadleaf crops. <p>Spray when most weeds have germinated and are still in the young rosette or pre- tillering stage. Use the lower rate when weeds are prior to tillering or seedlings.</p>
	Barnyard Grass, Liverseed Grass	NSW, ACT only		
	Apple of Peru, Mintweed, Noogoora Burr, Yellow Vine (Caltrop), Climbing Buckwheat (Black Bindweed), Barnyard Grass, Canary Grass (Annual Phalaris), Thorn Apple, Wild or Prickly Lettuce	Qld, NSW, ACT, NT only		Increase to the higher rate as weeds reach tillering or are 10-15cm diameter/high. In grazed situations if heavy grazing has occurred allow re- growth to 6-8cm high before spraying.
	Annual Ryegrass, Volunteer Field Peas, Tree Hogweed, Capeweed	NSW, ACT, Vic, SA, WA only		Visible symptoms of weed control may not be apparent for 10-21 days.
	Barnyard Grass, Liverseed Grass	Qld, NT only	160-240mL plus glyphosate 450g/L at 800mL-1.2L	
	Sorrel	WA, SA, Vic, NSW, ACT only	160-240mL plus 650-800mL glyphosate 450g/L or 800mL – 1L glyphosate 360g/L	
Direct Drilling, conservation tillage, non-till fallow, minimum tillage, and all between cropping applications	Clover	All States	200mL	Add to Glyphosate or Spray.Seed as required (mixtures of Spray.Seed plus DICAMBA 500 SL must not be used in Tas). Follow label recommended rates.
	Doublegee (Threecornered Jack, Spiny Emex), Wireweed, Vetch, Docks, Tares, Sunflower, Hexham Scent (Melilotus), Variegated Thistle	SA only	280mL	Spray when most weeds have germinated and are still in the young rosette stage, when they have not more than 8 true leaves.
	Doublegee (Threecornered Jack, Spiny Emex), Climbing Buckwheat (Black Bindweed), Hexham Scent (Melilotus), Docks	WA only		
	New Zealand Spinach	Qld, NSW, ACT, NT only		
	Climbing Buckwheat (Black Bindweed), Doublegee (Threecornered Jack, Spiny Emex), Wireweed (Hogweed), Hexham Scent (Melilotus), Docks, Fat-Hen, Variegated Thistle, Scotch (Cotton or Woolly) Thistle, Sunflowers, Vetch, Tree Hogweed	Qld, NSW, ACT, Tas, Vic, NT only		
	Chickweed, Sorrel, Wireweed	WA only	400mL	
	Common Ice-Plant, Hoary Cress, Sorrel, Saffron Thistle	All States	280mL plus 800mL of 2,4-D Amine 500g/L	
	Caltrop (Yellow Vine), Datura, Bathurst and Noogoora Burrs, Parthenium Weed, Amaranthus, Mintweed, Annual Gooseberry, Blackberry Nightshade, Fat-Hen, Sunflower, Seedling Perennial Gooseberry, Seedling Khaki Weed, Bellvine, Cobbler's Pegs, Star Thistle, Black or Spear Thistle, Illyrian Thistle, Doublegee (Threecornered Jack, Spiny Emex), Smart-Weed (Water Pepper), Stemless Thistle, Horehound and Rumex spp including Curled and Broadleaf Docks	All States	320-560mL for seedlings	Apply when weeds have at least 3 to 5 true leaves and are actively growing. Do not treat weeds which are beyond the rosette stage. Use higher rate on large weeds.
	Convolvulus, Sida-retusa, Sensitive Plant, Pink Burr, Cleome, Centro, Thickhead, Tar-Vine	Qld, NSW, ACT, NT, Vic, Tas, WA only		

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION. WITHHOLDING PERIOD: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS RESISTANT WEEDS WARNING

GROUP I HERBICIDE

OSPRAY DICAMBA 500 SL SL Selective Herbicide is a member of the Benzoic acid group of herbicides. DICAMBA 500 SL SL has the Disruptor of plant cell growth mode of action. For weed resistance management DICAMBA 500 SL is a Group I Herbicide. Some naturally occurring weed biotypes resistant to DICAMBA 500 SL and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by DICAMBA 500 SL or any other Group I Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, OSPRAY Pty. Ltd. accepts no liability for any losses that may result from the failure of DICAMBA 500 SL to control resistant weeds. Spray when weeds are young and actively growing, and unless stated otherwise, before the 3-5 leaf stage.

MIXING

Add the required amount of the product directly to the tank and mix well.

APPLICATION

For Boom Application: Apply by a properly calibrated boom spray using not less than 50 litres of water per hectare unless indicated otherwise in the Critical Comments column.

PLANT BACK PERIODS FOR CONSERVATION TILLAGE

Wait for the following periods before planting the following crops into soils sprayed with the product. If applied to a dry soil, at least 15 mm of rainfall is required before the Plant Back Period begins.

Note: If water-logging occurs shortly after sowing, the product may cause some reduction in crop vigour. Use as directed only in the States indicated.

Rate per Hectare	PLANT BACK PERIOD (DAYS)				
	Wheat	Barley	Oats	Triticale	Rye
200mL	1	1	1	1	1
280mL	7	7	7	7	7
560mL	14	14	14	14	14

Rate per Hectare	PLANT BACK PERIOD (DAYS)				
	Lupins	Clover/Medics	Mung Beans	Sunflower	Safflower
200mL	7	7	5	1	14
280mL	14	14	5	7	21
560mL	21	21	10	14	28

Rate per hectare	PLANT BACK PERIOD (DAYS)			
	Pigeon Peas	Soybeans	Millet	Cotton
200mL	5	5	1	7
280mL	5	5	3	7
560mL	10	10	7	14

Rate per Hectare	PLANT BACK PERIOD (DAYS)				
	Sorghum	Maize	Canola	Field Peas	Chick Peas
200mL	1	1	7	N.D.*	N.D.*
280mL	3	3	10	14	21
560mL	7	7	14	21	28

- Not determined

Rate	CROP	PLANT BACK PERIOD
160-240mL/ha	Wheat, Barley, Oats, Triticale, Commercial Rye, Sorghum, Maize, Millet, Sunflowers	1 day
	Soybeans, Mung Beans, Pigeon Peas	5 days
	Cotton, Canola	7 days

NOTE: FOR QLD, NT ONLY. FOR ALL CROPS AND RATES ABOVE 600mL/HA - PLANT BACK PERIOD 21 DAYS. FOLLOWING POTATO HAULM DESTRUCTION IN TAS: FOR RATES AND CROPS NOT INCLUDED ABOVE, A PLANT BACK PERIOD OF 6 WEEKS APPLIES GENERALLY AND 8 WEEKS FOR SEED CROPS.

COMPATIBILITY

DICAMBA 500 SL is compatible with Flowable Salvation†, Roundup CT, Roundup, Glyphosate, chloresulfuron, Ally, 2,4-D Amine, 2,4-D Ester and Diuron. Also compatible with Atrazine, but when used in combination with these products some temporary visible effects in the crop growth may be observed.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Spray equipment must be calibrated accurately prior to working on crops. Keep container sealed (airtight) when not in use. To prevent damage, DO NOT permit drift or vapour or spray mist to come into contact with fruit and ornamental trees, vines, vegetables, flowers, tobacco or any other field crop. Minimise spray drift by using low pressure and nozzles which DO NOT give a fine droplet size. DO NOT apply this product on or near desirable trees or plants or in locations where the chemical may be washed or moved into contact with their roots. DO NOT use in high wind. All spray equipment must be thoroughly washed out after use and must not be used for spraying horticultural crops. DO NOT use at higher rate then recommended.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT use container for any other purpose. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point (contact *drumMuster* Collection agency or Local Council for collection sites). If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury the empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

For Aerial Application: Apply at least 25 litres total spray volume per hectare. The aircraft should fly as low as practical under the prevailing conditions to minimise drift.

For High Volume Application: Apply a minimum 1500 litres per hectare.

For Knapsack Application: A 15 litre knapsack treats 150 square metres.

MIMOSA CONTROL

Broadacre application: Apply 2-2.4 L/ha water in 60 litres of water per hectare. Helicopter application preferred.

High Volume application: Apply 400 mL in 110 L water using ground equipment. Thoroughly wet all leaves and stems of the plant.

Basal Bark application: Apply 60 mL in 15 L water to plants of less than 5 cm basal diameter. Spray liberally all around the base and 25-40 cm up the stem.

Cut Stump application: Apply 60 mL in 15 L water on plants up to 10 cm basal diameter. Cut trees as close to ground level as is possible and spray the freshly cut surface before the sap dries.

Note: A temporary wilting may be evident in crops after application. If seasonal conditions are late or dry seek further advice before spraying as damage may result from spraying crops under these conditions. Growers should seek advice before spraying recently released cereal varieties. Certain maize varieties may develop clubfoot of the prop roots.