

POISON
KEEP OUT OF REACH OF CHILDREN

Casoron[®] **G**
HERBICIDE

ACTIVE CONSTITUENT: 67.5 g/kg DICHLOBENIL

GROUP 0 HERBICIDE

For pre-emergent weed control in orchards and vineyards; established shrubs and trees in flower beds and lawns in gardens and nursery stock in pots or field; blackcurrants, raspberries and gooseberries and for weed control around commercial, industrial and aquatic areas, as per DIRECTIONS FOR USE table

Net Contents: 25 kg

Chemtura Australia Pty Ltd
Level 7, 435 King William Street Adelaide SA 5000
Tel: (08) 8112 0900 Fax: (08) 8112 0999

 **Chemtura**
AGROSOLUTIONS™

DIRECTIONS FOR USE:**RESTRAINTS:**

- DO NOT apply within 4 weeks of transplanting.
- DO NOT graze livestock on treated areas.
- DO NOT use in glasshouses or enclosed shadehouses.
- DO NOT apply to annual plants.
- DO NOT apply to water which will be used for crop irrigation, for livestock watering or for human consumption.
- DO NOT use on commercial fish or shellfish waters.

Situation	Weeds Controlled	State	Rate	Critical Comments
Established shrubs and trees in beds and lawns, including: Acacia, Banksia, Birch, Box, Cedar, Cypress, Eucalyptus, Gardenia, Hakea, Hawthorn, Hazel, Heath, Holly, Lilly Pilly, Magnolia, Pines, Poplar, Rhododendron, Roses, Tea Tree, Willow.	Annual Grasses and Broadleaf weeds	All States	600 g / 100m ²	Spread the granules evenly over the soil of the area to be treated. Remove existing weeds by hand, desiccation or cultivation. Use higher rate on heavier soils.
			60 – 90 kg per treated hectare (2m x 5 km)	
Orchards, including: Apples, Apricots, Citrus, Peaches, Plums, and Vineyards		Tas. Only		Apply late winter to early spring before growth has started.
Blackcurrants, Raspberries, Gooseberries.		All States	60 g / 10 m ²	Stock not to be repotted.
			30 g / 10 m ²	New stock recently potted out. Apply two applications of the granules at 6-8 week intervals. Spread granules evenly over the soil of the area to be treated. Remove existing weeds by hand, desiccation or cultivation. Apply only to plants at least 6 months old.
Nursery stock in pots or field	Annual Weeds	All States	180 – 250 g per 10 m ²	Treatment prior to germination of the weeds will provide long lasting effect. Satisfactory result will be obtained if large weeds are chipped out, if treatment of seedling weeds is necessary. Use higher rate for heavier weed infestations.
			Perennial Weeds	250 – 370 g per 10 m ²
Commercial & Industrial Areas	Aquatic Weeds including Ribbonweed, Pondweeds, Milfoils (Cattail), Chara	All States	1.7 – 2.3 kg per 100 m ²	Exposed soil. Best results obtained when application is made to moist soil situation as close as is practicable to the time when these areas will be flooded. Use higher rate for heavier weed infestations.
			170 – 230 kg per hectare treated area OR 1.7 – 2.3 kg per 100 m ²	For water less than one metre deep. Apply to standing (ie. not moving) water and toward the end of the dormant period of the plant. Preferably July – August. Use higher rate for heavier weed infestations.
			230 – 340 kg per hectare treated area OR 2.3 – 3.4 kg per 100 m ²	For water greater than one metre deep. Apply to standing water (ie. not moving) and toward the end of the dormant period of the plant. Preferably July – August. Use higher rate for heavier weed infestations.

GENERAL INSTRUCTIONS

- DO NOT apply Casoron G Herbicide during periods of intense rainfall, when wind speed exceeds 30 kph for ground application or under conditions which cause granule movement to non-target areas.
- On slopes of 20° to 30°, apply Casoron G Herbicide by broadcast, strip or spot application only. On slopes greater than 30°, apply by spot application method only.
- DO NOT dispense via dust prone methods. Only use with applicators specifically designed to dispense granular products.

Resistant Weeds Warning**GROUP O HERBICIDE**

Casoron G Herbicide is a member of the Nitrile group of herbicides. This product has diverse sites for mode of action. For weed resistance management, the product is a Group O herbicide. Some naturally-occurring weed biotypes resistant to the product and other Group O herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group O herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Chemtura Australia Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

Application**Horticultural Situations:**

This product provides effective season-long control of many annual grasses and broadleaf weeds in a variety of horticultural situations. Application of the granules to a weed free surface ensures best results by preventing growth before it begins. Young, germinating weeds are killed. Take care to avoid letting granules lodge against buds.

Commercial, Industrial and Aquatic Situations:

Note: Consult your State Fisheries Department before applying this product to public waters.

This product is an easy to use form of highly active chemical. Simply sprinkle the granules, at the recommended rates evenly over the area to be treated. This product is suitable for the use on driveways, paths and tennis courts where desirable trees are in close proximity.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from equipment, that may cause product to be placed or moved onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF LIVESTOCK

DO NOT graze livestock on treated areas.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate dams, rivers or streams with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well ventilated area, out of direct sunlight. Do not dispose of undiluted chemical on-site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

FIRST AID

If poisoning occurs, contact a doctor or Poison Information Centre.
Phone Australia 131 126.

MATERIAL SAFETY DATA SHEET

For additional information refer to the Material Safety Data Sheet.

CONDITIONS OF SALE

Chemtura Australia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Chemtura Australia Pty Ltd is limited to the replacement of the goods or (at the option of Chemtura Australia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Chemtura Australia Pty Ltd within thirty days of delivery.

In an Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS - AUSTRALIA WIDE
---	--

© Casoron is a registered trademark of a subsidiary of Chemtura Corporation.



BN
DOM

APVMA Approval No. 54240/25/0107

4001778-0109

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

**WITHHOLDING PERIOD:
NOT REQUIRED WHEN USED AS DIRECTED.**