

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**Farmoz**

# Miti-Fol EC

## MITICIDE

ACTIVE CONSTITUENT: 240 g/L DICOFOL

SOLVENT: 587g/L XYLENE

GROUP **UN** INSECTICIDE

*For the control of mites on fruit and vegetable crops and ornamentals as specified in the Directions for Use Table.*

[www.farmoz.com.au](http://www.farmoz.com.au)

### DIRECTIONS FOR USE:

DO NOT use on crops grown in glasshouses in Tasmania.

CROP/SITUATION	PEST/DISEASE	STATE	RATE/100 L	CRITICAL COMMENTS
Babacos	Broad Mite, Two Spotted Mite	Qld only	200 mL	Apply at first appearance of mites (as shown by russetting along the veins in bananas) and repeat as necessary.
Bananas	Two Spotted Mite		250 mL	
	Strawberry Spider Mite		200 mL	
Citrus	Broad Mite, Brown Citrus Rust Mite, Citrus Rust Mite, Citrus Bud Mite, Two Spotted Mite			
Cotton	Spider Mites	NSW only	4 L/ha	Apply by ground rig at first appearance of mites, before row closure. For best results use a minimum spray volume of 160 L/sprayed ha and use 3-5 hollow cone nozzles per row. Multiple applications of this product could result in development of resistance to dicofol.
Grapevines	Bunch Mite, Grapeleaf Blister Mite	NSW, Vic, Tas, SA, WA only	140 mL	Apply just after bloom and repeat at first appearance of mites.
Hops	Two Spotted Mite	Tas only	200 mL	Apply at first appearance of mites and repeat as necessary.
Mangoes	Mango Bud Mite, Tea Red Spider Mite	Qld only		
Olives	Two Spotted Mite		250 mL	

CROP/SITUATION	PEST/DISEASE	STATE	RATE/100 L	CRITICAL COMMENTS
Ornamentals	Broad Mite, Cyclamen Mite, False Spider Mite	Qld, Tas, SA, WA, NT only	200 mL	Apply at first appearance of mites (as shown by distorted and twisted leaves, especially near the growing points) and repeat as necessary. NOTE: DO NOT use when control with predatory mites is practiced. NOTE: May damage some house plants, chrysanthemums, gardenias, hibiscus, roses and some varieties of fuchsia especially in hot weather.
	Tomato Russet Mite, Two Spotted Mite	All States		
Paw Paws	Broad Mite	Qld only		Apply at first appearance of mites (as shown by twisted new leaves or scarring on young leaves or fruit) and repeat 2 weeks later.
Pome fruit	Bryobia Mite, Two Spotted Mite, European Red Mite	All States		Apply at first appearance of mites and repeat as necessary.  NOTE: May cause russeting in pears or ring spot under some climatic conditions.
Stone fruit	Bryobia Mite, Two Spotted Mite, European Red Mite			
Strawberries	Two Spotted Mite			
Tobacco	Two Spotted Mite	Qld, NSW, Vic only	150 mL	
Tomatoes	Tomato Russet Mite	All States	200 mL	NOTE: DO NOT use on eggplants (aubergines).
Vegetables	Two Spotted Mite, Tomato Russet Mite			

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD:**

**BABACOS, BANANAS, CITRUS, COTTON, HOPS, MANGOES, OLIVES, PAW PAWS, POME FRUIT, STONE FRUIT, STRAWBERRIES, TOMATOES, VEGETABLES: DO NOT APPLY LATER THAN 7 DAYS BEFORE HARVEST.**  
**GRAPES: DO NOT APPLY LATER THAN 28 DAYS BEFORE HARVEST.**  
**DO NOT GRAZE OR CUT FOR STOCK FOOD.**

**GENERAL INSTRUCTIONS**

DO NOT use on crops grown in glass houses in Tasmania.

**INSECTICIDE RESISTANCE WARNING**

For insecticide resistance management FARMOZ MITI-FOL EC Miticide is a Group UN insecticide.

**GROUP UN INSECTICIDE**

Some naturally occurring insect biotypes resistant to MITI-FOL EC and other Group UN insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if MITI-FOL EC and other Group UN insecticides are used repeatedly. The effectiveness of MITI-FOL EC on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, FARMOZ Pty. Ltd. accepts no liability for any losses that may result from the failure of MITI-FOL EC to control resistant insects. MITI-FOL EC may be subject to specific resistance management strategies. For further information contact your local supplier, FARMOZ representative, or local agricultural department agronomist.

**MIXING**

Mix the required quantity of product with the water in the spray vat, while stirring or with agitators running.

**APPLICATION**

Thorough coverage, including the underside of leaves, is essential.

**COMPATIBILITY**

This product may be combined with any ONE of the following products: Copper Oxychloride, Maldison, Mancozeb, Thiram, Ziram. DO NOT mix with Bordeaux mixture. As formulations of other manufacturers' products are beyond the control of FARMOZ Pty. Ltd., all mixtures should be tested prior to mixing commercial quantities.

**PROTECTION OF LIVESTOCK**

Low hazard to bees. May be applied on any plants at any time. Low hazard to scale parasites, but will reduce predatory mite populations.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

**STORAGE AND DISPOSAL**

Store in the closed original container in a dry, cool, well-ventilated area, out of direct sunlight. The product is stable for a minimum two (2) years if stored as indicated above. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS**

Poisonous if inhaled or swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves and face shield or goggles. If product splashes on skin, immediately wash area with soap and water. If splashes in eyes, wash it out immediately with water. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre on 131126. If swallowed, DO NOT induce vomiting. Give a glass of water.

**MSDS**

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for MITI-FOL EC is available from FARMOZ on request. Call Customer Service on (02) 9431 7800.

**CONDITIONS OF SALE:** The use of FARMOZ MITI-FOL EC Miticide being beyond the control of the manufacturer, no warranty expressed or implied is given by FARMOZ Pty. Ltd. regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and FARMOZ Pty. Ltd. accepts no responsibility for any consequence whatsoever resulting from the use of this product.

**UN No: 1993 FLAMMABLE LIQUID N.O.S.\***  
**PACKAGING GROUP: III HAZCHEM CODE: 2[Y]**

**PACK SIZE: 5 L, 20 L**

