

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KELTHANE[®] MF MITICIDE

ACTIVE CONSTITUENT: 480 g/L DICOFOL

GROUP	2B	INSECTICIDE
--------------	-----------	--------------------

For the control of mites on certain crops and ornamentals
as specified in the directions for use.

IMPORTANT: READ THE ATTACHED LEAFLET
BEFORE USING THIS PRODUCT

CONTENTS: 5 & 20 LITRES

Crop Care Australasia Pty Ltd, 77 Tingira Street Pinkenba Qld 4008

STORAGE AND DISPOSAL

DO NOT STORE NEAR NAKED FLAMES. Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse empty containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of water ways desirable vegetation and tree roots. Empty containers and product should not be burnt

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing the spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

Additional information is listed in the Material Safety Data Sheet No. 41915

Conditions of sale

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether from consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Care Australasia Pty Ltd within thirty days of delivery.

UN No. 1170	ETHANOL SOLUTIONS
In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS - AUSTRALIA WIDE
P.G. III	HAZCHEM Z



Crop Care Australasia Pty Ltd
ACN 061 362 347
77 Tingira Street,
PINKENBA Qld 4008

bar codes to be
inserted here

(Item Print Code to be inserted here)

December 7, 2000

page 2 of 8

NRA Approval No.: 48201/1000

BN

DOM

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

KELTHANE[®] MF MITICIDE

ACTIVE CONSTITUENT: 480 g/L DICOFOL

GROUP	2B	INSECTICIDE
-------	-----------	-------------

For the control of mites on certain crops and ornamentals
as specified in the directions for use.

READ THIS LEAFLET BEFORE USING THIS PRODUCT

Crop Care Australasia Pty Ltd, 77 Tingira Street Pinkenba Qld 4008

DIRECTIONS FOR USE:

DO NOT use on crops grown in glasshouses in Tasmania

Crop/ Situation	Pest/Disease	State	Rate/ 100 L	Critical Comments
Babacos	Broad mite Two spotted mite	Qld, WA Only	100 mL	Apply at first appearance of mites (as shown by russetting along the veins in bananas) and repeat as necessary.
Bananas	Two spotted mite		125 mL	
	Strawberry spider mite			
Citrus	Broad mite Brown citrus rust mite Citrus rust mite Citrus bud mite Two spotted mite		100 mL	
Cotton	Spider mites		NSW, WA Only	2.0 L/ha
Grapevines	Two spotted mite	NSW, WA only	63 mL	Commence regular checking of foliage early in the season following extended periods of warm, dry weather or late in the season after harvest for signs of mottling damage indicating the presence of two spotted mite on the undersides of leaves. Spray a miticide early in the season if high numbers and mottling damage on the leaves is clearly visible. Apply thoroughly to the undersides of leaves. If necessary, spray twice 10 days apart.
	Grapeleaf blister mite (<i>Colomerus vitis</i>)	NSW, SA, Vic, WA & NT only	75 mL	Spray at first sign of mite infestation. Repeat as necessary. Thoroughly wet all surfaces.
	Bunch mite (<i>Brevipalpus lewisi</i>) Grapeleaf rust mite (<i>Calepitrimerus vitis</i>)	Qld., NSW, SA, Vic, WA & NT only		
Hops	Two spotted mite	Tas, WA only	100 mL	Apply at first appearance of mites and repeat as necessary.
Mangoes	Mango bud mite Tea red spider mite	Qld, WA Only		
Olives	Two spotted mite			

Crop/ Situation	Pest/Disease	State	Rate/ 100 L	Critical Comments
Ornamentals and Roses	Broad mite Cyclamen mite False spider mite	Qld, Tas, SA, WA, NT only	100 mL	Apply at first appearance of mites (as shown by distorted and twisted leaves, especially near the growing points) and repeat as necessary. NOTE: DO NOT use when control with predatory mites is practiced. NOTE: May damage some house plants, chrysanthemums, gardenias, hibiscus, roses and some varieties of fuchsia especially in hot weather
	Tomato russet mite Two spotted mite	All States		
Paw Paws	Broad mite Two spotted mite	Qld, NSW, WA, NT Only		Apply at first appearance of mite activity if predators are not present (as shown by twisted new leaves or scarring on young leaves or fruit) and repeat 2 weeks later.
Pome fruit Stone fruit	Bryobia mite Two spotted mite European red mite	All States		Apply at first appearance of mites and repeat as necessary. NOTE: May cause russetting in pears or ring spot under some climatic conditions.
	Bryobia mite Two spotted mite European red mite			
Strawberries	Two spotted mite			
Tobacco	Two spotted mite	Qld, NSW, Vic, WA Only	75 mL	
Tomatoes	Tomato russet mite	All States	100 mL	
Vegetables	Two spotted mite Tomato russet mite		115 mL	NOTE: DO NOT use on eggplant (aubergines).

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

DO NOT APPLY TO THE FOLLOWING CROPS LATER THAN 7 DAYS BEFORE HARVEST:

BABACOS, BANANAS, CITRUS, COTTON, HOPS, MANGOES, OLIVES, PAW PAWS, POME FRUIT, STONE FRUIT, STRAWBERRIES, TOMATOES, VEGETABLES:

DO NOT APPLY TO GRAPES LATER THAN 4 WEEKS BEFORE HARVEST.

DO NOT GRAZE OR CUT FOR STOCK FOOD.

GENERAL INSTRUCTIONS

This product mixes readily with water. Add the required quantity to the spray tank with the agitator operating. Add Agral wetting and spreading agent at a rate of 16mL/100 litres to the spray solution, to improve spray coverage.

Insecticide Resistance Warning

GROUP 2B INSECTICIDE

For Insecticide resistance management Kelthane MF Miticide is a Group 2B Insecticide. Some naturally occurring insect biotypes resistant to Kelthane and other Group 2B Insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Kelthane or other Group 2B Insecticides are used repeatedly. The effectiveness of Kelthane on resistant individuals could be significantly reduced. Since the occurrence of resistant insects is difficult to detect prior to use, Crop Care Australasia Pty Ltd accepts no liability for any losses that may result from the failure of Kelthane to control resistant insects.

Kelthane may be subject to specific resistance management strategies. For further information contact your local supplier, Crop Care Australasia Pty Ltd representative or local agricultural department agronomist.

Compatibility

This product is compatible with most commonly used insecticides and fungicides. Do not mix with Bordeaux mixture, lime sulphur, zinc sulphate or carbaryl. As changes in formulation and climatic conditions can alter the sensitivity of plants to mixtures of sprays, Crop Care Australasia Pty Ltd accepts no responsibility for such mixture.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied on any plants at any time. Do not graze livestock in treated areas. Do not feed clippings or residue from sprayed crops to meat or milk-producing animals.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

DO NOT STORE NEAR NAKED FLAMES. Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse empty containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of water ways desirable vegetation and tree roots. Empty containers and product should not be burnt

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing the spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

Additional information is listed in the Material Safety Data Sheet No. 41915

Conditions of sale

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether from consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the

goods to enable proper examination being returned to Crop Care Australasia Pty Ltd within thirty days of delivery.


UN No. 1170	ETHANOL SOLUTIONS
In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS - AUSTRALIA WIDE
P.G. III	HAZCHEM Z



Kelthane is a Registered Trade Mark of Rohm & Haas Company.
Agral is a Registered Trade Mark of Zeneca Limited.



Crop Care Australasia Pty Ltd
ACN 061 362 347
77 Tingira Street,
PINKENBA Qld 4008



(Item Print Code to be inserted here)

NRA Approval No.: 48201/1000