

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# DIFLUFENICAN 500

## Selective Herbicide

ACTIVE CONSTITUENT: 500g/L DIFLUFENICAN

**GROUP F HERBICIDE**

**For control of certain weeds in clover-based Pasture Field Peas, Lentils, Lupins and Oilseed Poppy**

NOT CLASSIFIED AS DANGEROUS GOODS -  
FOR TRANSPORT BY ROAD AND RAIL

**4FARMERS PTY. LTD.**

A.C.N. 067 443 485  
643 MURRAY STREET, WEST PERTH,  
WESTERN AUSTRALIA 6005  
PH (08) 9356 3445 FAX: (08) 9356 3447

APVMA Approval No.: 55669/0402

### GENERAL INSTRUCTIONS

For use as an early post-emergence spray in clover-based pasture, field peas, lentils, and lupins. 4Farmers DIFLUFENICAN 500 may also be used as a post-sowing pre-emergence spray on lupins in NSW, ACT, Vic, SA and Tas. This product provides both contact and residual activity. Residual activity can be expected for up to 8 weeks after application under favourable growing conditions. This product is taken up by the shoots of germinating seeds and seedlings. Susceptible weeds germinate but show immediate chlorosis followed by a mauve-pink discoloration. The chlorosis spreads with the aerial growth and the plants become necrotic and die back. After application, some transient crop discoloration may occur. In lentils and lupins, this usually appears as yellow or white banding on the leaves, while in clover and field peas, white/pink coloration of the leaf veins and tips may occur. Some crop height reduction may also occur. Provided the crop is not under stress from pre-emergent herbicide, disease, insect damage, nutrient deficiency, frost, extremes of pH, dry or excessively moist conditions, the development of the crop and all subsequent growth will not be affected. Some pre-emergence herbicides, such as atrazine, can cause stress to certain crops resulting in an increase in crop damage when using this product. Field peas are particularly sensitive.

#### CLOVER TOLERANCE TABLE VARIETY EFFECT ON VEGETATIVE GROWTH

Arrowleaf (Zulu)	Moderate	Subterranean (Larissa)	Moderate
Balansa (Paradana)	Moderate	Subterranean (Mt Barker)	Moderate
Persian (Kyambro)	Minimal	Subterranean (Seaton Park)	Minimal
Strawberry (Palestine)	Moderate	Subterranean (Trikkala)	Minimal
Subterranean (Clare)	Moderate	Subterranean (Woogenellup)	Moderate
Subterranean (Junee)	Moderate	White (Haifa)	Moderate
Subterranean (Karridale)	Moderate		
Reduction in growth - Minimal (0-20%), Moderate (20-50%)			

The following varieties of subterranean clover have been tested for effects on seed yield: Seaton Park, Trikkala and Woogenellup. Some reduction in seed yield may occur with cv. Trikkala.

Subsequent Crop Tolerance: To reduce the effect on subsequent susceptible crops (eg. canola), ensure thorough cultivation of soil prior to the sowing of these crops.

#### Mixing

Stir product or invert container several times before use as settling may occur after storage for some weeks. To ensure even mixing, half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly then add the remainder of the water. Agitate thoroughly while carrying out spray operations. Reseal part-used container immediately after use.

#### Application

Ground: A minimum water rate of 50 L/ha should be used, however, for optimum results water rates of 70-100 L/ha are recommended. Increase the water volume where weed infestation is heavy or the crop cover is dense. Complete coverage of weeds is essential. Higher water volumes (up to 100 L/ha) will ensure improved activity of the product on the weeds but may increase the symptoms of crop damage.

The following settings are examples that will ensure excellent coverage of exposed weeds:

WATER RATE	50 L/ha	75 L/ha	75 L/ha
NOZZLE	Hardi No. 10 or equivalent	Hardi No. 12 or equivalent	Hardi No. 14 or equivalent
SPEED	10 KpH	10 KpH	12 KpH
PRESSURE	240 KpA (2.4 bar)	220 KpA (2.2 bar)	210 KpA (2.1 bar)

#### COMPATIBILITY:

4Farmers DIFLUFENICAN 500 is physically compatible with most currently registered grass herbicides as two-way tank mixtures.

4FARMERS DIFLUFENICAN 500	COMPATIBLE PRODUCT
Up to 150 mL	Simazine (500g/L product) up to 1.0 L/ha.
All rates	Decis Forte 25EC1, Dimethoate formulations, Alpha Cypermethrin 100EC, Hallmark 50EC, Karate <sup>1</sup> , Omethoate 290 SC, Metribuzin, Talstar <sup>1</sup> and Quizalofop-p-ethyl.

#### Warning:

For tank-mixtures with grass herbicides, use the recommended rates for both herbicides as well as the surfactant recommendations of the grass herbicide. Read the label for the grass herbicide before mixing and using the tank mixtures.

**DO NOT** use crop oils with 4Farmers DIFLUFENICAN 500 or 4Farmers DIFLUFENICAN 500 /grass herbicide tank-mixtures. Applications to lupins and field peas under stressed conditions may cause significant damage to the crop. Tank mixes with simazine should be applied post-emergence to lupins crops only. Increased crop effects may be experienced with the tank mix.

**DO NOT** apply tank-mixtures to clover.

When tank-mixing 4Farmers DIFLUFENICAN 500 and Quizalofop-p-ethyl, use a surfactant only. Mixtures of 4Farmers DIFLUFENICAN 500 and Verdict1 applied to lupins or field peas can cause damage which may result in yield losses. Consult your local 4Farmers Pty Ltd representative or the relevant grass herbicide manufacturer for advice on application and timing of tank-mixtures.

As formulations of other manufacturers' products are beyond the control of 4Farmers Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

#### RESISTANT WEEDS WARNING:

4Farmers DIFLUFENICAN 500 is a member of the Nicotinanalide group of herbicides and acts by inhibiting carotenoid biosynthesis. For weed resistance management 4Farmers DIFLUFENICAN 500 is a group F herbicide. Some naturally occurring weed biotypes resistant to 4Farmers DIFLUFENICAN 500 and other Nicotinanalides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by 4Farmers DIFLUFENICAN 500 or other Nicotinanalide herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, 4Farmers Pty Ltd accepts no liability for any losses that may result from the failure of 4Farmers DIFLUFENICAN 500 to control resistant weeds.

**GROUP F HERBICIDE**

### DIRECTIONS FOR USE

#### RESTRAINTS

DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

DO NOT apply to crops under stress due to pre-emergence herbicide, root disease, insect damage, nutrient deficiency, excessively moist or dry conditions or extremes of pH.

DO NOT apply to frost affected crops or if frosts are imminent.

DO NOT apply if heavy rain is expected within 4 hours.

CROP	WEEDS CONTROLLED	STATE	WEED STAGE	RATE	CRITICAL COMMENTS		
Clover based Pasture Field Peas, Lentils, Lupins	Wild Radish ( <i>Raphanus raphanistrum</i> )	WA only	Up to 2 leaf stage and not more than 60mm in diameter	100mL/ha	<b>CROP STAGE</b> Sow crop evenly to a depth of 20 to 50mm. <b>CLOVER-BASED PASTURE:</b> Apply postemergence, not before the 3rd trifoliate leaf stage. <b>Warning:</b> Some species and varieties of clover may be more sensitive than others. Refer to legume tolerance table in the general instructions. DO NOT apply to medics or yellow serradella. <b>FIELD PEAS:</b> Apply early post-emergence after the third node stage and before the start of flowering. <b>Warning:</b> Field peas grown on high pH soils in the Presence of free lime may be less tolerant to 4Farmers DIFLUFENICAN 500 . <b>LENTILS:</b> Apply early post-emergence after the third node stage of the crop. <b>Warning:</b> Some lentil varieties may be more Sensitive than others. DO NOT apply to Northfield variety. Avoid spray overlap. <b>LUPINS:</b> Post-emergence of Crop: Apply post-Emergence from the 2 leaf to the 6 leaf stage of crop (40 to 100mm high). <b>Post-sowing, Pre-emergence of Crop (Not WA):</b> Apply in a tank mix with the recommended rate of post-sowing pre-emergence treatment of simazine. (4Farmers DIFLUFENICAN 500 should not be incorporated). <b>APPLICATION AND WEED CONTROL</b> Apply when weeds are actively growing. For optimum results apply 4 to 6 weeks post-sowing. Application beyond 8 weeks post-sowing may result in reduced levels of weed control. In most situations the rate specified for each weed size will give satisfactory control. Under certain conditions such as: <ul style="list-style-type: none"> <li>high crop and weed density,</li> <li>late season germinations,</li> <li>abnormal weed growth (including early flowering); higher rates of product (up to the maximum rate of application specified for that weed) may be required.</li> </ul> 4Farmers DIFLUFENICAN 500 will not effectively control : <ul style="list-style-type: none"> <li>regrowth of suppressed weeds,</li> <li>transplanted weeds,</li> <li>regrowth from rhizomes or roots,</li> <li>weeds growing under stress from previous herbicide applications.</li> </ul>		
			Up to 4 leaf stage and not more than 120mm in diameter	150mL/ha			
			Up to 6 leaf stage and not more than 180mm in diameter	200mL/ha			
	Hedge Mustard ( <i>Sisymbrium officinale</i> ) Indian Hedge Mustard ( <i>Sisymbrium orientale</i> ) Wild Turnip ( <i>Brassica tournefortii</i> )	NSW, ACT, Vic, Tas, SA only	Up to 2 leaf stage and not more than 60mm in diameter	100mL/ha			
			Up to 4 leaf stage and not more than 120mm in diameter	150mL/ha			
			Up to 6 leaf stage and not more than 180mm in diameter	200mL/ha			
Turnip Weed ( <i>Rapistrum rugosum</i> )	Charlock (Wild Mustard) ( <i>Sinapis arvensis</i> ) Deadnettle ( <i>Lamium amplexicaule</i> )	NSW, ACT, Vic, Tas, SA, WA only	Up to 4 leaf stage and not more than 120mm in diameter				
						Prickly Lettuce ( <i>Lactuca serriola</i> )	Up to 2 leaf stage and not more than 60mm in diameter

CROP	WEEDS CONTROLLED	STATE	WEED STAGE	RATE	CRITICAL COMMENTS				
<b>SUPPRESSION OF THE FOLLOWING WEEDS</b>									
Clover based Pasture Field Peas, Lentils, Lupins	Capeweed ( <i>Arctotheca calendula</i> ) Crassula ( <i>Crassula spp.</i> ) Corn Gromwell ( <i>Buglossoides arvensis</i> ) Marshmallow ( <i>Malva parviflora</i> ) Shepherd's Purse ( <i>Capsella bursapastoris</i> )	NSW, ACT, Vic, Tas, SA, WA only	Up to 4 leaf stage and not more than 120mm in diameter	200mL/ha	The level of effective residual control may be Reduced where: <ul style="list-style-type: none"> <li>rates lower than 200mL/ha are used,</li> <li>dry conditions prevail, poor coverage of the soil surface is achieved,</li> <li>crop is planted in non-wetting sand,</li> <li>soils have a high content of clay or organic matter.</li> </ul> Where weeds are present at application, good spray coverage of the weeds is important. Apply before weeds are obscured by the crop canopy. Weed control may be reduced in areas where trash or burnt straw from previous harvest is dense, such as in header trails. Best results will be obtained if good soil moisture exists at and after application.				
						Chickweed ( <i>Stellaria media</i> ) Hyssop Loosestrife ( <i>Lythrum hyssopifolia</i> ) Mouse-eared Chickweed ( <i>Cerastium glomeratum</i> ) Night-scented Stock ( <i>Matthiola longipetala</i> ) Skeleton Weed ( <i>Chondrilla juncea</i> ) Speedwell ( <i>Veronica hederifolia</i> )	NSW, ACT, Vic, Tas, SA only		
								Amsinckia ( <i>Amsinckia spp.</i> ) Wireweed ( <i>Polygonum aviculare</i> )	NSW, ACT, Vic, Tas, SA only
						Sorel ( <i>Rumex acetosella</i> ) Toad Rush ( <i>Juncus bufonius</i> )	NSW, ACT Vic, Tas, SA only		
								Stinging Nettle ( <i>Urtica urens</i> )	NSW, ACT only
						Oilseed Poppy	Charlock ( <i>Sinapis arvensis</i> ) Hedge Mustard ( <i>Sisymbrium officinale</i> ) Indian Hedge Mustard ( <i>Sisymbrium orientale</i> ) Wild Radish ( <i>Raphanus raphanistrum</i> ) Wild Turnip ( <i>Brassica tournefortii</i> )	Tas only	Early postemergence up to the 4 leaf stage and not more than 120mm in diameter

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD** NOT REQUIRED WHEN USED AS DIRECTED

#### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

#### STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

#### SAFETY DIRECTIONS

Avoid contact with eyes and skin. Wash hands after use.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (telephone 13 11 26).

#### MATERIAL SAFETY DATA SHEET

If additional hazard information is required refer to the Material Safety Data Sheet.

For a copy phone 08 9356 3445 or visit our website at www.4farmers.com.au

#### CONDITIONS OF SALE

This product as supplied is of a high grade and suitable for the purpose for which it is expressly intended and must be used in accordance with the directions. The user must monitor the performance of any product as climatic, geographical or biological variables and/or developed resistance may affect the results obtained. No responsibility is accepted in respect of this product, save for those non-excludable conditions implied by the Trade Practices Act or any State or Federal legislation.

NOT CLASSIFIED AS DANGEROUS GOODS FOR TRANSPORT BY ROAD.

**IN A TRANSPORT EMERGENCY  
DIAL 000  
POLICE OR FIRE BRIGADE**

**FOR 24 HOUR SPECIALIST ADVICE IN EMERGENCY ONLY**

**PHONE 1800 638 556**

<sup>1</sup> is not a Registered Trademark of 4Farmers Pty Ltd.