

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm
Diuron
900DF

Herbicide

ACTIVE CONSTITUENT: 900g/kg DIURON

GROUP C HERBICIDE

For selective weed control in the situations as per the Directions for Use table.

THE BAG MUST NOT BE SOLD SEPARATELY.
DO NOT DESTROY BOX WHILE PRODUCT STILL REMAINS.

**IMPORTANT: READ THE ATTACHED LEAFLET
BEFORE USING THIS PRODUCT.**

Net Contents: 10kg

Nufarm Australia Limited
ACN 004 377 780
103-105 Pipe Road
Laverton North Victoria 3026
Tel: (03) 9282 1000
Fax: (03) 9282 1001



STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN.

Store in the closed, original container in a dry, cool, well ventilated area out of direct sunlight. Shake empty bag into container. **DO NOT** dispose of undiluted chemicals on site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should **NOT** be burnt.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. **DO NOT** inhale dust or spray mist. If product on skin immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS).

CONDITIONS OF SALE

"Nufarm Australia Limited('Nufarm') shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Nufarm's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Nufarm has any authority to add to or alter these conditions."

In case of emergency: Phone 1800 033 498

Ask for shift supervisor. Toll free 24 hours.

APVMA Approval No.: 39201/10kg/0507

Batch number:

Date of manufacture:

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm
Diuron
900DF

Herbicide

ACTIVE CONSTITUENT: 900g/kg DIURON

GROUP  HERBICIDE

For selective weed control in the situations as per the Directions for Use table.

**NOT TO BE SOLD SEPARATELY.
BEFORE USING PRODUCT, READ ALL DIRECTIONS
ON OUTER PACK AND LABEL LEAFLET.**

SALE OF THIS BAG EXCEPT IN OUTER PACK IS ILLEGAL.

Net Contents: 10kg

APVMA Approval No. 39201/10kg/0507

Nufarm Australia Limited
ACN 004 377 780
103-105 Pipe Road
Laverton North Victoria 3026
Tel: (03) 9282 1000
Fax: (03) 9282 1001



STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN.

Store in the closed, original container in a dry, cool, well ventilated area out of direct sunlight. Shake empty bag into container. DO NOT dispose of undiluted chemicals on site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. If product on skin immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 13 11 26.

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm
Diuron
900DF

Herbicide

ACTIVE CONSTITUENT: 900g/kg DIURON

GROUP **C** HERBICIDE

For selective weed control in the situations as per the Directions For Use table.

READ THIS LEAFLET BEFORE USING THIS PRODUCT.

APVMA Approval No. 39201/0507

Nufarm Australia Limited
ACN 004 377 780
103-105 Pipe Road
Laverton North Victoria 3026
Tel: (03) 9282 1000
Fax: (03) 9282 1001



DIRECTIONS FOR USE:

RESTRAINTS:

DO NOT replant treated areas within 2 years of application of Diuron 900DF except when otherwise stated on label. **DO NOT** use on very light sandy or gravelly soils as crop damage may occur.

Crop	Weeds Controlled	State	Rate	Critical Comments
Apples & Pears (established at least 1 year)	Barnyard grass, Capeweed, Barley grass, Fat hen, Amaranthus, Mustards, Ryegrass, Wild oats, Wild radish, Wild turnip, Blackberry nightshade, Cobbler's pegs, Crowsfoot grass, Guinea grass, Pigweed, Sowthistle, Summer grass	All States	Single application 4kg/ha Split application 2kg/ha	Apply under trees as a broadcast or band spray. Area should be cleared of weeds prior to spraying Diuron 900DF using Weedmaster® Duo Herbicide. SINGLE APPLICATION: Apply in winter. SPLIT APPLICATION: Apply post harvest, with a second application in spring.
	Wireweed, Toad rush	Qld, NSW, Vic, ACT, SA, WA only		
	Willow herb	Tas only		
Asparagus	Annual Broadleaf weeds and some Annual grasses	All States	1.5 - 2kg/ha	Apply in established beds after they have "hilled up" and before emergence of spikes. This will give 10-14 weeks freedom from weeds. Post emergence application may be made, but the spray should be directed so as to contact only the base of the stems. Use higher rates on heavier soils DO NOT USE ON SEEDLING ASPARAGUS.
Bananas	Amaranthus, Barley grass, Barnyard grass, Capeweed, Cobblers pegs, Crowsfoot grass, Fat hen, Guinea grass, Mustard, Pigweed, Prince of Wales feather, Rye grass, Sowthistle, Summer grass, Wild radish, Wild turnip.	NSW, Qld, WA only	2 - 4kg/ha	At planting: Apply following fill-in planting. DO NOT spray banana foliage. Use higher rate on heavier soil types. Apply in 200 to 500L water/ sprayed ha. Established plantations: Apply avoiding contact with banana plants. Use higher rate on heavier soil types. Add Nuquat® 250 if weed growth is above 8cm high.
			275g - 500g/ha Plus 1.2 or 1.6L/ha Nuquat 250	Add to Nuquat® 250 to enhance weed control under favourable growing conditions and in open sunny conditions. Diuron 900DF at rates up to 500g/ha can be blanket sprayed. Use a directed spray for higher rates of Diuron 900DF. Complete spray coverage is essential. For grasses and broadleaved weeds up to 5cm high use a minimum of 250L spray solution/ha, increase to 350 L/ha for weeds up to 10cm high. Use a minimum spray volume of 400L/ha for weeds >10cm high and for dense mature weeds. Refer to Nuquat label for rates and full directions. If Nuquat 250 rate is less than the ratio 500mL/100L add 100mL Surfactant 600 or 60mL Chemwet 1000 /100L spray mix.
Citrus (established at least 1 year)	Seedlings of: Guinea grass, Barnyard grass, Billygoat weed, Wild turnip, Crowsfoot grass, Cobbler's pegs, Wild mustard, Red natal grass, Feather-top, Wild radish, Rhodes grass, Pigweed, Blackberry nightshade, Spiny burr grass, Winter grass, Wild oats, Capeweed, Wireweed, Fat hen, Sowthistle, Summer grass, Barley grass, Ryegrass, Prince of Wales feather, Amaranthus	All States	2 - 4kg/ha plus 500mL Surfactant 600 per 100L spray mixture	Apply only to trees one year or more old. Apply after cultivation in March or April. DO NOT add Surfactant 600 if spray is likely to contact citrus foliage. DO NOT use on very light sands. Apply in up to 4000L water/ha. Use higher rate on heavier soils and if weeds up to 15 cm in height are present.
	Couch Grass		350g/100L	SPOT SPRAYING - Spray to the point of run-off.
Coffee	Seedlings of: Summer grass, Barnyard grass, Billygoat weed, Sourgrass	Qld, NSW only	3.9kg/ sprayed ha	Initial application: Destroy all vegetative growth by spraying with Nuquat 250 before applying Diuron 900DF.
			1 - 1.9kg/ha plus 600 - 1.1L Nuquat 250 plus Surfactant 600	Subsequent application: Frequency of application will depend on regrowth. Use higher rates if weed growth is dense.
	Couch grass		710g/200L	SPOT SPRAYING - Spray to the point of run-off.

Crop	Weeds Controlled	State	Rate	Critical Comments
Cotton	Annual broadleaf weeds and some Annual Grasses	Qld, NSW only	1 - 2kg/ha	<p>Pre-emergence: Apply at planting or within 7 days of planting and prior to cotton emergence. Seed should be planted at least 4 to 5cm deep and soil compacted over seeds by a normal planter packer wheel. Use high rate on soils high in clay or organic matter content. DO NOT use on light sandy soils as injury to crop may occur. DO NOT make a second pre-emergent application to the same field during the same crop year as injury to the crop may result. Apply in 300 to 450L water/ha.</p> <p>Post-emergence (lay-by): Use the higher rate on heavier soil types high in clay and/or organic matter. DO NOT use on light sandy soils low in organic matter as injury to the cotton may result. Apply when cotton is at least 30cm high, or immediately after the last cultivation. If emerged annual weeds (15cm or less in height) are present add 170mL Nufarm Surfactant 600/100L water to the spray solution. Apply Diuron 900DF as a directed spray to cover the area beneath cotton plants and between rows. Avoid spray contact or drift on to cotton plants as injury may result. DO NOT apply more than once per season.</p> <p>DO NOT re-plant treated areas to any crop within one year after last application as injury may result, except cotton, corn or sorghum (grain sorghum only) which may be planted in the spring following treatment. Apply in 300-500 L water/ha.</p>
Gladioli, Tulips, Daffodils, Lilium and Iris (Apply at least 4 weeks before bulb emergence)	Seedlings of: Summer grass, Barnyard grass, Mustard, Barley grass, Ryegrass, Wild oats, Capeweed, Wild radish, Wild turnip, Blackberry nightshade, Cobbler's pegs	All States	2kg/ha	<p>Apply in a high volume spray, to moist, bare soil before weeds emerge. DO NOT replant areas to any crop within 1 year of treatment, as crop damage may occur. DO NOT use on light sandy or gravelly soils, or soils low in organic matter as it may cause damage to crops, particularly Liliums.</p>
	Amaranthus, Annual ryegrass, Barley grass, Barnyard grass, Blackberry nightshade, Capeweed, Cobblers pegs, Crowsfoot grass*, Fat hen, Guinea grass*, Mustards, Pigweed, Redshank*, Sow thistles*, Summer grass, Thistles, Wild oats, Wild radish, Wild turnip, Winter grass, Wireweed	Tas only	1.5 - 2.0kg/ha	<p>Sufficient moisture in the form of rain or irrigation is necessary within 3-4 days of application, to move the product into the weed root-zone. Heavy rains after application may cause severe crop damage.</p> <p>* may require 2.0kg/ha for control.</p>
Grass Seed Crops Established (at least 1 year old) fescue, phalaris and cocksfoot Phalaris cultivars Sirolan and Sirosa	Annual Broadleaf weeds and some Annual Grass weeds	SA only	2.8kg/ha	Apply at weed emergence by boom spray. Use higher rate for heavier soils. Brome grasses will not be controlled.
		Qld, NSW, Vic, ACT & WA only	3 - 3.3kg/ha	
		All States	1.9kg/ha	This rate should be used for follow-up sprays in the 2nd year.
			1.7kg/ha	DO NOT use as a tank mix with Nuquat 250
Lucerne Established for 1 year or more	Seedlings of Capeweed, Wireweed, Chickweed, Summer grass, Docks, Barnyard grass, Thistles, Billygoat weed, Spiny emex, Ryegrass, Barley grass, Sheperds' purse, Fat hen	All States	1 - 1.9kg/ha	Apply when lucerne has been grazed. Weeds should be very small when treated. Use only in areas where lucerne becomes winter dormant. Use higher rate on heavier soil.
	Black nightshade, Wild oats, Wild radish, Winter grass, Amaranthus, Mustard, Wild turnip	Tas only		

Crop	Weeds Controlled	State	Rate	Critical Comments
Lupins (on light soils only)	Annual ryegrass, Capeweed, Clovers, Medics, Doublegee, Wild radish, Wild turnip and suppression only of Dock, Sorrel, Wireweed, Wild oats and Brome grass	WA only	1.1kg/ha	Use through boom spray equipment only. Apply in a minimum of 50L water/ha. Higher volumes of water will give more even application. Must be applied prior to crop emergence. DO NOT use on white or grey sands
Paw Paws (not for use on trees under 9 months old)	Seedlings of Guinea Grass, Crowsfoot Grass, Barnyard Grass, Billygoat weed, Summer grass, Red natal grass, Giant sensitive plant	Qld, NSW, only	4 kg/ha	Avoid contact with plants. Use higher rates on heavier soils.
			2 kg/ha	Apply a second spray if required. DO NOT exceed 6.5 kg/ha per year. Add Nuquat 250 to Diuron 900DF if weed growth is more than 8cm high.
Peas	Annual ryegrass, Barley grass, Brome grass, Capeweed, Doublegee, Wild Oats, Wild turnip, Wild radish, Mustards	WA only	835g - 1.2kg/ha	DO NOT USE ON SANDY SOILS. Use the lower rate pre-sowing in the low rainfall eastern wheat belt and use the higher rate immediately after sowing in the Great Southern Area. Weed control may be increased by mixing with Trifluralin pre-sowing.
Pineapples	Annual broadleaf weeds and some Annual grasses	Qld, NSW only	3.9kg/ha	Note: DO NOT apply more than 8kg/ha before fruit differentiation of flower bud formation. Apply as a blanket spray after planting and before weed emergence. Ratoon Crop: Apply immediately prior to the first rains on weed free soil. Apply in up to 4000L water/ha.
			1.9kg/ha	Follow-up sprays, if necessary, should be made at this lower rate.
Pulses (Including Chickpeas, Faba Beans, Lentils, Narbon beans, Field Peas and Vetch)	Cape weed, Double Gee, Wild radish, Wild Turnip, Toad rush, Crassula, and Erodium.	All States	IBS 830g-1.1kg/ha PSPE 550-830g/ha	Incorporated by sowing (IBS) Apply to bare moist soil prior to sowing or at sowing and incorporate by the sowing operation. Use the lower rate on light sandy soils. Post Sowing Pre-Emergent Application. (PSPE) Apply as a post sowing pre-emergent application to moist soil before weed or crop emergence. Use the lower rate on light sandy soils. Application should not be made to excessively ridged or waterlogged soils. Sow the crop at least 5cm deep. Trifluralin or Spinnaker may be tank mixed at the recommended rates.
Sugar Cane (plant and ratoon crops)	Barnyard grass, Crowsfoot grass, Guinea grass, Summer grass, Barley grass, Ryegrass, Wild oats, Cobbler's pegs, Pigweed, Prince of Wales feather, Amaranthus, Fat hen, Mustard, Sowthistle, Capeweed, Wild radish, Wild turnip	Qld, NSW, only	3.9kg/ha	Pre-emergence of cane and weeds. Soil surface must be moist at time of application. Rain or irrigation is required within 10 days of application
			1.9 - 3.9kg/ha	Post-emergence of cane, as for pre-emergence. Use lower rate if following an application at planting. If weeds at early post-emergence stage addition of Surfactant 600 at 250mL/100L of water will enhance activity, but direct spray to below cane leaves.
			275g/ha - 3.9kg/ha plus 1.2L - 1.6L Nuquat 250	Add to Nuquat 250 to enhance weed control under favourable growing conditions and in open sunny conditions. Diuron 900DF at rates up to 500g/ha can be blanket sprayed. Use a directed spray for higher rates of Diuron 900DF. Complete spray coverage is essential. For grasses and broadleaved weeds up to 5 cm high use a minimum of 250L spray solution/ha, increase to 350 L/ha for weeds up to 10cm high. Use a minimum spray volume of 400L/ha for weeds > 10cm high and for dense mature weeds. Refer to Nuquat label for rates and full directions. Note: Where broad leaved weeds are a serious problem add Tornado® DF to the mixture.
Summer Fallows	Potato weed (<i>Heliotropium europaeum</i>)	SA only	275g/ha	Only effective when in mixture with Spray•Seed * 250 at 1.6L/ha (See Spray•Seed * 250 label)
Tea plantation	Pre-emergence weed control, especially Guinea grass and other grass weeds.	Qld only	1kg/ha	Use ONLY on tea headlands where soil movement or erosion into cropping areas or waterways is minimal or nil. Apply one boom spray application per year prior to onset of wet season. Avoid contact of spray with the crop. DO NOT use in crop inter-row areas.

Crop	Weeds Controlled	State	Rate	Critical Comments	
Vineyards	Barnyard grass, Crowsfoot grass, Guinea grass, Summer grass, Barley grass, Ryegrass, Wild oats, Cobbler's pegs, Prince of Wales feather, Amaranthus, Fat hen, Mustard, Sowthistle, Capeweed, Wild radish, Wild turnip	All States	2 - 4kg/ha	Band Treatment to crop rows. Apply to established vines (at least 3 years old). Apply to moist soil prior to emergence of weeds. DO NOT apply to vines of butt diameter less than 4cm. Use the lower rate where annual and winter weeds are to be controlled. Avoid spray drift or contact onto fruit or foliage.	
	Winter grass, Blackberry nightshade, Thistles	Tas only			
	Annual broadleaf weeds & Some annual grasses	All States	85g/20m ²	Under racks - apply after harvest.	
Wheat, Barley, Triticale, Cereal Rye, Oats	Capeweed, Doublegee, Mustard, Wild Radish [supression only], Water Weeds [Toads Rush, Crassula]	WA only	275g - 550g *Barley to 390g only	Post planting application only for Full cut sowing systems that leave a relatively smooth surface, eg. standard combine or airseeder. Apply post plant pre-emergent to crop within 3-4 days of planting. Pre plant application only for sowing systems (eg NoTill or ZeroTill) which leave open slots or furrows, or where seed is sown along the bottom of the furrow. Even the soil surface where harrowed after sowing. Apply within 7 - 0 days pre plant. Soil Types - use lower rates on sandy surfaced soils with a low organic carbon or clay content. Caution: DO NOT use on grey or white sands. Crop to be planted at least 35mm deep. Crop tipping and yellowing may occur particularly when heavy rain falls after sowing. Shallow sown crops are more at risk. Caution: Severe crop damage will occur if herbicides containing Paraquat or Diquat contact an emerging cereal crop either as a salvage treatment or via spray drift unless rainfall has been exceptionally low.	
	As above plus suppression of Annual Ryegrass, Barley Grass and Silver Grass		550g/ha plus 500ml of a 720g/L metolachlor to improve suppression of Ryegrass		
Wheat, Barley, Triticale, Cereal Rye			275g - 550g plus 830mL/ha of TriflurX	As Above Must be incorporated within 4 hours of application.	
Wheat, Barley, Oats	Capeweed, Charlock, Iceplant, Melilotus, Mustards, Rough Poppy, Saffron thistle, Sheepweed, Three cornered Jack, Wild turnip, Yellow burr weed, Deadnettle	NSW, Vic, ACT & SA only	500g/ha	Apply when weeds are at the 2-4 leaf stage and soon after the crop is past the 2-5 leaf stage which is normally within 6 weeks of sowing. DO NOT SPRAY LATER THAN EARLY TILLERING STAGE.	
Wheat, Barley, Triticale, Oats	Capeweed, Doublegee, Erodium, London Rocket, Lupin, Mustard, Rapistrum, Wild Radish, Wild turnip, Storksbill, Wild mustard	WA only	200g/ha plus 400mL MCPA 500	Apply when the crop has at least 3 to 5 leaves on the main stem. (Zadoks 13-15)	Apply by boom spray or aircraft. Weeds must be in the 2-5 leaf stage. Crop and weeds should be dry at the time of spraying. DO NOT spray if rain imminent. Undersown sub-clovers may be slightly retarded. DO NOT apply to undersown medics. Apply in 40-70L water/ha.
			280g/ha plus 200mL Amicide® 625	Apply when crop has at least 4 to 5 leaves on main stem. (Zadoks 14-15)	
Wheat, Barley	Amsinkia (Yellow burr), Capeweed, Deadnettle, Double gee (Three cornered Jack), Mustard, White ironweed (Sheep weed), Wild radish	NSW, Vic & ACT, SA only	280g/ha plus 500mL/ha (NSW, Vic, ACT only) 350mL/ha (SA only) MCPA 500	Apply by boomspray or aircraft when the crop is at the 3-5 leaf stage (NSW, SA only) 5 leaf stage (Vic only). Weeds must be in the 2-5 leaf stage. DO NOT spray if rain is imminent. DO NOT apply to under-sown medic or lucerne. DO NOT use on Durati, hybrid Titan, Kite, Kamilaroi, or Shortim Wheat.	
	Soursob	SA only	710 - 980g/ha	Boom Spray: Sow crop at least 5cm deep. Apply when crop is in the 2-5 true leaf stage. Use the lower rate on lighter soils.	
Irrigation Channels & Drainage Ditches	Annual and Perennial grasses and Broadleaf weeds. (Except Nutgrass, Bindweed and Russian knapweed)	SA, WA, Vic, Qld, Tas only	22kg/ha OR 220g/100m ²	High Volume or Power Spraying. Apply in 2000L water. Apply during the non-crop season and when channel is not in use. Soil should be moist and preferably bare ie. after regrading and cleaning channels. Apply prior to expected seasonal rainfall so as to "fix" it into the soil. If 50-100mm of rain has not fallen after application and it is necessary to use the irrigation channel, fill it with water and allow to stand for 72 hours, then drain off the water in the channel and run to waste.	
		NSW only	40kg/ha maximum		

Crop	Weeds Controlled	State	Rate	Critical Comments
Rights of Way, Commercial & Industrial Areas	Annual grasses, Perennial grasses and Broadleaf weeds. (Except Nutgrass, Johnson grass, Bindweed and Russian knapweed)	All States		If moderate weed growth ie. greater than 15cm in height is present at the time of spraying add a knockdown herbicide such as Amitrole T to the mixture. Apply to short growth. Apply just before or during the active growing season. DO NOT treat during the dry season. Time application so that rainfall afterwards will take the chemical down to the root zone for uptake by the roots. For application by Railway Tanker or Boom Spray apply in 400 - 800L water/ha. For power spraying or hand spraying equipment apply the equivalent of at least 2000L water/ha. Ensure entire area is treated - including bare ground.
			4.5 - 18kg/ha 18 - 36kg/ha 27 - 40kg/ha	Railway Tanker or Boom Sprayer - Initial Treatment: Annual rainfall 0 - 500mm Annual rainfall 500 - 1000mm Annual rainfall 1000mm and over Use higher rates in the ranges shown on soils with high clay or organic matter content and where maximum duration of control is required.
			3.5 - 7kg/ha 7 -13.5kg/ha 9 - 18kg/ha	Re - Treatment Annual rainfall 0 - 500mm. Annual rainfall 500 - 1000mm. Annual rainfall 1000mm and over.
			220 - 890g/100L 890g - 1.8kg/100L 1.3 - 2kg/100L	High Volume or Power Spraying - Initial Treatment: Annual rainfall 0 - 500mm. Annual rainfall 500 - 1000mm. Annual rainfall 1000mm and over.
			180 - 355g/100L 355 - 670g/100L 450 - 890g/100L	Re - Treatment Annual rainfall 0 - 500mm. Annual rainfall 500 - 1000mm. Annual rainfall 1000mm and over.
Small Areas			90g in 15L knapsack/ 20m ²	Apply evenly.
Bore Drains	Prickly acacia, Mimosa bush	Qld only	35.5kg/ha	Block off bore drain 24 hours prior to spraying and allow drain to empty. Spray 1m wide strip onto mud. Wait 48 - 72 hours before opening drain again. Water MUST by-pass house tanks and desirable tree stands for 7 days after opening drain. DO NOT delve for 2 weeks.
Non-Crop Areas	Sensitive plant (Giant)	WA only	2 - 4kg/ha or 20 - 40g/10L	Apply to moist soil.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: DO NOT ALLOW ANIMALS TO DRINK WATER FROM TREATED BORE DRAINS FOR 3 DAYS BEFORE SLAUGHTER FOR HUMAN CONSUMPTION.

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.

GRAZING:

PULSE CROPS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 35 DAYS AFTER APPLICATION. ALL OTHER CROPS: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions and First Aid Instructions.

Application should be made to moist soil before weeds emerge unless otherwise recommended. Sufficient moisture in the form of rain or irrigation is necessary after application to move the chemical into the weed root zone. Add Nufarm Surfactant where recommended. When spraying emerged weeds best results are obtained when weeds are actively growing. Thoroughly clean all traces of this product from application equipment immediately after use, otherwise injury to desirable vegetation may result when equipment is used again. Flush tank, pump, hose and boom with several changes of water after removing nozzle tip and screens (clean these parts separately). Keep from contact with fertilizers, insecticides, fungicides and seeds.

MIXING

Partly fill spray tank with water, and agitate while adding this product steadily. Continue agitation after filling and during spraying.

Agitation: An efficient by-pass agitator or means of mechanical agitation should be employed when using this product.

COMPATIBILITY

This product is compatible with Spray•Seed* 250 and the following Nufarm products Amicide® 625, Nuquat® 250, MCPA 500. Add the required amount of product when the spray tank containing Diuron 900DF is nearly full and being continually agitated. This product is not compatible with Tribunil*, Rogor*, Mataven*, Hoegrass*/Nugrass®, and Bayleton*.

HERBICIDE RESISTANCE WARNING



'Diuron 900DF' Herbicide is a member of the urea group of herbicides and has the inhibitors of photosynthesis at photosystem II mode of action. For weed resistance management Diuron 900DF is a Group C herbicide. Some naturally occurring weed biotypes resistant to Diuron 900DF and other Group C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Diuron 900DF or other Group C herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from the failure of Diuron 900DF Herbicide to control resistant weeds.

WEED CONTROL ON NON-CROP LAND

Diuron 900DF Herbicide is an effective herbicide for the residual control of many annual and perennial grasses and broadleaf weeds on non-crop land such as rail tracks, roadways, firebreaks, industrial sites, fence lines, storage areas, irrigation and drainage

ditches where bare ground is desired. Established perennials such as Bindweed, Johnson grass, Nutgrass, and Russian Knapweed are not satisfactorily controlled at rates recommended. Duration of control depends on the amount of chemical applied, soil type, rainfall and particular weed species.

SELECTIVE USE IN CROPS

For selective use in crops it is important to apply recommended rates correctly. Diuron 900DF Herbicide is most effective when applied to soil which has been well tilled (clod free) and is free of weeds and trash. Where not recommended for overall application, care should be taken to avoid spray drift onto fruit or foliage. Wetting agents should only be used where specified.

REPLANTING RESTRICTIONS

Crop Rotations: DO NOT replant treated areas to any crop within two years after last spray unless otherwise stated.

Cotton and Lucerne - DO NOT replant treated areas to any crop within 1 year after last spray. Except cotton, corn or grain sorghum which may be planted in the spring of the following year.

Gladioli, Tulip, Daffodils, Lilium, Iris - DO NOT replant treated areas to any crop within 1 year after treatment.

Pineapples and Sugarcane - treated areas can be replanted to pineapple or sugarcane 1 year after last spray.

CROP SAFETY

Heavy rains after application of this product may cause severe crop damage, risk must be assumed by the user.

Crop Safety: Apples and Pears - DO NOT use under dwarf or semi-dwarf trees.

Grapevines - DO NOT apply to vines of butt diameter less than 4cm. On soils low in clay or organic matter, plant injury may occur if heavy rains follow treatment.

DO NOT USE ON SANDY OR GRAVELLY SOILS OR SOILS LOW IN ORGANIC MATTER.

AVOID CONTACT OF SPRAYS WITH CROP FOLIAGE OR FRUIT.

PROTECTION OF CROPS, NATIVE & NON - TARGET PLANTS.

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used container.

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN.

Store in the closed, original container in a dry, cool, well ventilated area out of direct sunlight. Shake empty bag into container. **DO NOT** dispose of undiluted chemicals on site. Puncture or shred and bury the empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should **NOT** be burnt.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. **DO NOT** inhale dust or spray mist. If product on skin immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS), which can be obtained from your supplier or from the Nufarm website - www.nufarm.com.au.

CONDITIONS OF SALE

"Nufarm Australia Limited ('Nufarm') shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Nufarm's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Nufarm has any authority to add to or alter these conditions."

In case of emergency: Phone 1800 033 498

Ask for shift supervisor. Toll free 24 hours.

® Amicide, Nuquat, Nugrass and Tornado are registered trademarks of Nufarm Australia Limited.

* Other registered trademarks