

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

DIUREX® WG Herbicide

ACTIVE CONSTITUENT: 900 g/kg DIURON

GROUP	C	HERBICIDE
--------------	----------	------------------

For selective weed control in the situations as per the directions for use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

Crop Care Australasia Pty Ltd
ACN 061 362 347
Unit 15/16 Metroplex Avenue
Murarrie QLD 4172
Tel: (07) 3909 2000



APVMA Approval No.: 45772/63080

Contents: 10kg, 15kg

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN. Store in the closed, original container in a dry, cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Shake empty box or bag into spray tank. Single rinse bag before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500mm in a disposal pit specifically marked and set up for the purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from your supplier, or from the Crop Care website – www.cropcare.com.au


Conditions of sale

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Care Australasia Pty Ltd within thirty days of delivery.

In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 498 ALL HOURS - AUSTRALIA WIDE
---	--

® Diurex is a registered trademark of, or used under licence by Crop Care Australasia Pty Ltd.

BN / DOM / Barcode / Drummuster / 10kg, 15kg



ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (CONTAINS DIURON)
UN No: 3077
PACKING GROUP: III
HAZCHEM: 2Z
IN AN EMERGENCY DIAL: 000 POLICE OR FIRE BRIGADE
IN AN EMERGENCY, SPECIALIST ADVICE NUFARM AUSTRALIA LIMITED 1800 033 498

"NOTE: MANUFACTURED IN AUSTRALIA – DG CLASS 9 MARKING ONLY REQUIRED FOR SHIPMENT TO TASMANIA – THIS IS DONE BY NUFARM LOGISTICS"

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

DIUREX[®] WG Herbicide

ACTIVE CONSTITUENT: 900 g/kg DIURON

GROUP	C	HERBICIDE
--------------	----------	------------------

For selective weed control in the situations as per the directions for use table.

BEFORE USING PRODUCT, READ DIRECTIONS ON OUTER PACK.

SALE OF THIS BAG EXCEPT IN OUTER PACK IS ILLEGAL

Crop Care Australasia Pty Ltd
ACN 061 362 347
Unit 15/16 Metroplex Avenue
Murarrie QLD 4172
Tel: (07) 3909 2000



APVMA Approval No.: 45772/63080

Contents: 10kg (inner bag to box)

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN. Store in the closed, original container in a dry, cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Single rinse bag before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. Puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500mm in a disposal pit specifically marked and set up for the purpose

SAFETY DIRECTIONS

Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from your supplier, or from the Crop Care website – www.cropcare.com.au

Conditions of sale

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Care Australasia Pty Ltd within thirty days of delivery.

In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 498 ALL HOURS - AUSTRALIA WIDE
---	--

® Diurex is a registered trademark of, or used under licence by Crop Care Australasia Pty Ltd

BN / DOM / Barcode / Drummuster / 10kg

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

DIUREX[®] WG Herbicide

ACTIVE CONSTITUENT: 900 g/kg DIURON

GROUP	C	HERBICIDE
--------------	----------	------------------

For selective weed control in the situations as per the directions for use table.

READ COMPLETE DIRECTIONS FOR USE BEFORE USING THIS PRODUCT.

Crop Care Australasia Pty Ltd
ACN 061 362 347
Unit 15/16 Metroplex Avenue
Murarrie QLD 4172
Tel: (07) 3909 2000



APVMA Approval No.: 45772/63080

DIRECTIONS FOR USE

RESTRAINTS

DO NOT replant treated areas within 2 years of application of Diurex WG except when otherwise stated on label.
DO NOT use on very light sandy or gravelly soils as crop damage may occur.
DO NOT use in water-logged areas.
DO NOT apply if greater than 50mm rainfall is expected within 3 days of application.
DO NOT irrigate within 3 days of application.
DO NOT apply to fields where the slope exceeds 3%.
DO NOT spot spray more than 5% of total farm areas.
DO NOT apply more than once per calendar year.

Cotton

DO NOT use in cotton UNLESS ALL irrigation tailwater and/or up to 25mm rainfall can be captured and held on farm.

Pulse crops

DO NOT USE in pulse crops UNLESS application can be made using no-till farming practices.

Lucerne

DO NOT use in lucerne UNLESS ALL irrigation tailwater can be captured and held on farm.

Irrigation channels and drains

DO NOT use in irrigation channels or drains UNLESS ALL irrigation tailwater and rainfall can be captured and held on farm.

Desert channels (bore drains)

DO NOT apply between 1 December and 30 March.
DO NOT open drains for 72 hours following treatment.

SPRAY DRIFT RESTRAINTS

DO NOT apply by air.
DO NOT apply when wind speed is less than 3 or more than 20 kilometres per hour as measure at the application site.
DO NOT apply with spray droplets smaller than COARSE spray droplet size category according to nozzle manufacturer specifications that refer to the ASAE S572 Standards or the BCPC Guideline.

Users of this product MUST make an accurate written record of the details of each spray application within 24 hours following application and KEEP this record for a minimum of 2 years. The spray application details that must be recorded are: 1. Date with start and finish times of application; 2. Location address and paddock/s sprayed; 3. Full name of this product; 4. Amount of products used per hectare and number of hectares applied to; 5. Crop/situation and weed/pest; 6. Wind speed and direction during application; 7. Air temperature and relative humidity during application; 8. Nozzle brand, type, spray angle, nozzle capacity and spray system pressure measured during application; 9. Name and address of person applying this product. (Additional record details may be required by the State or Territory where this product is used.)

MANDATORY NO-SPRAY ZONES

DO NOT apply when there is non-target vegetation downwind from the application area and within the mandatory no-spray zones shown in table below.

DO NOT apply when there are aquatic and wetland areas including aquacultural ponds, surface streams and rivers downwind from the application area and within the mandatory no-spray zones shown in table below.

SITUATION	RATE (g ac/ha)	DOWNWIND BUFFER ZONE (M)	
		AQUATIC	TERRESTRIAL
Wheat, Barley, Triticale, Cereal rye, Oats (WA only)	250-500	30	60
Wheat, Barley, Triticale and Oats	180-250	15	30
Wheat, Barley, Oats (NSW, Vic, ACT, SA only)	450	25	50
Wheat and Barley – SA only	640-880	50	100
Wheat and Barley – NSW, Vic, ACT, SA only	250	15	30
Summer fallows – SA only	250	15	30
Cotton	900-1800	100	200
Lucerne	750	50	100
Lupins – WA only	990	30	80
Pulses – incorporated by sowing	750-990	30	80
Pulses – post-sowing pre-emergent	495-750	50	100
Sugarcane and Bananas	250-450	25	50
Sugarcane	1800	100	200
Asparagus	1800	100	200

DIRECTIONS FOR USE

Crop/Situation	Weeds Controlled	State	Rate	Critical Comments
Asparagus	Amaranthus, Barley grass, Barnyard grass (Water grass), Capeweed, Cobbler's peg, Crowsfoot grass, Fat hen (White goosefoot), Guinea grass, Milk thistle (Sowthistle), Mustards, Pigweed, Summer grass, Wild oats, Wild radish, Wild turnip, Wimmera ryegrass (Annual ryegrass)	Vic only	1.5-1.9kg/sprayed ha	Apply in established beds after they have been "hilled up" and before emergence of spikes. This will give 10 to 14 weeks freedom from weeds. Post-emergence applications may be made, but the spray should be directed so as to contact only the base of the stems. DO NOT use on seedling asparagus. Use the higher rates on heavier soils.
Bananas	Billygoat weed (Bluetop) Amaranthus, Barley grass, Barnyard grass, Capeweed, Cobbler's pegs, Crowsfoot grass, Fat hen, Guinea grass, Mustard, Pigweed, Prince of Wales feather, Ryegrass, Sowthistle, Summer grass, Wild oats, Wild radish, Wild turnip	Qld only Qld, NSW, WA only	275-500g/ha Plus 1.2 or 1.6L/ha Shirquat@ 250	Add to Shirquat 250 to enhance weed control under favourable growing conditions and in open sunny conditions. Diurex WG at rates up to 500g/ha can be blanket sprayed. Use a directed spray for higher rates of Diurex WG. Complete spray coverage is essential. For grasses and broadleaved weeds up to 5cm high use a minimum of 250L spray solution/ha, increase to 350L/ha for weeds up to 10cm high. Use a minimum spray volume of 400L/ha for weeds > 10cm high and for dense mature weeds. Refer to Shirquat 250 label for rates and full directions. If Shirquat 250 rate is less than the ratio 500mL/100L add 100mL Shirwet 600 or 60mL BS1000 per 100L spray mix.

Crop/Situation	Weeds Controlled	State	Rate	Critical Comments
Cotton	Amaranthus, Barley grass, Barnyard grass, Capeweed, Cobbler's pegs, Crowsfoot grass, Fat hen, Guinea grass, Mustard, Pigweed, Prince of Wales feather, Ryegrass, Sowthistle, Summer grass, Wild oats, Wild radish, Wild turnip	Qld, NSW only	1-2kg/ha	<p>Pre-emergence: Apply at planting or within 7 days of planting and prior to cotton emergence. Seed should be planted at least 4-5cm deep and soil compacted over seeds by a normal planter packer wheel. Use high rate on soils high in clay or organic matter. DO NOT use on light, sandy soils as injury to crop may result. DO NOT make a second pre-emergence application to the same field during the same crop year as injury to the crop may result. Apply in 300-450L water/ha.</p> <p>Post emergence (lay-by): Use the higher rates on heavier soil types high in clay and/or organic matter. DO NOT use on light, sandy soils low in organic matter as injury to the cotton may result. Apply when cotton is at least 30cm high, or immediately after the last cultivation. If emerged annual weeds (15cm or less in height) are present add Shirwet 600 to the spray solution at the rate of 250-500mL/1000L. Apply Diurex WG as a directed spray to cover the area beneath cotton plants and between rows. Avoid spray contact or drift onto cotton plants as injury may result. DO NOT apply more than once per season. DO NOT re-plant treated areas to any crop within one year after last application as injury may result, except cotton, corn or sorghum (grain sorghum only) which maybe planted in the Spring following treatment. Apply in 300-500L water/ha.</p>
Lucerne (established for 1 year or more)	Amsinckia (Yellow burr), Charlock, Dead nettle, Poppies, Sheepweed (White ironweed), Three cornered Jack (Double gee) Seedlings of Capeweed, Spiny emex Suppression of Wild radish, Wild turnip	All states	830g/ha	<p>Apply when lucerne has been grazed. Weeds should be very small (emerging – 4 leaf) when treated. Use only in areas where lucerne becomes Winter dormant.</p> <p>A wider spectrum of weeds will be controlled when tank mixed with Shirquat 250 or Revolver® (Refer to Shirquat 250 or Revolver label for rates and full directions).</p>
Lupins (on light soils only)	Annual ryegrass, Capeweed, Clovers, Double gee, Medics, Wild radish, Wild turnip and suppression only of Dock, Sorrel, Wireweed, Wild oats and Brome grass	WA only	1.1kg/ha	Use through boom spray equipment only. Apply in a minimum of 50L water/ha. Higher volumes of water will give more even application. Must be applied prior to crop emergence. DO NOT use on white or grey sands
			1.1kg/ha plus 1.25L/ha Trifluralin 480	Use tank mix of Diurex plus Trifluralin 480 where Annual ryegrass is present. Apply at pre-sowing stage.

Crop/Situation	Weeds Controlled	State	Rate	Critical Comments
Pulses (Including Chickpeas, Faba beans, Lentils, Narbon beans, Field peas and Vetch)	Capeweed, Double gee, Wild radish, Wild turnip, Toad rush, Crassula, and Erodium	All States	IBS 830g-1.1kg/ha PSPE 550-830g/ha	Incorporated by sowing (IBS) Apply to bare moist soil prior to sowing or at sowing and incorporate by the sowing operation. Use the lower rate on light sandy soils. Post Sowing Pre-Emergent Application. (PSPE) Apply as a post sowing pre-emergent application to moist soil before weed or crop emergence. Use the lower rate on light sandy soils. Application should not be made to excessively ridged or waterlogged soils. Sow the crop at least 5cm deep. Trifluralin or Spinnaker* may be tank mixed at the recommended rates.
Sugar cane (Plant and ratoon crops)	Barnyard grass, Crowsfoot grass, Guinea grass, Summer grass, Barley grass, Ryegrass, Wild oats, Cobbler's pegs, Pigweed, Prince of Wales feather, Amaranthus, Fat hen, Mustard, Sowthistle, Capeweed, Wild radish, Wild turnip	Qld, NSW only	1.9kg/ha	Post-emergence of cane APPLY ONLY as a directed band spray over a maximum of 60% of the crop area DO NOT apply in the Wet Tropics DO NOT apply in the Burdekin between 1 January and 29 February DO NOT apply in Mackay/Whitsunday between 1 December and 30 April DO NOT apply in Mary-Burnett between 1 November and 29 February DO NOT apply in NSW between 1 November and 30 April Soil surface must be moist at time of application. Rain or irrigation is required within 10 days of application if weeds at early post-emergence stage addition of Shirwet 600 at 250mL/100L of water will enhance activity, but direct spray to below cane leaves.
			275-500g/ha plus 1.2-1.6L/ha Shirquat 250	Add to Shirquat 250 to enhance weed control under favourable growing conditions and in open sunny conditions. Diurex WG at rates up to 500g/ha can be blanket sprayed. Complete spray coverage is essential. For grasses and broadleaved weeds up to 5cm high use a minimum of 250L spray solution/ha, increase to 350L/ha for weeds up to 10cm high. Use a minimum spray volume of 400L/ha for weeds > 10cm high and for dense mature weeds. Refer to Shirquat 250 label for rates and full directions. Note: Where broad leaved weeds are a serious problem add Tornado® DF to the mixture.

Crop/Situation	Weeds Controlled	State	Rate	Critical Comments
			500g-1.9kg/ha plus 1.2-1.6L/ha Shirquat 250	<p>APPLY ONLY as a directed band spray over a maximum of 60% of the crop area DO NOT apply in the Wet Tropics DO NOT apply in the Burdekin between 1 January and 29 February DO NOT apply in Mackay/Whitsunday between 1 December and 30 April DO NOT apply in Mary-Burnett between 1 November and 29 February DO NOT apply in NSW between 1 November and 30 April</p> <p>Add to Shirquat 250 to enhance weed control under favourable growing conditions and in open sunny conditions. Use only as a directed spray. Complete spray coverage is essential. For grasses and broadleaved weeds up to 5cm high use a minimum of 250L spray solution/ha, increase to 350L/ha for weeds up to 10cm high. Use a minimum spray volume of 400L/ha for weeds > 10cm high and for dense mature weeds. Refer to Shirquat 250 label for rates and full directions. Note: Where broad leaved weeds are a serious problem add Tornado® DF to the mixture.</p>
Summer fallows	Potato weed (<i>Heliotropium europaeum</i>)	SA only	275g/ha	Only effective when in mixture with Revolver® at 1.6L/ha (See Revolver label)
Wheat, Barley, Oats	Amsinckia, Capeweed, Charlock, Corn gromwell (Ironweed), Dead nettle, Ice plant, Melilotus, Mustard, Rough poppy, Saffron thistle, Sheepweed, Three cornered Jack (Double gee, Spiny emex), Wild radish, Wild turnip, Yellow burr weed	NSW, Vic, ACT & SA only	500g/ha	<p>Apply when weeds are at the 2-4 leaf stage and soon after the crop is past the 2-5 leaf stage which is normally within 6 weeks of sowing.</p> <p>DO NOT SPRAY LATER THAN THE EARLY TILLERING STAGE.</p>
Wheat, Barley, Triticale, Cereal rye, Oats	Capeweed, Double gee, Mustard, Wild radish [suppression only], Water weeds [Toads rush, Crassula]	WA only	275-550g/ha ^Barley to 390g/ha only	<p>Post planting application only for Full cut sowing systems that leave a relatively smooth surface, eg. standard combine or airseeder. Apply post plant pre-emergent to crop within 3-4 days of planting.</p> <p>Pre plant application only for sowing systems (eg No Till or Zero Till) which leave open slots or furrows, or where seed is sown along the bottom of the furrow. Even the soil surface where harrowed after sowing. Apply within 7-10 days pre plant.</p> <p>Soil Types - use lower rates on sandy surfaced soils with a low organic carbon or clay content. Caution: DO NOT use on grey or white sands. Crop to be planted at least 35mm deep. Crop tipping and yellowing may occur particularly when heavy rain falls after sowing. Shallow sown crops are more at risk.</p> <p>Caution: Severe crop damage will occur if herbicides containing Paraquat or Diquat contact an emerging cereal crop either as a salvage treatment or via spray drift unless rainfall has been exceptionally low.</p>
	As above plus suppression of Annual ryegrass, Barley grass and Silver grass		550g/ha plus 500mL of a 720g/L metolachlor to improve suppression of Ryegrass	
Wheat, Barley, Triticale, Cereal rye			275-550g plus 830mL/ha of Trifluralin 480	As Above Must be incorporated within 4 hours of application.
Wheat, Barley,	Capeweed, Double gee,	WA only	200g/ha plus	Apply when the Apply by boom spray.

Crop/Situation	Weeds Controlled	State	Rate	Critical Comments
Triticale, Oats	Erodium, London rocket, Lupin, Mustard, Rapistrum, Storkbill, Wild mustard, Wild radish, Wild turnip,		270mL/ha Thistlekille® 750	crop has at least 3-5 leaves on the main stem. (Z13-15)
			280g/ha plus 200mL/ha Amine 625	Apply when the crop has at least 4-5 leaves on the main stem. (Z14-15)
Wheat, Barley, Triticale, Cereal rye	Amsinckia (Yellow burr), Capeweed, Charlock, Corn gromwell, Dead nettle, Double gee (Three cornered Jack), Erodium spp., Mustard, Spiny emex, Turnip weed, White ironweed, (Sheepweed), Wild radish, Wild turnip	NSW, Vic, ACT, SA only	280g/ha plus 330mL/ha (NSW, Vic only) 230mL/ha (SA only) Thistlekille® 750	Apply by boomspray when the crop is at the 3-5 leaf stage (NSW, SA only) 5 leaf stage (Vic only). Weeds must be in the 2-5 leaf stage. DO NOT spray if rain is imminent. DO NOT apply to under sown medic or lucerne. In NSW DO NOT use on Durati, Kite or Shortim wheat. Apply in 20-100L/ha.
Wheat, Barley, Triticale	Sour sob	SA only	710-980g/ha	Boom Spray: Sow crop at least 5cm deep. Apply when crop is in the 2-5 true leaf stage. Use the lower rate on lighter soils. NSW: Some crop damage may occur at 1.1kg/ha.
		NSW only	1.1kg/ha	
Irrigation Channels and Drainage Ditches	Annual and perennial grasses and broadleaf weeds (Except Nutgrass, Bindweed and Russian knapweed)	SA, WA, Vic, Qld, Tas only	22kg/ha OR 220g/100m ²	High Volume or Power Spraying. Apply in 2000L water. Apply during the non-crop season and when channel is not in use. Soil should be moist and preferably bare ie. after regrading and cleaning channels. Apply prior to expected seasonal rainfall so as to "fix" it into the soil. If 50-100mm of rain has not fallen after application and it is necessary to use the irrigation channel, fill it with water and allow to stand for 72 hours, then drain off the water in the channel and run to waste.
Bore Drains	Prickly acacia, Mimosa bush	Qld only	35.5kg/ha	Block off bore drain 24 hours prior to spraying and allow drain to empty. Spray 1m wide strip onto mud. Wait 48-72 hours before opening drain again. Water MUST by-pass house tanks and desirable tree stands for 7 days after opening drain. DO NOT delve for 2 weeks.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: DO NOT ALLOW ANIMALS TO DRINK WATER FROM TREATED BORE DRAINS FOR 3 DAYS BEFORE SLAUGHTER FOR HUMAN CONSUMPTION.

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.

GRAZING:

PULSE CROPS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 35 DAYS AFTER APPLICATION.

ALL OTHER CROPS: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

Diurex WG Herbicide is a powerful weedkiller which remains active in the soil for a long time. Use only at recommended rates and read label directions carefully before using. The effects of this product on weeds are slow to appear and will not become apparent until it is carried into the root zone of the weeds or weed seeds by moisture. Results vary with soil type - lower rates are effective on lighter soils and higher rates on heavier soils. This product is most effective and economical when applied to moist bare soil or soil carrying a small amount of vegetation. Application should be made shortly before the period of main weed growth and best results will be achieved if

rainfall or irrigation occurs within 3-4 days of application. Where additional "knockdown" is required, Shirwet 600 at the rate of 250-500mL/100L of water should be added. If existing herbage is excessive it is advisable to mix Shirquat 250 with Diurex WG. This product is inactivated by absorption onto charcoal and other carboniferous materials.

RESISTANT WEEDS WARNING

GROUP	C	HERBICIDE
-------	----------	-----------

Diurex WG Herbicide is a member of the urea group of herbicides. Diurex WG has the inhibitor of photosynthesis at photosystem II mode of action. For weed resistance management Diurex WG is a Group C herbicide. Some naturally occurring weed biotypes resistant to Diurex WG and other Group C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Diurex WG or other Group C herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Crop Care Australasia Pty Ltd accepts no liability for any losses that may result from the failure of Diurex WG to control resistant weeds.

MIXING

This product mixes readily with water. Steadily add the required quantity to the spray vat which should be 50-75% full. The agitation system should be running during addition of the product and during the spraying operation. If left to settle, stir/agitate thoroughly before spraying. Avoid drawing accumulated particles into outlet pipe. DO NOT pre-mix as a paste.

Wetting Agent

Add Shirwet 600 at 125mL/100L of spray mixture when applying to emerged weeds as a foliar spray.

COMPATIBILITY

Diurex WG Herbicide is compatible with the following Crop Care products: Shirquat® 250, Spray*Seed* 250, Revolver®, Thistlekill® 750, Amine 625, Trifluralin 480.

Selective use in crops

For selective use in crops it is important to apply recommended rates correctly. Diurex WG Herbicide is most effective when applied to soil which has been well tilled (clod free) and is free of weeds and trash. Where not recommended for overall application care should be taken to avoid spray drift onto fruit or foliage. Wetting agents should only be used where specified.

REPLANTING RESTRICTIONS

Crop Rotations: DO NOT replant treated areas to any crop within two years after last spray unless otherwise stated.

Cotton and Lucerne - DO NOT replant treated areas to any crop within 1 year after last spray. Except cotton, corn or grain sorghum which may be planted in the Spring of the following year.

Sugarcane - treated areas can be replanted to pineapple or sugarcane 1 year after last spray.

CROP SAFETY

Heavy rains after application of this product may cause severe crop damage, risk must be assumed by the user.

Crop Safety:

DO NOT USE ON SANDY OR GRAVELLY SOILS OR SOILS LOW IN ORGANIC MATTER. AVOID CONTACT OF SPRAYS WITH CROP FOLIAGE OR FRUIT.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment which may cause spray drift onto nearby susceptible plants/crops, cropping lands or pastures. DO NOT use for soil sterilisation near or under trees or shrubs.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN. Store in the closed, original container in a dry, cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Shake empty box or bag into spray tank. Single rinse bag before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. Puncture or shred and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500mm in a disposal pit specifically marked and set up for the purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from your supplier, or from the Crop Care website www.cropcare.com.au

Conditions of sale

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes

the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Care Australasia Pty Ltd within thirty days of delivery.

In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 498 ALL HOURS - AUSTRALIA WIDE
---	--

® Shirquat, Diurex, Thistekillem and Revolver are registered trademarks of, or used under licence by Crop Care Australasia Pty Ltd

* Other registered trademarks