


Please Note: This product has been superseded by

David Grays Crabgrass & Clover Killer.



David Grays D.S.M.A. Crab Grass Killer.

Available in 50gm, 4kg and 25kg.

Active Constituent 630g/kg D.S.M.A.

A selective herbicide to control Crab Grass in Lawns, Tennis Courts, Bowling Greens and playing fields.

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTION BEFORE OPENING OR USING

Directions for use:

USE ONLY ON COUCH OR KIKUYU LAWNS.

Water the grass well for a few days prior to application.

Mix 25 grams in 4 litres of water and stir until dissolved.

Apply at the rate of 4 litres of the solution per 30 square metres.

Repeat application at 7 day intervals.

Avoid watering for 4 days after application or until necessary for the survival of the grass or turf.

SAFETY DIRECTIONS: This pesticide is dangerous if swallowed, inhaled or absorbed. Avoid contact with eyes and skin and avoid breathing the vapour. Wash concentrate from eyes and skin immediately. Wash exposed parts of the body after use and before eating, drinking or smoking.

FIRST AID: If poisoning occurs contact a Doctor or Poisons Information Centre Ph: 131126. If swallowed and more than 15 minutes from hospital induce vomiting preferably using IPECAC SYRUP APF.

PASSTOX FLOWABLE: TURF HERBICIDE

Active Ingredient: 526g/L DSMA

Poison Schedule: DANGEROUS POISON S7

Registered for use In: N.S.W., V.I.C., S.A., QLD., ACT, W.A.

Resistance Grouping: Group K Herbicide

Available In: 10 Litre & 1 Litre (10 x 1L per carton)

DIRECTIONS FOR USE

RESTRAINTS:

DO NOT apply to dry turf during hot weather.

DO NOT water or cut grass for 24 hours after treatment.

SITUATION	WEEDS CONTROLLED	STATE	RATE	PERCENTAGE	CRITICAL COMMENTS
			100	h m ² a	

Common Couch Lawns & Turf	Paspalum, Summer Grass, Mullumbimby couch.	NSW, 18	1	Apply as a foliar spray in a minimum 100L water/1000 sq. metres or 1000L/ha. Add Ultrawet wetter at 300mL/100sq. m or 3L/ha
		Vic, SA, 0 Qld, ACT only		

WA only	10 0 m L	1	Apply again 7-10 days later and again 21- 30 days after that.
---------	-------------------	---	--

Bent and Fescue lawns and Turf	Paspalum, Summer Grass, Mullumbimby couch.	NSW, 75	7.	If necessary irrigate prior to treatment but do not water for 24 hours after treatment.
		Vic, QLD, m ACT only		

SA only	90 m L	9
---------	--------------	---

General Instructions:

DO NOT use on Buffalo, Kikuyu, Queensland Blue Couch, Carpet or Centipede grass as injury will occur. In hot weather temporary yellowing may occur. Risk of damage increases with temperatures over 30C. Injury will also occur on Bent or Fescues if the recommended rates are exceeded. Passtox Flowable is most effective on actively growing weeds at air temperatures above 21 degrees celcius. Do NOT treat new lawns until after 3 mowings.

Mixing:

Add the required amount of Passtox Flowable to water and stir thoroughly until dissolved. Then add Ultrawet as required. Do NOT store spray solution in tank for prolonged periods.

RESISTANT WEEDS WARNING:

CAMPBELL PASSTOX FLOWABLE HERBICIDE is a member of the organoarsenic group of herbicides. CAMPBELL PASSTOX FLOWABLE HERBICIDE has the 'herbicides with diverse sites of action' mode of action. For weed resistance management CAMPBELL PASSTOX HERBICIDE is a Group K herbicide.

Some naturally-occurring weed biotypes resistant to CAMPBELL PASSTOX HERBICIDE and other 'diverse sites of action herbicides' may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by CAMPBELL PASSTOX FLOWABLE HERBICIDE or other herbicides with diverse sites of action herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Colin Campbell (Chemicals) Pty Ltd accepts no liability for any losses that may result from the failure of CAMPBELL PASSTOX FLOWABLE HERBICIDE to control resistant weeds.

Environmental Information:

DO NOT feed grass clippings from treated areas to poultry or other animals. Do not allow livestock to graze treated area.

Toxic to fish. Do not contaminate streams, rivers or waterways with the chemical or used container.

Do not apply under weather conditions or from spraying equipment which may cause spray to drift onto nearby susceptible plants, crops, crop lands or pastures.

Storage and Disposal:

Store in the closed original container in a well ventilated area, as cool as possible out of direct sunlight. Store in a locked room or place away from children, animals, food, feed-stuffs, seed and fertilizers. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. Break, crush, puncture and bury empty containers in a local authority landfill. If not available bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

Safety Directions:

