

Main panel label

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Ospray
Olympus
Fungicide



ACTIVE CONSTITUENT: 125 g/L EPOXICONAZOLE

GROUP	3	FUNGICIDE
-------	----------	-----------

For the control of certain fungal diseases including stripe rust of wheat and specified fungal diseases of barley, as per the DIRECTIONS FOR USE table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE.

Contents: (5 Litres, 10 Litres, 20 Litres, 200 Litres)

Ospray Pty Ltd
50 Middleton Road
Cottonvale QLD 4375
Freecall 1800 624 597
Fax 07 4685 2446
www.ospray.com.au

APVMA Approval No.: 62840/57247

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes, nose, throat and skin. Avoid contact with eyes and skin. Do not inhale vapours. If product on skin, immediately wash area with soap and water. Wash hands after use. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and half facepiece respirator. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat. After each day's use, wash gloves, respirator and if rubber, wash with warm water. Wash contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS) available from Ospray Pty Ltd.

CONDITIONS OF SALE

This product must be used strictly as directed, and in accordance with all instructions appearing on the label. Ospray Pty Ltd accepts no liability or responsibility for loss or damage arising from incorrect storage, handling or use.

Batch Number:

Date of Manufacture:

Barcode

Drummuster logo

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Ospray
Olympus
Fungicide

ACTIVE CONSTITUENT: 125 g/L EPOXICONAZOLE

GROUP	3	FUNGICIDE
-------	----------	-----------

For the control of certain fungal diseases including stripe rust of wheat and specified fungal diseases of barley, as per the DIRECTIONS FOR USE table.

IMPORTANT: READ THIS LEAFLET BEFORE USE.

Ospray Pty Ltd
50 Middleton Road
Cottonvale QLD 4375
Freecall 1800 624 597
Fax 07 4685 2446
www.ospray.com.au

APVMA Approval No.: 62840/57247.

Ancillary panel leaflet

DIRECTIONS FOR USE:

CROP	DISEASE	RATE PER HA	CRITICAL COMMENTS
Wheat	Leaf rust (<i>Puccinia recondita</i>)	500 mL	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59) if disease is present. DO NOT apply later than ZGS 59.
	Stripe rust (<i>Puccinia striiformis</i>) Septoria nodorum blotch (<i>Phaeosphaeria nodorum</i>)	250 to 500 mL	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59) if disease is present. DO NOT apply later than ZGS 59. Apply the higher rate when disease is present on the top leaf or conditions are favourable for disease development at these stages of crop development. Repeat spraying may be required, particularly if infection occurs early. Regularly monitor the crop from 3-4 weeks after the first application for signs of re-infection.
	Powdery mildew (<i>Blumeria graminis</i> f.sp. <i>tritici</i>)	250 mL	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply during early tillering (ZGS 21-22) stage of the crop if disease is present. DO NOT apply later than ZGS 59. Repeat spraying may be required, particularly if infection pressure persists. Regularly monitor the crop from 3-4 weeks after the first application for signs of re-infection.

Ancillary panel leaflet continued

CROP	DISEASE	RATE PER HA	CRITICAL COMMENTS
Barley			DO NOT exceed a rate of 500 mL/ha in a growing season.
	Leaf rust (<i>Puccinia hordei</i>)	250 to 500 mL	Apply when conditions favour disease development and preferably prior to development of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59) if disease is present. DO NOT apply later than ZGS 59. Apply the higher rate when disease is present on the top leaf or conditions are favourable for disease development at these stages of crop development.
	Net form of net blotch (<i>Pyrenophora teres</i> f. sp. <i>teres</i>)	250 to 500 mL	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59) if disease is present. DO NOT apply later than ZGS 59. Two applications at the lower rate may provide better control than a single application under certain conditions. If applying a two spray program, only apply 250 mL/ha per application, with the first application at ZGS 31-32 if the disease is present. Aim to apply the second application around ZGS 37-39, but no later than ZGS 59, if infection pressure persists. DO NOT exceed a total rate of 500 mL/ha in barley per growing season.
	Leaf scald (<i>Rynchosporium secalis</i>) Powdery mildew (<i>Blumeria graminis</i> f. sp. <i>hordei</i>)	250 mL	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply between early tillering and flag leaf emergence (ZGS 21-39) if disease is present. DO NOT apply later than ZGS 59. Repeat spraying may be required, particularly if infection pressure persists. Regularly monitor the crop from 3-4 weeks after the first application for signs of re-infection. DO NOT exceed a total rate of 500 mL/ha in barley per growing season.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION.

DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 6 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Ospray Olympus Fungicide is a broad spectrum contact and systemic fungicide. It is taken up via the stem and foliage and is translocated upwards and outwards to provide protection to new growth. Thorough coverage of the crop is necessary for best results.

Check the crop regularly for any symptoms of disease infection. Apply as soon as disease is present as indicated in the Directions for Use.

MIXING

Ospray Olympus Fungicide is a suspension concentrate formulation. The addition of a non-ionic surfactant is generally not required, although under certain environmental conditions and lower total application volumes, the addition of a recommended surfactant may assist in providing better coverage of sprayed surfaces.

Add the product to the half filled spray tank while agitating. Continue to agitate while topping up the tank and during spraying.

APPLICATION

Ground

Apply in a water volume of between 50 and 100 L/ha, using flat fan nozzles operating at around 50 cm above the top of the crop. Use the higher water volume in crops with heavier canopies.

Aerial

Apply with suitable aircraft, set up and operated to apply fungicides to cereal crops in a minimum water volume of 20 L/ha.

COMPATIBILITY

For information on compatibility please contact your local reseller or Ospray representative.

FUNGICIDE RESISTANCE WARNING

GROUP	3	FUNGICIDE
-------	---	-----------

Ospray Olympus Fungicide is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 3 fungicides, thus resulting in a reduction of efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Ospray Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

PRECAUTIONS

Re-entry period: DO NOT allow entry into the treated areas until the spray has dried.

CAUTION

Export of treated cereals

Growers should note that the Maximum Residue Limits (MRLs) or import tolerances do not exist in all markets for cereals treated with Ospray Olympus Fungicide. Additionally, some export markets have established MRLs different to those in Australia. If you are growing cereals for export, please check with Ospray or your grain exporter for the latest information on MRLs and import tolerances BEFORE using Ospray Olympus Fungicide.

LIVESTOCK DESTINED FOR EXPORT MARKETS

The grazing withholding period only applies to stock slaughtered for the domestic market. Some export markets apply different standards. To meet these standards, ensure that in addition to complying with the grazing withholding period, the Export Slaughter interval is observed before stock are sold or exported.

EXPORT SLAUGHTER INTERVAL (ESI):

AFTER OBSERVING THE WITHHOLDING PERIOD FOR GRAZING OR CUTTING FOR STOCKFOOD, LIVESTOCK THAT HAVE BEEN GRAZED ON OR FED TREATED CROPS SHOULD BE PLACED ON CLEAN FEED FOR 3 DAYS PRIOR TO SLAUGHTER.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET CROPS

DO NOT apply under weather conditions, or from spraying equipment, that may cause drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Highly toxic to aquatic organisms. DO NOT apply Ospray Olympus Fungicide aerially if water bodies, watercourses or wetlands are within 30 metres downwind of the application area. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes, nose, throat and skin. Avoid contact with eyes and skin. Do not inhale vapours. If product on skin, immediately wash area with soap and water. Wash hands after use. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and half facepiece respirator. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat. After each day's use, wash gloves, respirator and if rubber, wash with warm water. Wash contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS) available from Ospray Pty Ltd.

CONDITIONS OF SALE

This product must be used strictly as directed, and in accordance with all instructions appearing on the label. Ospray Pty Ltd accepts no liability or responsibility for loss or damage arising from incorrect storage, handling or use.