

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

OVERTURE[®]

FUNGICIDE

ACTIVE CONSTITUENT: 125 g/L EPOXICONAZOLE

GROUP 3 FUNGICIDE

For the control of leaf rust, stripe rust and septoria nodrum blotch of wheat and leaf rust of barley, as specified in the DIRECTIONS FOR USE table.



IMPORTANT:
READ THE ATTACHED BOOKLET BEFORE USE

10L

APVMA APPROVAL NO: 63994/10L/0410

OVERTURE[®]

FUNGICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse the containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes, nose and throat and skin. Avoid contact with eyes and skin. Do not inhale vapours. When opening the container and preparing the spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and half facepiece respirator. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat. If product is on skin, immediately wash area with soap and water. Wash hands after use. After each day's use, wash gloves, respirator and if rubber, wash with warm water. Wash contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Telephone 131 126 Australia-wide.

MATERIAL SAFETY DATA SHEET

If additional hazard information is required, refer to the Material Safety Data Sheet available from the supplier or from our web-site, www.sipcam.com.au

NOTICE TO BUYER

Sipcam Pacific Australia Pty Limited (Sipcam) shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence, use under abnormal conditions or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Sipcam's skill or judgment in purchasing or using the product and every person dealing with this product does so at their own risk.

Date Of Manufacture:

Batch Number:

This product is NOT classified as a Dangerous Good under the Australian code for the transport of dangerous goods by road and rail.

For specialist advice in emergency only, call **1800 033 111** all hours, Australia wide

Sipcam Pacific Australia Pty Ltd

Level 1, 191 Malop Street, Geelong, VICTORIA 3220

© Registered Trademark of Sipcam Pacific Australia Pty Ltd



File Name	63994_47155_leaflet_MPL_V01.pdf
Full-Size Printed Label Dimensions	130 mm wide x 140 mm high_single page 260 mm wide x 140 mm high_spread
Approximate Scale (%) at which the pdf will print on A4 (or A3)	A4 - 100%
Latest Preparation Date	29/04/10

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

OVERTURE[®]
FUNGICIDE

ACTIVE CONSTITUENT: 125 g/L EPOXICONAZOLE

GROUP 3 FUNGICIDE

For the control of leaf rust, stripe rust and septoria nodrum blotch of wheat and leaf rust of barley, as specified in the DIRECTIONS FOR USE table.

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

APVMA Approval No: 63994/0410



SIPCAM PACIFIC AUSTRALIA PTY LTD
Level 1, 191 Malop Street Geelong VICTORIA 3220

GENERAL INSTRUCTIONS

OVERTURE is a broad spectrum contact and systemic fungicide. It is taken up via the stem and foliage and is translocated upwards and outwards to provide protection to new growth. Thorough coverage of the crop is necessary for best results. Check the crop regularly for any symptoms of disease infection. Apply as soon as disease is present on the flag leaf or flag-2. Application beyond flag leaf emergence (ZGS 39) may not provide any yield benefit, particularly under dry seasonal conditions.

MIXING

OVERTURE is a suspension concentrate formulation. Add the product to the half filled spray vat while agitating. Continue to agitate while topping up the tank and during spraying.

APPLICATION

Ground

Apply in a water volume of between 50 and 100L/ha, using flat fan nozzles operating at around 50cm above the top of the crop. Use the higher water volume in crops with heavier canopies.

Aerial

Apply with suitable aircraft, set up and operated to apply fungicides to cereal crops in a minimum water volume of 20L/ha.

COMPATIBILITY

For information on compatibility please contact your local reseller or Sipcam representative.

FUNGICIDE RESISTANCE WARNING

OVERTURE is a member of the DMI group of fungicides.

GROUP 3 FUNGICIDE

For fungicide resistance management the product is a Group 3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Sipcam accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

RE-ENTRY PERIOD

DO NOT allow entry into treated areas until the spray has dried.

DIRECTIONS FOR USE:

CROP	DISEASE	RATE	CRITICAL COMMENTS
Wheat	Leaf rust (<i>Puccinia recondita</i>), Stripe rust (<i>Puccinia striiformis</i>)	500mL plus 200mL Chemwet per 100L of spray mix	Apply when conditions favour disease development and preferably prior to development of high levels of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59). DO NOT apply later than ZGS 59. Repeat spraying may be required for stripe rust. Regularly monitor crops from 3-4 weeks after spraying for signs of reinfection and treat accordingly.
	Septoria nodorum blotch (<i>Septoria nodorum</i>)	500mL plus 200mL Chemwet per 100L of spray mix	
Barley	Leaf rust (<i>Puccinia hordei</i>)	250 to 500mL plus 200L Chemwet per 100L of spray mix	Apply when conditions favour disease development and preferably prior to development of high levels of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59). DO NOT apply later than ZGS 59. Apply the higher rate when disease is present on the top leaf or conditions are favourable for disease development at these stages of crop development.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION.

DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 6 WEEKS AFTER APPLICATION.

EXPORT SLAUGHTER INTERVAL (ESI):

AFTER OBSERVING THE PERIOD FOR GRAZING OR CUTTING FOR STOCKFOOD, LIVESTOCK THAT HAVE BEEN GRAZED ON OR FED TREATED CROPS SHOULD BE PLACED ON CLEAN FEED FOR 3 DAYS PRIOR TO SLAUGHTER.

CAUTION**Export of treated cereals**

Growers should note that Maximum Residue Limits (MRLs) or import tolerances do not exist in all markets for cereals treated with OVERTURE Fungicide. Additionally, some export markets have established MRLs different to those in Australia. If you are growing cereals for export, please check with Sippcam or your grain exporter for the latest information on MRLs and import tolerances BEFORE using OVERTURE Fungicide.

LIVESTOCK DESTINED FOR EXPORT MARKETS

The grazing withholding period only applies to stock slaughtered for the domestic market. Some export markets apply different standards. To meet these standards, ensure that in addition to complying with the grazing withholding period, the Export Slaughter Interval is observed before stock are sold or slaughtered.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT


Highly toxic to aquatic organisms. DO NOT apply under weather conditions or from spraying equipment which could be expected to cause spray drift onto adjacent crops, crop lands, livestock, pastures, wetland, waterbodies or watercourses. DO NOT apply OVERTURE aerially if waterbodies, watercourses or wetlands are within 30 metres downwind of the application area.
DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes, nose, throat and skin. Avoid contact with eyes and skin. Do not inhale vapours. When opening the container and preparing the spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and half facepiece respirator. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or



equivalent clothing) and a washable hat. If product on skin, immediately wash area with soap and water. Wash hands after use. After each day's use, wash gloves, respirator and if rubber, wash with warm water. Wash contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre, telephone 131126 Australia wide.

MSDS

If additional hazard information is required, refer to the Material Safety Data Sheet available from the supplier or from our web-site, www.sipcam.com.au

NOTICE TO BUYER

Sipcam Pacific Australia Pty Ltd (Sipcam) shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence, use under abnormal conditions or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Sipcam's skill or judgement in purchasing or using the product and every person dealing with this product does so at their own risk.

