

Product Name: OVERTURE FUNGICIDE
APVMA Approval No: 63994/112327



Label Name:	OVERTURE FUNGICIDE
Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	125 g/L EPOXICONAZOLE
Mode of Action:	GROUP 3 FUNGICIDE
Statement of Claims:	For the control of certain fungal diseases including stripe rust of wheat, and specified fungal diseases of barley, as specified in the DIRECTIONS FOR USE.
Net Contents:	10L 20L 5L
Restrains:	
Directions for Use:	
Other Limitations:	NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.
Withholding Periods:	DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION. DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 6 WEEKS AFTER APPLICATION.

Trade Advice:	<p>EXPORT SLAUGHTER INTERVAL (ESI): AFTER OBSERVING THE PERIOD FOR GRAZING OR CUTTING FOR STOCKFOOD, LIVESTOCK THAT HAVE SEEN GRAZED ON OR FED TREATED CROPS SHOULD BE PLACED ON CLEAN FEED FOR 3 DAYS PRIOR TO SLAUGHTER.</p> <p>Export of treated cereals Growers should note that Maximum Residue Limits (MRLs) or import tolerances do not exist in all markets for cereals treated with OVERTURE Fungicide. Additionally, some export markets have established MRLs different to those in Australia. If you are growing cereals for export, please check with Sipcam or your grain exporter for the latest information on MRLs and import tolerances BEFORE using OVERTURE Fungicide.</p> <p>LIVESTOCK DESTINED FOR EXPORT MARKETS The grazing withholding period only applies to stock slaughtered for the domestic market. Some export markets apply different standards. To meet these standards, ensure that in addition to complying with the grazing withholding period, the Export Slaughter Interval is observed before stock are sold or slaughtered.</p>
---------------	--

General Instructions:	<p>OVERTURE Fungicide is a broad spectrum contact and systemic fungicide. It is taken up via the stem and foliage and is translocated upwards and outwards to provide protection to new growth. Thorough coverage of the crop is necessary for best results. Check the crop regularly for any symptoms of disease infection. Apply as soon as disease is present on the flag leaf or flag-2. Application beyond flag leaf emergence (ZGS 39) may not provide any yield benefit, particularly under dry seasonal conditions.</p> <p>MIXING OVERTURE Fungicide is a suspension concentrate formulation. The addition of a non-ionic surfactant is generally not required, although under certain environmental conditions and lower total application volumes, the addition of Chemwet 1000 or other recommended surfactant may assist in providing better coverage of sprayed surfaces. Add the product to the half-filled spray vat while agitating. Continue to agitate while topping up the tank and during spraying.</p> <p>APPLICATION Ground Apply in a water volume of between 50 and 100L/ha, using flat fan nozzles operating at around 50 cm above the top of the crop. Use the higher water volume in crops with heavier canopies. Aerial Apply with suitable aircraft, set up and operated to apply fungicides to cereal crops in a minimum water volume of 20L/ha.</p> <p>COMPATIBILITY For information on compatibility please contact your local reseller or Sipcam representative.</p>
-----------------------	--

Resistance Warning:	<p>GROUP 3 FUNGICIDE</p> <p>OVERTURE Fungicide is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Sipcam accepts no liability for any losses that may result from the failure of this product to control resistant fungi.</p>
---------------------	---

Precautions:	Re-Entry Period DO NOT allow entry into treated areas until the spray has dried.
--------------	---

Protections:	PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT Highly toxic to aquatic organisms. DO NOT apply under weather conditions or from spraying equipment which could be expected to cause spray drift onto adjacent crops, crop lands, livestock, pastures, wetland, waterbodies or watercourses. DO NOT apply OVERTURE Fungicide aerially if waterbodies, watercourses or wetlands are within 30 meters downwind of the application area. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.
--------------	--

Storage and Disposal:	Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.
-----------------------	---

Safety Directions:	Harmful if inhaled or swallowed. Will irritate the eyes, nose and throat and skin. Avoid contact with eyes and skin. Do not inhale vapours. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and half facepiece respirator. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat. If product on skin, immediately wash area with soap and water. Wash hands after use. After each day's use, wash gloves, respirator and if rubber, wash with warm water. Wash contaminated clothing.
--------------------	---

First Aid Instructions:	If poisoning occurs contact a doctor or Poisons Information Centre. Phone 131 126 Australia wide.
-------------------------	---

First Aid Warnings:	
---------------------	--

DIRECTIONS FOR USE:

CROP	DISEASE	RATE PER HA	CRITICAL COMMENTS
Wheat	Leaf rust (<i>Puccinia recondita</i>)	500 mL	Apply when conditions favour disease development and preferably prior to development of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59) if disease is present. DO NOT apply later than ZGS 59.
	Stripe rust (<i>Puccinia striiformis</i>), Septoria nodorum blotch (<i>Septoria nodorum</i>)	250 - 500 mL	
	Powdery mildew (<i>Blumeria graminis</i> f.sp. <i>tritici</i>)	250 mL	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply during early tillering (ZGS 21-22) stage of the crop if disease is present. DO NOT apply later than ZGS 59. Repeat spraying may be required, particularly if infection pressure persists. Regularly monitor the crop from 3-4 weeks after the first application for signs of re-infection.
Barley	DO NOT exceed a total rate of 500mL/Ha of Overture in a growing season.		
	Leaf rust (<i>Puccinia hordei</i>)	250 to 500 mL	Apply when conditions favour disease development and preferably prior to development of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59). DO NOT apply later than ZGS 59. Apply the higher rate when disease is present on the top leaf or conditions are favourable for disease development at these stages of crop development.
	Net form of net blotch (<i>Pyrenophora teres</i> f. sp. <i>teres</i>)	250 to 500 mL	Apply when conditions favor disease development and prior to development of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59) if disease is present. DO NOT apply later than ZGS 59. Two applications of Overture Fungicide at the lower rate may provide better control than a single application under certain conditions. If applying a 2-spray program only apply 250 mL/Ha per application, the first application at ZGS 31-32 if the disease is present. Aim to apply the second application around ZGS 37-39 but no later than ZGS 59, if infection pressure persists. DO NOT exceed a total rate of 500 mL/Ha of Overture Fungicide in barley per growing season.
	Leaf scald (<i>Rynchosporium secalis</i>), Powdery mildew (<i>Blumeria graminis</i> f. sp. <i>hordei</i>)	250 mL	Apply when conditions favour disease development and prior to development of disease in the crop. Aim to apply between early tillering and flag leaf emergence (ZGS 21-39) if disease is present. DO NOT apply later than ZGS 59. Repeat spraying may be required, particularly if infection pressure persists. Regularly monitor the crop from 3-4 weeks after the first application for signs of re-infection. DO NOT exceed a total rate of 500 mL/ha of Overture in barley per growing season.