

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

LEGION[®]

Insecticidal Seed Treatment

ACTIVE CONSTITUENT: 500 g/L FIPRONIL

GROUP	2B	INSECTICIDE
--------------	-----------	--------------------

For the control of redlegged earth mite in canola and false wireworm in sorghum and sunflowers, and to protect sorghum and sunflower seedlings from attack by black field earwig as per the DIRECTIONS FOR USE Table.

Contents: 500mL - 200L

Crop Care Australasia Pty Ltd, Unit 17/16 Metroplex Avenue, Murarrie Qld 4172

DIRECTIONS FOR USE

CROP	PEST	STATE	RATE	CRITICAL COMMENTS
Canola	Redlegged earth mite (<i>Halotydeus destructor</i>)	Qld, NSW, Vic, SA & WA only	400mL/100kg seed	Ensure thorough coverage of seed. Add 400mL of LEGION to 600mL of water/100kg seed (total volume 1.0L/100 kg seed). Seedling protection will be satisfactory under conditions of low pest pressure (ie. less than 1000 per m ²). Where greater pest numbers are anticipated or where there are more than 8 mites per plant, alternative or additional treatments will be required.
Sorghum Sunflowers	False wireworm (<i>Pterohelaeus darlingensis</i> and <i>Gonocephalum macleayi</i>) Protection from: Black field earwig (<i>Nala lividipes</i>)	Qld, NSW, Vic WA & NT only	150mL/100kg seed	Thoroughly coat seed using commercial seed coating equipment. Add 150mL of LEGION to 350mL of water/100kg seed (total volume 500mL/100 kg seed).

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

Harvest

Canola, Sorghum, Sunflower: NOT REQUIRED WHEN USED AS DIRECTED.

Grazing

Canola: DO NOT GRAZE PLANTS GROWN FROM TREATED SEED OR CUT FOR STOCK FOOD WITHIN 9 WEEKS OF SOWING.

Sorghum: DO NOT GRAZE PLANTS GROWN FROM TREATED SEED OR CUT FOR STOCK FOOD WITHIN 5 WEEKS OF SOWING.

Sunflower: DO NOT GRAZE PLANTS GROWN FROM TREATED SEED OR CUT FOR STOCK FOOD WITHIN 3 WEEKS OF SOWING.

DO NOT FEED TREATED SEED TO ANIMALS.

GENERAL INSTRUCTIONS

Where pests are in plague numbers, seed treatment alone may not provide adequate protection and additional methods such as grain baiting may need to be employed.

Insecticide Resistance Warning

GROUP	2B	INSECTICIDE
--------------	-----------	--------------------

For insecticide resistance management, LEGION Insecticidal Seed Treatment is a Group 2B insecticide. Some naturally occurring insect biotypes resistant to LEGION and other Group 2B insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if LEGION or other Group 2B insecticides are used repeatedly. The effectiveness of LEGION on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Crop Care Australasia Pty Ltd accepts no liability for any losses that may result from the failure of LEGION to control resistant insects.

LEGION may be subject to specific resistance management strategies. For further information contact your local supplier, Crop Care Australasia Pty Ltd representative or local agricultural department agronomist.

Mixing And Application

Shake container before use.

Canola

LEGION should be applied in a total volume of 1.0L water/100kg seed.

Sorghum and sunflowers

Mix LEGION with water to a total volume of up to 500mL/100kg seed and apply to seed in a manner which ensures thorough and even coverage.

Compatibility

LEGION is compatible with Semevin+, quintozene, Apron⁺ and Peridiam⁺.

PRECAUTIONS

Treated seed should be allowed to dry before re-handling.

DO NOT use treated seeds for human or animal consumption. DO NOT allow treated seed to contaminate grain/other seed intended for animal or human consumption. DO NOT feed treated seed, or otherwise expose, to wild or domestic birds.

PROTECTION OF LIVESTOCK

DO NOT feed treated seed to animals, including poultry.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. Toxic to galliform birds and related species. Highly toxic to fish and very highly toxic to aquatic invertebrates. Any spillage of treated seed resulting during handling and loading operations must be cleaned up as soon as practicable, preferably by recovery and re-use. If disposal is required, ensure treated seeds are thoroughly buried and not accessible to livestock, birds and other wildlife.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Rinse containers before disposal. Add rinsings to the mixing tank. Do not dispose of undiluted chemicals on-site. Dispose of at an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Poisonous if inhaled or swallowed. Avoid contact with skin. When opening the container, preparing product for use and using the prepared product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from your supplier or from the Crop Care website: www.cropcare.com.au

Conditions of Sale

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Care Australasia Pty Ltd within thirty days of delivery.

UN No: 2902	PESTICIDE, LIQUID, TOXIC, N.O.S. (contains fipronil)
In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 498 ALL HOURS – AUSTRALIA WIDE
PG III	HAZCHEM 2X

® Legion is a registered trademark used under licence by Crop Care Australasia Pty Ltd..
+ other registered trademarks.



Crop Care Australasia Pty Ltd
ACN 061 362 347
Unit 17/16 Metroplex Avenue
Murarrie Qld 4172
Tel : (07) 3909 2000

Insert barcodes, label codes and *drumMUSTER* logo as appropriate

APVMA Approval No. : 70278/101505

BN :

DOM :