

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Farmoz

Jubilee®

IN-FURROW & FOLIAR FUNGICIDE

ACTIVE CONSTITUENT: 250 g/L FLUTRIAFOL

GROUP **C** FUNGICIDE

For the control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser or applied as a foliar spray as specified in the Directions for Use table.

www.farmoz.com.au

DIRECTIONS FOR USE: ALL STATES

A. JUBILEE APPLIED IN-FURROW AS A FERTILISER TREATMENT

CROP	DISEASE	RATE PER HECTARE	WHP	CRITICAL COMMENTS																								
Canola	Blackleg (<i>Leptosphaeria maculans</i>)	400 mL	4 weeks Grazing	Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties. Application: Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit 400 mL of the product on the quantity of fertiliser required to be sown in furrow per hectare, using the table below as a guide. <table border="1"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100 kg of fertiliser (mL)</th> </tr> </thead> <tbody> <tr><td>60</td><td>667*</td></tr> <tr><td>70</td><td>571</td></tr> <tr><td>80</td><td>500</td></tr> <tr><td>100</td><td>400</td></tr> <tr><td>120</td><td>333</td></tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)	60	667*	70	571	80	500	100	400	120	333												
Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)																											
60	667*																											
70	571																											
80	500																											
100	400																											
120	333																											
Wheat	Stripe Rust (<i>Puccinia striiformis</i>)	200 or 400 mL		For control of stripe rust of wheat and control of Powdery Mildew and Scald of barley, use the high rate where conditions conducive to severe disease are expected. Application: Spray product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit the specific rate of the product on the quantity of fertiliser required to be sown in furrow per hectare, using the table below as a guide. <table border="1"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th colspan="2">Volume of product per 100 kg of fertiliser (mL)</th> </tr> <tr> <th></th> <th>200 mL rate</th> <th>400 mL rate</th> </tr> </thead> <tbody> <tr><td>50</td><td>400</td><td>800*</td></tr> <tr><td>60</td><td>333</td><td>667*</td></tr> <tr><td>70</td><td>286</td><td>571</td></tr> <tr><td>80</td><td>250[†]</td><td>500</td></tr> <tr><td>100</td><td>200[†]</td><td>400</td></tr> <tr><td>120</td><td>167[†]</td><td>333</td></tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)			200 mL rate	400 mL rate	50	400	800*	60	333	667*	70	286	571	80	250 [†]	500	100	200 [†]	400	120	167 [†]	333
Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)																											
	200 mL rate	400 mL rate																										
50	400	800*																										
60	333	667*																										
70	286	571																										
80	250 [†]	500																										
100	200 [†]	400																										
120	167 [†]	333																										
	Septoria Tritici Blotch (<i>S. tritici</i>), Take-all (<i>Gaeumannomyces graminis</i>)	400 mL																										
Barley	Powdery Mildew (<i>Erysiphe graminis</i>), Scald (<i>Rhynchosporium secalis</i>)	200 or 400 mL		<table border="1"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th colspan="2">Volume of product per 100 kg of fertiliser (mL)</th> </tr> <tr> <th></th> <th>200 mL rate</th> <th>400 mL rate</th> </tr> </thead> <tbody> <tr><td>50</td><td>400</td><td>800*</td></tr> <tr><td>60</td><td>333</td><td>667*</td></tr> <tr><td>70</td><td>286</td><td>571</td></tr> <tr><td>80</td><td>250[†]</td><td>500</td></tr> <tr><td>100</td><td>200[†]</td><td>400</td></tr> <tr><td>120</td><td>167[†]</td><td>333</td></tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)			200 mL rate	400 mL rate	50	400	800*	60	333	667*	70	286	571	80	250 [†]	500	100	200 [†]	400	120	167 [†]	333
Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)																											
	200 mL rate	400 mL rate																										
50	400	800*																										
60	333	667*																										
70	286	571																										
80	250 [†]	500																										
100	200 [†]	400																										
120	167 [†]	333																										

* Application volumes over 600 mL per 100 kg of fertiliser may adversely effect the flow rate and handling characteristics of dusty/poorly granulated fertilisers and some high nitrate fertilisers.

[†] Where application volume is low, product may be diluted with water to improve coverage.

B. JUBILEE APPLIED AS A FOLIAR SPRAY

CROP	DISEASE	RATE PER HECTARE	WHP	CRITICAL COMMENTS
Wheat	Stripe Rust (<i>Puccinia striiformis</i>)	250 - 500 mL Ground Application Apply in 100-150 L water plus 200 mL 'BS 1000' per 100 L of spray solution.	7 weeks Harvest or Grazing	Apply when stripe rust appears and before disease reaches 1% of leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential, and where conditions favourable to severe disease persist, use a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.
	Leaf Rust (<i>P. recondita</i>), Septoria Nodorum Blotch (<i>Septoria nodorum</i>), Septoria Tritici Blotch (<i>S. tritici</i>)	Aerial Application Apply in a minimum of 15 L water plus 200 mL 'BS 1000' per 100 L of spray solution.		Apply after flag leaves have begun to emerge and before disease reaches 1% of flag leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential, and where conditions favourable to severe diseases persist during flowering, use a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.
Barley	Powdery Mildew (<i>Erysiphe graminis</i>)		10 weeks Harvest or Grazing	Apply when powdery mildew first appears. The high rate of application gives a longer period of protection than the low rate. Where conditions favourable to severe disease persist, a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

A. JUBILEE APPLIED IN-FURROW AS A FERTILISER TREATMENT

CANOLA, WHEAT AND BARLEY: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING.

B. JUBILEE APPLIED AS A FOLIAR SPRAY

HARVEST

BARLEY: DO NOT HARVEST FOR 10 WEEKS AFTER APPLICATION.

WHEAT: DO NOT HARVEST FOR 7 WEEKS AFTER APPLICATION.

GRAZING

BARLEY: DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 10 WEEKS AFTER APPLICATION.

WHEAT: DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 7 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Fertiliser Application

Ensure that treated fertiliser is dry before sowing with the crop.

Sowing: Treated fertiliser should be sown in the same furrow as the seed, not broadcast. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered. DO NOT mix seed with treated fertiliser.

Foliar Application

Check the crop carefully and regularly. Spray only if there is sufficient time in the season for the rust to develop to a level on the variety in question when it will cause significant yield loss.

FUNGICIDE RESISTANCE WARNING

FARMOZ JUBILEE® In-Furrow & Foliar Fungicide is a member of the DMI group of **GROUP C FUNGICIDE** fungicides. For fungicide resistance management JUBILEE is a Group C fungicide. Some naturally-occurring individual fungi resistant to JUBILEE and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by JUBILEE and other Group C fungicides thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant individuals is difficult to detect prior to use, FARMOZ Pty. Ltd. accepts no liability for any losses that may result from the failure of JUBILEE to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the time of application.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL (5 L, 10 L, 20 L, 200 L)

Store in the closed original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If not available bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

For refillable containers, empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

1000 L

Store in the closed, original container in a cool, well-ventilated areas. DO NOT store for prolonged periods in direct sunlight. Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs FARMOZ Pty. Ltd. should be advised immediately. This minibulk container is reusable and remains the property of FARMOZ Pty. Ltd. DO NOT rinse empty container. Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return to FARMOZ Pty. Ltd. for cleaning, relabelling and refilling.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. May irritate the nose and throat and skin. Avoid contact with eyes and skin. Avoid inhaling spray mist. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and face shield and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for JUBILEE is available from FARMOZ on request. Call Customer Service on (02) 9431 7800.

CONDITIONS OF SALE: The use of FARMOZ JUBILEE In-Furrow & Foliar Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by FARMOZ Pty. Limited regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and FARMOZ Pty. Limited accepts no responsibility for any consequence whatsoever resulting from the use of this product.

© Registered trademark of FARMOZ Pty. Ltd.

NOT A DANGEROUS GOOD ACCORDING TO THE AUSTRALIAN DANGEROUS GOODS (ADG) CODE.

PACK SIZE: 5 L, 10 L, 20 L, 200 L, 1000 L