

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Farmoz

Jubilee®

IN-FURROW & FOLIAR FUNGICIDE

ACTIVE CONSTITUENT: 250 g/L FLUTRIAFOL

GROUP **C** FUNGICIDE

For control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser or applied as a foliar spray as per the Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING

www.farmoz.com.au



Farmoz Pty. Ltd. ABN 55 050 328 973
Suite 1, Level 4, Building B, 207 Pacific Highway
St Leonards NSW 2065 Australia
Tel: (02) 9431 7800 Fax: (02) 9431 7700

CONTENTS

110 L



STORAGE AND DISPOSAL

Store the original sealed Envirodrum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the Envirodrum to the point of purchase. The Envirodrum remains the property of FARMOZ Pty. Ltd.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. May irritate the nose and throat and skin. Avoid contact with eyes and skin. Avoid inhaling spray mist. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and face shield and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

PLACE LEAFLET HERE

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for JUBILEE is available from FARMOZ on request. Call Customer Service on (02) 9431 7800.

CONDITIONS OF SALE: The use of FARMOZ JUBILEE In-Furrow & Foliar Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by FARMOZ Pty. Limited regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and FARMOZ Pty. Limited accepts no responsibility for any consequence whatsoever resulting from the use of this product.

© Registered trademark of FARMOZ Pty. Ltd.



**FOR SPECIALIST ADVICE
IN AN EMERGENCY ONLY
PHONE 1800 024 973
TOLL FREE – ALL HOURS – AUSTRALIA WIDE**



9 3265 12 013927

Batch No.

Date of
Manufacture

APVMA Approval No: 55454/110L/1008

EXECUTIVE PRESS 09/08 – 0418 263 830

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Farmoz

Jubilee[®]

IN-FURROW & FOLIAR FUNGICIDE

ACTIVE CONSTITUENT: 250 g/L FLUTRIAFOL

GROUP C FUNGICIDE

For control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser or applied as a foliar spray as per the Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING

www.farmoz.com.au



Farmoz Pty. Ltd. ABN 55 050 328 973
Suite 1, Level 4, Building B, 207 Pacific Highway
St Leonards NSW 2065 Australia
Tel: (02) 9431 7800 Fax: (02) 9431 7700

CONTENTS

110 L





STORAGE AND DISPOSAL

Store the original sealed Envirodrum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the Envirodrum to the point of purchase. The Envirodrum remains the property of FARMOZ Pty. Ltd.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. May irritate the nose and throat and skin. Avoid contact with eyes and skin. Avoid inhaling spray mist. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and face shield and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for JUBILEE is available from FARMOZ on request. Call Customer Service on (02) 9431 7800.

CONDITIONS OF SALE: The use of FARMOZ JUBILEE In-Furrow & Foliar Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by FARMOZ Pty. Limited regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and FARMOZ Pty. Limited accepts no responsibility for any consequence whatsoever resulting from the use of this product.

® Registered trademark of FARMOZ Pty. Ltd.



PLACE LEAFLET HERE

**FOR SPECIALIST ADVICE
IN AN EMERGENCY ONLY
PHONE 1800 024 973**
TOLL FREE – ALL HOURS – AUSTRALIA WIDE



Batch No.

Date of
Manufacture

APVMA Approval No: 55454/110/1008
EXECUTIVE PRESS 09/08 – 0418 263 830

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Farmoz
Jubilee®

IN-FURROW & FOLIAR FUNGICIDE

ACTIVE CONSTITUENT: 250 g/L FLUTRIAFOL

GROUP C FUNGICIDE

For control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser or applied as a foliar spray as per the Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING

www.farmoz.com.au



Farmoz Pty. Ltd. ABN 55 050 328 973
 Suite 1, Level 4, Building B, 207 Pacific Highway
 St Leonards NSW 2065 Australia
 Tel: (02) 9431 7800 Fax: (02) 9431 7700

CONTENTS

1000 L



STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated areas. **DO NOT** store for prolonged periods in direct sunlight. Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs FARMOZ Pty. Ltd. should be advised immediately. This minibulk container is reusable and remains the property of FARMOZ Pty. Ltd. **DO NOT** rinse empty container. Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return to FARMOZ Pty. Ltd. for cleaning, relabelling and refilling.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. May irritate the nose and throat and skin. Avoid contact with eyes and skin. Avoid inhaling spray mist. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and face shield and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

PLACE LEAFLET HERE

Label image proudly supplied by



MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for JUBILEE is available from FARMOZ on request. Call Customer Service on (02) 9431 7800.

CONDITIONS OF SALE: The use of FARMOZ JUBILEE In-Furrow & Foliar Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by FARMOZ Pty. Limited regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and FARMOZ Pty. Limited accepts no responsibility for any consequence whatsoever resulting from the use of this product.

® Registered trademark of FARMOZ Pty. Ltd.

**FOR SPECIALIST ADVICE IN
 AN EMERGENCY ONLY
 PHONE 1800 024 973
 TOLL FREE – ALL HOURS – AUSTRALIA WIDE**



9 3265 12 011855

Batch No.

Date of
 Manufacture

APVMA Approval No: 55454/1000/0505

EXECUTIVE PRESS 05/05 – 0418 263 830

DIRECTIONS FOR USE: ALL STATES

A. JUBILEE APPLIED IN-FURROW AS A FERTILISER TREATMENT

CROP	DISEASE	RATE PER HECTARE	WHP	CRITICAL COMMENTS														
Canola	Blackleg (<i>Leptosphaeria maculans</i>)	400 mL	4 weeks Grazing	Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties. Application: Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit 400 mL of the product on the quantity of fertiliser required to be sown in furrow per hectare, using the table below as a guide. <table border="1"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100 kg of fertiliser (mL)</th> </tr> </thead> <tbody> <tr><td>60</td><td>667*</td></tr> <tr><td>70</td><td>571</td></tr> <tr><td>80</td><td>500</td></tr> <tr><td>100</td><td>400</td></tr> <tr><td>120</td><td>333</td></tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)	60	667*	70	571	80	500	100	400	120	333		
Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)																	
60	667*																	
70	571																	
80	500																	
100	400																	
120	333																	
Wheat	Stripe Rust (<i>Puccinia striiformis</i>)	200 or 400 mL		For control of stripe rust of wheat and control of Powdery Mildew and Scald of barley, use the high rate where conditions conducive to severe disease are expected. Application: Spray product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit the specific rate of the product on the quantity of fertiliser required to be sown in furrow per hectare, using the table below as a guide. <table border="1"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100 kg of fertiliser (mL)</th> </tr> </thead> <tbody> <tr><td>60</td><td>667*</td></tr> <tr><td>70</td><td>571</td></tr> <tr><td>80</td><td>500</td></tr> <tr><td>100</td><td>400</td></tr> <tr><td>120</td><td>333</td></tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)	60	667*	70	571	80	500	100	400	120	333		
	Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)																
60	667*																	
70	571																	
80	500																	
100	400																	
120	333																	
	Septoria Tritic Blotch (<i>S. tritici</i>), Take-all (<i>Gaeumannomyces graminis</i>)	400 mL																
Barley	Powdery Mildew (<i>Erysiphe graminis</i>), Scald (<i>Rhynchosporium secalis</i>)	200 or 400 mL		<table border="1"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100 kg of fertiliser (mL)</th> </tr> </thead> <tbody> <tr><td>50</td><td>800*</td></tr> <tr><td>60</td><td>667*</td></tr> <tr><td>70</td><td>571</td></tr> <tr><td>80</td><td>500</td></tr> <tr><td>100</td><td>400</td></tr> <tr><td>120</td><td>333</td></tr> </tbody> </table> <p>* Application volumes over 600 mL per 100 kg of fertiliser may adversely effect the flow rate and handling characteristics of dusty/poorly granulated fertilisers and some high nitrate fertilisers. † Where application volume is low, product may be diluted with water to improve coverage.</p>	Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)	50	800*	60	667*	70	571	80	500	100	400	120	333
Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)																	
50	800*																	
60	667*																	
70	571																	
80	500																	
100	400																	
120	333																	

B. JUBILEE APPLIED AS A FOLIAR SPRAY

CROP	DISEASE	RATE PER HECTARE	WHP	CRITICAL COMMENTS
Wheat	Stripe Rust (<i>Puccinia striiformis</i>)	250 - 500 mL Ground Application Apply in 100-150 L water plus 200 mL 'BS 1000' per 100 L of spray solution. Aerial Application Apply in a minimum of 15 L water plus 200 mL 'BS 1000' per 100 L of spray solution.	7 weeks Harvest or Grazing	Apply when stripe rust appears and before disease reaches 1% of leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential, and where conditions favourable to severe disease persist, use a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.
	Leaf Rust (<i>P. recondita</i>), Septoria Nodorum Blotch (<i>Septoria nodorum</i>), Septoria Tritic Blotch (<i>S. tritici</i>)			Apply after flag leaves have begun to emerge and before disease reaches 1% of flag leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential, and where conditions favourable to severe diseases persist during flowering, use a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.
Barley	Powdery Mildew (<i>Erysiphe graminis</i>)		10 weeks Harvest or Grazing	Apply when powdery mildew first appears. The high rate of application gives a longer period of protection than the low rate. Where conditions favourable to severe disease persist, a single application at the higher rate, or two applications 3-4 weeks apart at the low rate.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

A. JUBILEE APPLIED IN-FURROW AS A FERTILISER TREATMENT CANOLA, WHEAT AND BARLEY: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING.

B. JUBILEE APPLIED AS A FOLIAR SPRAY

HARVEST

BARLEY: DO NOT HARVEST FOR 10 WEEKS AFTER APPLICATION.

WHEAT: DO NOT HARVEST FOR 7 WEEKS AFTER APPLICATION.

GRAZING

BARLEY: DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 10 WEEKS AFTER APPLICATION.

WHEAT: DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 7 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Fertiliser Application

Ensure that treated fertiliser is dry before sowing with the crop.

Sowing: Treated fertiliser should be sown in the same furrow as the seed, not broadcast. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered. DO NOT mix seed with treated fertiliser.

Foliar Application

Check the crop carefully and regularly. Spray only if there is sufficient time in the season for the rust to develop to a level on the variety in question when it will cause significant yield loss.

FUNGICIDE RESISTANCE WARNING

FARMOZ JUBILEE® In-Furrow & Foliar Fungicide is a member of the DMI group of fungicides. For fungicide resistance management JUBILEE is a Group C fungicide. Some naturally-occurring individual fungi resistant to JUBILEE and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by JUBILEE and other Group C fungicides thus resulting in a reduction in efficacy and and possible yield loss. Since the occurrence of resistant individuals is



1000 L

Store in the closed, original container in a cool, well-ventilated areas. DO NOT store for prolonged periods in direct sunlight. Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs FARMOZ Pty. Ltd. should be advised immediately. This minibulk container is reusable and remains the property of FARMOZ Pty. Ltd. DO NOT rinse empty container. Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return to FARMOZ Pty. Ltd. for cleaning, relabelling and refilling.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. May irritate the nose and throat and skin. Avoid contact with eyes and skin. Avoid inhaling spray mist. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and face shield and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for JUBILEE is available from FARMOZ on request. Call Customer Service on (02) 9431 7800.

CONDITIONS OF SALE: The use of FARMOZ JUBILEE In-Furrow & Foliar Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by FARMOZ Pty. Limited regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and FARMOZ Pty. Limited accepts no responsibility for any consequence whatsoever resulting from the use of this product.

© Registered trademark of FARMOZ Pty. Ltd.

APVMA Approval No.: 55454/1008
EXECUTIVE PRESS 09/08 - 0418 263 830

difficult to detect prior to use, FARMOZ Pty. Ltd. accepts no liability for any losses that may result from the failure of JUBILEE to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the time of application.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL (5 L, 10 L, 20 L, 200 L)

Store in the closed original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If not available bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

For refillable containers, empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Envirodrum Micro Matic Valve (110 L)

Store the original sealed Envirodrum in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the Envirodrum to the point of purchase. The Envirodrum remains the property of FARMOZ Pty. Ltd.



POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



ACTIVE CONSTITUENT: 250 g/L FLUTRIAFOL



For the control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser or applied as a foliar spray as specified in the Directions for Use table.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT

www.farmoz.com.au



Farmoz Pty. Ltd. ABN 55 050 328 973
Suite 1, Level 4, Building B, 207 Pacific Highway
St Leonards NSW 2065 Australia
Tel: (02) 9431 7800 Fax: (02) 9431 7700