

POISON**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

FLUTRIAFOL 250 SC Fungicide

ACTIVE CONSTITUENT : 250g/L FLUTRIAFOL**GROUP C FUNGICIDE****IN-FURROW AND FOLIAR FUNGICIDE****For control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser or applied as a foliar spray as per the directions for use table.****SHAKE WELL BEFORE USE****READ THE ATTACHED INFORMATION THOROUGHLY BEFORE USING THIS PRODUCT****10 Litres****4FARMERS PTY. LTD.**

A.C.N. 067 443 485

643 MURRAY STREET, WEST PERTH,
WESTERN AUSTRALIA 6005

PH (08) 9356 3445 FAX: (08) 9356 3447

FOR DIRECTION OF USE SEE ATTACHED LEAFLET**GENERAL INSTRUCTIONS:****FERTILISER APPLICATION**

Ensure that treated fertiliser is dry before sowing with the crop.

Sowing: Treated fertiliser should not be broadcast nor be sown in contact with the seed, but in close proximity to the seed. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered. Do not mix seed with treated fertiliser.**FOLIAR APPLICATION**

Check the crop carefully and regularly. Spray only if there is sufficient time in the season for the rust to develop to a level on the variety in question when it will cause significant yield loss.

FUNGICIDE RESISTANCE WARNING **GROUP C FUNGICIDE**

4Farmers FLUTRIAFOL 250 SC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management 4Farmers FLUTRIAFOL 250 SC Fungicide is a Group C fungicide.

Some naturally occurring individual fungi resistant to 4Farmers FLUTRIAFOL 250 SC Fungicide and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by 4Farmers FLUTRIAFOL 250 SC Fungicide and other Group C fungicides thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use 4Farmers Pty Ltd accepts no liability for any losses that may result from the failure of 4Farmers FLUTRIAFOL 250 SC Fungicide to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL (10, 20 & 200L)

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

STORAGE AND DISPOSAL (100 & 1000L)

DO NOT store or handle near heat or naked flame. Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Store this tightly closed minibulk container in a safe and secure place away from children, animals, food and feedstuffs. Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs 4Farmers Pty Ltd should be advised immediately.

This minibulk container is reusable and remains the property of 4Farmers Pty Ltd.

DO NOT rinse empty container. Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return to 4Farmers Pty Ltd for cleaning, relabelling and refilling.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. May irritate the nose and throat and skin. Avoid contact with eyes and skin. Avoid inhaling spray mist. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEETIf additional hazard information is required refer to the Material Safety Data Sheet. For a copy phone 08 9356 3445 or visit our website at www.4farmers.com.au**Condition of sale**

The use of 4Farmers FLUTRIAFOL 250 SC Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by 4Farmers Pty Ltd regarding its suitability, fitness or efficacy for any purpose for which it is used by the buyer, whether in accordance with the directions or not and 4Farmers Pty Ltd accepts no responsibility for any consequences whatsoever resulting from the use of this product.

**IN A TRANSPORT EMERGENCY DIAL 000
POLICE OR FIRE BRIGADE****Not classified as Dangerous Goods by the criteria of the Australian Dangerous Goods Code (ADG Code) for Transport by road or rail.**

Batch Number:

Date of Manufacture:

APVMA Approval No. : 58328/10/0604

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



FLUTRIAFOL 250

SC FUNGICIDE

ACTIVE CONSTITUENT : 250g/L FLUTRIAFOL

GROUP C FUNGICIDE

IN-FURROW AND FOLIAR FUNGICIDE

For control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser or applied as a foliar spray as per the directions for use table.

SHAKE WELL BEFORE USE

4FARMERS PTY. LTD.

A.C.N. 067 443 485
643 MURRAY STREET, WEST PERTH,
WESTERN AUSTRALIA 6005
PH (08) 9356 3445 FAX: (08) 9356 3447

APVMA Approval No.: 58328/0604

GENERAL INSTRUCTIONS

Fertiliser Application

Ensure that treated fertiliser is dry before sowing with the crop.

Sowing: Treated fertiliser should not be broadcast nor be sown in contact with the seed, but in close proximity to the seed. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered. Do not mix seed with treated fertiliser.

Foliar Application

Check the crop carefully and regularly. Spray only if there is sufficient time in the season for the rust to develop to a level on the variety in question when it will cause significant yield loss.

FUNGICIDE RESISTANCE WARNING

GROUP C FUNGICIDE

4Farmers FLUTRIAFOL 250 SC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management 4Farmers FLUTRIAFOL 250 SC Fungicide is a Group C fungicide.

Some naturally occurring individual fungi resistant to 4Farmers FLUTRIAFOL 250 SC Fungicide and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by 4Farmers FLUTRIAFOL 250 SC Fungicide and other Group C fungicides thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use 4Farmers Pty Ltd accepts no liability for any losses that may result from the failure of 4Farmers FLUTRIAFOL 250 SC Fungicide to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL (10, 20 & 200L)

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

STORAGE AND DISPOSAL (100 & 1000L)

DO NOT store or handle near heat or naked flame. Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Store this tightly closed minibulk container in a safe and secure place away from children, animals, food and feedstuffs. Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs 4Farmers Pty Ltd should be advised immediately. This minibulk container is reusable and remains the property of 4Farmers Pty Ltd. DO NOT rinse empty container. Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return to 4Farmers Pty Ltd for cleaning, relabelling and refilling.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. May irritate the nose and throat and skin. Avoid contact with eyes and skin. Avoid inhaling spray mist. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

If additional hazard information is required refer to the Material Safety Data Sheet. For a copy phone 08 9356 3445 or visit our website at www.4farmers.com.au

Condition of sale

The use of 4Farmers FLUTRIAFOL 250 SC Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by 4Farmers Pty Ltd regarding its suitability, fitness or efficacy for any purpose for which it is used by the buyer, whether in accordance with the directions or not and 4Farmers Pty Ltd accepts no responsibility for any consequences whatsoever resulting from the use of this product.

DIRECTIONS FOR USE : ALL STATES

A. FLUTRIAFOL 250 SC APPLIED IN-FURROW AS A FERTILISER TREATMENT

Crop	Disease	Rate per hectare	WHP	Critical Comments																							
Canola	Blackleg (<i>Leptosphaeria maculans</i>)	400mL	4 weeks Grazing	Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties. Application: Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit 400 mL of the product on the quantity of fertiliser required to be sown in furrow per hectare, using the table below as a guide.																							
	<table border="1"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100kg (mL) of fertiliser</th> </tr> </thead> <tbody> <tr><td>60</td><td>667*</td></tr> <tr><td>70</td><td>571</td></tr> <tr><td>80</td><td>500</td></tr> <tr><td>100</td><td>400</td></tr> <tr><td>120</td><td>333</td></tr> </tbody> </table>		Fertiliser rate per hectare (kg)	Volume of product per 100kg (mL) of fertiliser	60	667*	70	571	80	500	100	400	120	333													
Fertiliser rate per hectare (kg)	Volume of product per 100kg (mL) of fertiliser																										
60	667*																										
70	571																										
80	500																										
100	400																										
120	333																										
Wheat	Stripe rust (<i>Puccinia striiformis</i>)	200 or 400mL		For control of stripe rust of wheat and control of powdery mildew and scald of barley, use the high rate where conditions conducive to severe disease are expected.																							
	Septoria tritici blotch (<i>S. tritici</i>) Take-all (<i>Geaumannomyces graminis</i>)	400mL		Application: Spray product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit the specific rate of the product on the quantity of fertiliser required to be sown in-furrow per hectare, using the table below as a guide.																							
Barley	Powdery mildew (<i>Erysiphe graminis</i>) Scald (<i>Rhynchosporium secalis</i>)	200 or 400mL		<table border="1"> <thead> <tr> <th rowspan="2">Fertiliser rate per hectare (kg)</th> <th colspan="2">Volume of product per 100kg (mL) of fertiliser</th> </tr> <tr> <th>200mL rate</th> <th>400mL rate</th> </tr> </thead> <tbody> <tr><td>50</td><td>400</td><td>800*</td></tr> <tr><td>60</td><td>333</td><td>667*</td></tr> <tr><td>70</td><td>286</td><td>571</td></tr> <tr><td>80</td><td>250†</td><td>500</td></tr> <tr><td>100</td><td>200†</td><td>400</td></tr> <tr><td>120</td><td>167†</td><td>333</td></tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100kg (mL) of fertiliser		200mL rate	400mL rate	50	400	800*	60	333	667*	70	286	571	80	250†	500	100	200†	400	120	167†	333
	Fertiliser rate per hectare (kg)	Volume of product per 100kg (mL) of fertiliser																									
200mL rate		400mL rate																									
50	400	800*																									
60	333	667*																									
70	286	571																									
80	250†	500																									
100	200†	400																									
120	167†	333																									
				* Application volumes over 600mL per 100kg of fertiliser may adversely effect the flow rate and handling characteristics of dusty/poorly granulated fertilisers and some high nitrate fertilisers. † Where application volume is low, product may be diluted with water to improve coverage.																							

B. FLUTRIAFOL 250 SC APPLIED AS A FOLIAR SPRAY

Crop	Disease	Rate per hectare	WHP	Critical Comments
Wheat	Stripe rust (<i>Puccinia striiformis</i>)	250mL - 500mL	7 weeks Harvest or Grazing	Apply when stripe rust appears and before disease reaches 1% of leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential, and where conditions favourable to severe disease persist, use a single application at the higher rate, or two applications 3 - 4 weeks apart at the low rate.
	Leaf rust (<i>P. recondita</i>) Septoria nodorum blotch (<i>Septoria nodorum</i>) Septoria tritici blotch (<i>S. tritici</i>)	<p>Ground Application</p> <p>Apply in 100-150L water plus 200mL 'Wetter 1000' per 100L of spray solution.</p> <p>Aerial Application</p> <p>Apply in a minimum of 15L water plus 200mL 'Wetter 1000'</p>		Apply after flag leaves have begun to emerge and before disease reaches 1% of flag leaf area. The high rate of application gives a longer period of protection than the low rate. In crops with high yield potential, and where conditions favourable to severe disease persist during flowering, use a single application at the higher rate, or two applications 3 - 4 weeks apart at the low rate.
Barley	Powdery mildew (<i>Erysiphe graminis</i>)	per 100L of spray solution.	10 weeks Harvest or Grazing	Apply when powdery mildew first appears. The high rate of application gives a longer period of protection than the low rate. Where conditions favourable to severe disease persist, a single application at the higher rate, or two applications 3 - 4 weeks apart at the low rate, may be required.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

A. APPLIED IN-FURROW AS A FERTILISER TREATMENT :

CANOLA, WHEAT AND BARLEY

- DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING.

B. APPLIED AS A FOLIAR SPRAY :

DO NOT HARVEST EARLIER THAN THE NUMBER OF WEEKS SHOWN AFTER APPLICATION:

BARLEY - 10 WEEKS

WHEAT - 7 WEEKS

DO NOT GRAZE OR CUT FOR STOCK FOOD EARLIER THAN THE NUMBER OF WEEKS SHOWN AFTER APPLICATION:

BARLEY - 10 WEEKS

WHEAT - 7 WEEKS

IN A TRANSPORT EMERGENCY

DIAL 000

POLICE OR FIRE BRIGADE

Not classified as Dangerous Goods by the criteria of the Australian Dangerous Goods Code (ADG Code) for Transport by road or rail.

FOR 24 HOUR SPECIALIST ADVICE IN EMERGENCY ONLY PHONE 1800 638 556