

RECEIVED

By Elabels Jclareance at 8:32 am, Sep 17, 2010

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FARMALINX

Flufol 500 SC

FUNGICIDE

ACTIVE CONSTITUENT: 500g/L FLUTRIAFOL

GROUP **C** FUNGICIDE

*For control of certain fungal diseases on wheat,
barley and canola when mixed with fertiliser as per
the Directions for Use table*

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING

farmalinx

FARMALINX Pty Ltd ABN 95 134 353 245
South Sydney Corporate Park, Unit W2E, 75-85 O'Riordan Street, Alexandria NSW 2015
T: 02 8338 9473 F: 02 9313 4352 W: www.farmalinx.com

CONTENTS
**10, 110,
200, 1000L**

FARMALINX

Flufol 500 SC

FUNGICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will irritate the eyes. May irritate the skin. Avoid contact with the eyes and skin. When using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow length chemical resistant gloves and a face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. Additional information is listed in the Material Safety Data Sheet.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX Flufol 500 SC Fungicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 8338 9473.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

APVMA Approval No. 65535/51148

	IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE
	SPECIALIST ADVICE IN AN EMERGENCY ONLY 1800 024 973 ALL HOURS – AUSTRALIA WIDE

Batch No.

BARCODE

Date of Manufacture

farmalinx

FARMALINX

Flufol 500 SC

FUNGICIDE

DIRECTIONS FOR USE: ALL STATES

CROP	DISEASE	Rate/ha	WHP	Critical Comments																								
Canola	Blackleg (<i>Leptosphaeria maculans</i>)	200mL	4 weeks Grazing	Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties. Application: Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit 200mL of the product on the quantity of fertiliser required to be sown in furrow per hectare, using the table below as a guide. <table border="1"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100kg of fertiliser (mL)</th> </tr> </thead> <tbody> <tr><td>60</td><td>333*</td></tr> <tr><td>70</td><td>285</td></tr> <tr><td>80</td><td>250</td></tr> <tr><td>100</td><td>200</td></tr> <tr><td>120</td><td>166</td></tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)	60	333*	70	285	80	250	100	200	120	166												
Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)																											
60	333*																											
70	285																											
80	250																											
100	200																											
120	166																											
Wheat	Stripe rust (<i>Puccinia striiformis</i>) Take -all (<i>Gaeumannomyces graminis</i>)	100 or 200mL		For control of stripe rust of wheat and control of powdery mildew and scald of barley, use the high rate where conditions conducive to severe disease are expected. Application: Spray product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit the specific rate of the product on the quantity of fertiliser required to be sown in- furrow per hectare, using the table below as a guide. <table border="1"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th colspan="2">Volume of product per 100kg of fertiliser (mL)</th> </tr> <tr> <td></td> <th>100mL rate</th> <th>200mL rate</th> </tr> </thead> <tbody> <tr><td>50</td><td>200</td><td>400*</td></tr> <tr><td>60</td><td>166</td><td>333*</td></tr> <tr><td>70</td><td>143</td><td>285</td></tr> <tr><td>80</td><td>125†</td><td>250</td></tr> <tr><td>100</td><td>100†</td><td>200</td></tr> <tr><td>120</td><td>88†</td><td>166</td></tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)			100mL rate	200mL rate	50	200	400*	60	166	333*	70	143	285	80	125†	250	100	100†	200	120	88†	166
	Fertiliser rate per hectare (kg)	Volume of product per 100kg of fertiliser (mL)																										
	100mL rate	200mL rate																										
50	200	400*																										
60	166	333*																										
70	143	285																										
80	125†	250																										
100	100†	200																										
120	88†	166																										
	Septoria tritici blotch (<i>S. tritici</i>)	200mL																										
Barley	Powdery mildew (<i>Erysiphe graminis</i>) Scald (<i>Rhynchosporium secalis</i>)	100 or 200mL		* Application volumes over 300mL per 100kg of fertiliser may adversely affect the flow rate and handling characteristics of dusty/poorly granulated fertilisers and some high nitrate fertilisers. † Where application volume is low, product may be diluted with water to improve coverage.																								

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

APPLIED IN-FURROW AS A FERTILISER TREATMENT

CANOLA, WHEAT AND BARLEY - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING

GENERAL INSTRUCTIONS

Fertiliser Application

Ensure that treated fertiliser is dry before sowing with the crop. Sowing: Treated fertiliser should be sown in the same furrow as the seed, not broadcast. Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered. Do not mix seed with treated fertiliser.

FUNGICIDE RESISTANCE WARNING

GROUP **C** FUNGICIDE

FARMALINX Flufol 500 SC Fungicide is a member of the DMI group of fungicides. For fungicide resistance management Flufol 500 SC Fungicide is a Group C fungicide. Some naturally occurring individual fungi resistant to Flufol 500 SC Fungicide and other Group C fungicides may exist through normal genetic variability in any

fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Flufol 500 SC Fungicide or other Group C fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, FARMALINX Pty Ltd accepts no liability for any losses that may result from the failure of Flufol 500 SC Fungicide to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will irritate the eyes. May irritate the skin. Avoid contact with the eyes and skin. When using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow length chemical resistant gloves and a face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre Phone Australia 13 11 26. Additional information is listed in the Material Safety Data Sheet.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for FARMALINX Flufol 500 SC Fungicide is available from FARMALINX Pty Ltd on request. Call Customer Service on 02 8338 9473.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions, or under off-label permits not endorsed by FARMALINX Pty Ltd or under abnormal conditions. FARMALINX Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Flufol 500 SC

FUNGICIDE

ACTIVE CONSTITUENT: 500g/L FLUTRIAFOL

GROUP **C** FUNGICIDE

For control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser as per the Directions for Use table

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

farmalinx

FARMALINX Pty Ltd ABN 95 134 353 245
South Sydney Corporate Park, Unit W2E,
75-85 O'Riordan Street, Alexandria NSW 2015

farmalinx

FARMALINX Pty Ltd ABN 95 134 353 245
South Sydney Corporate Park, Unit W2E,
75-85 O'Riordan Street, Alexandria NSW 2015
T: 02 8338 9473 F: 02 9313 4352
Website: www.farmalinx.com

APVMA Approval NO.: #####/#####