

# CAUTION

**KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**Farmoz**

# Imazagard® 750 WG

## FUNGICIDE

ACTIVE CONSTITUENT: 750 g/kg IMAZALIL

present as the SULPHATE

<b>GROUP</b>	3	<b>FUNGICIDE</b>
--------------	---	------------------

***For control of post-harvest disease of citrus, apples, pears and storage diseases in potatoes and rockmelons as specified in the Directions for Use table.***

[www.farmoz.com.au](http://www.farmoz.com.au)

### DIRECTIONS FOR USE

TREE CROP	DISEASE	RATE PER 100 L WATER	CRITICAL COMMENTS
Apples and Pears	Post-harvest storage decay: <i>Penicillium expansum</i> (Blue mould)	68 g provides a concentration of 500 ppm	Apply to fruit preferably within 24 hours of harvest. Use as a bulk dip, flood or drench for a maximum of 30 seconds. After treatment allow fruit to dry before storage.
Citrus Fruit	Post-harvest storage decays: <i>Penicillium italicum</i> (Blue mould), <i>Penicillium digitatum</i> (Green mould)		Apply to fruit preferably within 24 hours of harvest. Use as a bulk dip, flood or spray before waxing.
Rockmelon	Fusarium Fruit Rot ( <i>Fusarium</i> spp.), Sour Rot ( <i>Geotrichum candidum</i> ), Alternaria Fruit Rot ( <i>Alternaria</i> spp.), Rhizopus Soft Rot ( <i>Rhizopus stolonifer</i> ), Pink Mould Rot ( <i>Trichothecium roseum</i> )		<b>DO NOT use on rockmelons intended for export (except New Zealand).</b> Dip fruit for 45 seconds within 24 hours of harvest.
Potato Tubers	STORAGE DISEASES <i>Helminthosporium solani</i> (Silver Scurf), <i>Phoma exigua</i> (Gangrene), <i>Fusarium</i> spp. (Dry rot)	20 g to a maximum of 2 L water (15 15 g a.i imazalli)	Apply the solution as a mist at the rate of a maximum of 2 litres per tonne over the potatoes on the conveyor line prior to storage. IMAZAGARD 750 WG will give control of <i>Fusarium semitectum</i> and significant reduction of dry rot caused by <i>Fusarium trichoteciodes</i> . For good control of <i>Fusarium solani</i> a tank mix with thiabendazole (TBZ) is advised.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.**

## GENERAL INSTRUCTIONS

### FUNGICIDE RESISTANCE WARNING

FARMOZ IMAZAGARD 750 WG Fungicide is a member of the DMI group of fungicides. For **GROUP 3 FUNGICIDE** fungicide resistance management IMAZAGARD 750 WG is a Group 3 fungicide. Some naturally occurring individual fungi resistant to IMAZAGARD 750 WG and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, FARMOZ Pty. Ltd. accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

IMAZAGARD 750 WG is freely soluble in water and can be used without the addition of a wetting agent in water having a pH below 7. Use buffering agent to correct water pH to a neutral pH (ie pH = 6-7). Use IMAZAGARD 750 WG before waxing. DO NOT use in waxes. At the indicated rate IMAZAGARD 750 WG has curative and preventative action as well as antisporeulation properties.

### MIXING

Add the required amount of product to water and mix thoroughly. No wetting agent is required in water having a pH below 7.

### Recharging dip solution

To maintain the required volume of dip solution in the tank, add the required volume of water and product and mix thoroughly. The dip solution should be replaced when sediment accumulates.

### COMPATIBILITY

Laboratory tests have shown this product to be physically compatible with various DPA formulations. Because of diversity of field conditions, small scale tests should be carried out in the first instance.

### PROTECTION OF LIVESTOCK

DO NOT feed treated produce or by-products to food producing animals, including poultry.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate ponds, rivers, waterways or dams with the chemical or used containers.

### STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to dip solution. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the empty containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### DIP DISPOSAL

Dispose of dip in an authorized dip disposal facility.

### SAFETY DIRECTIONS

Harmful if swallowed. Avoid contact with eyes and skin. Wash hands after use.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

### MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for IMAZAGARD 750 WG is available from FARMOZ on request. Call Customer Service on (02) 9431 7800.

**CONDITIONS OF SALE:** The use of FARMOZ IMAZAGARD 750 WG Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by FARMOZ Pty. Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and FARMOZ Pty. Ltd. accepts no responsibility for any consequence whatsoever resulting from the use of this product.

® IMAZAGARD is a registered trademark of a Makhteshim-Agan company

**UN No. 3077  
ENVIRONMENTALLY HAZARDOUS SUBSTANCE,  
SOLID, N.O.S.  
HAZCHEM 2X  
PACKAGING GROUP III**



**PACK SIZE: 680 g, 1 kg**