

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Farmoz

Imazagard[®] 500 EC

FUNGICIDE

ACTIVE CONSTITUENT: 500 g/L IMAZALIL

GROUP 3 FUNGICIDE

For control of post-harvest disease of citrus as specified in the Directions for Use table.

www.farmoz.com.au

DIRECTIONS FOR USE

CROP	DISEASE	RATE	CRITICAL COMMENTS
Citrus	Post-harvest storage decays: Blue Mould (<i>Penicillium italicum</i>), Green Mould (<i>Penicillium digitatum</i>)	<u>In water:</u> 100 mL per 100 litres of water provides a concentration of 500 ppm <u>In citrus wax:</u> 1000 mL – 1600 mL per 200 litres of wax provides a concentration of 2500-4000 ppm	Apply to citrus fruit preferably within 24 hours of harvest. Use as a bulk dip, flood or spray before waxing. Apply in wax without further dilution using conventional waxing equipment. Sporulation control can be obtained at the rate of 1600 mL per 200 litres of wax (4000 ppm). No top up for wax.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:
CITRUS: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

FUNGICIDE RESISTANCE WARNING

FARMOZ IMAZAGARD 500 EC Fungicide is a member of the DMI group of fungicides. For **GROUP 3 FUNGICIDE** fungicide resistance management IMAZAGARD 500 EC Fungicide is a Group 3 fungicide. Some naturally occurring individual fungi resistant to IMAZAGARD 500 EC Fungicide and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, FARMOZ Pty. Ltd. accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

To avoid build up of resistant strains, label recommendations should be followed and a strict sanitation programme should be carried out throughout the season.

MIXING IN WATER

Make up a premix of at least five parts of water to one part of product and stir thoroughly. Fill up to the required volume with water and agitate. The emulsion will remain stable if agitated every four hours.

MIXING IN WAX

Make a premix of at least five parts of water to one part of product and stir thoroughly. Add the premix to the wax and stir thoroughly until possible creaming has disappeared. Constant agitation in the wax is advisable during application. At indicated rate used as above IMAZAGARD 500 EC has curative (up to max. 24 hours) and preventative action. At the higher use rate IMAZAGARD 500 EC has antispore properties. For disease control it is not necessary to apply consecutive applications of IMAZAGARD 500 EC to citrus fruit using both methods described above. One application in water or wax at the label rate is sufficient.

COMPATIBILITY

Compatible with most fungicides including Iprodione. Also compatible with water based citrus waxes. As formulations of other manufacturer's products are beyond the control of FARMOZ Pty. Ltd., all mixtures should be tested prior to mixing commercial quantities.

PROTECTION OF LIVESTOCK

DO NOT feed treated produce or by products to food producing animals, including poultry.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways or dams with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers. Dispose of at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt. Dispose of spent treatment solutions in a suitable waste pit clear of waterways, desirable vegetation and tree roots.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate skin. Will damage eyes. Avoid contact with eyes and skin. When opening the container and preparing the solution wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow length PVC gloves and goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for IMAZAGARD 500 EC is available from FARMOZ on request. Call Customer Service on (02) 9431 7800.

CONDITIONS OF SALE: The use of FARMOZ IMAZAGARD 500 EC Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by FARMOZ Pty. Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and FARMOZ Pty. Ltd. accepts no responsibility for any consequence whatsoever resulting from the use of this product.

® Registered trademark of FARMOZ Pty. Ltd.

**UN No. 3082
ENVIRONMENTALLY HAZARDOUS
SUBSTANCE, LIQUID, N.O.S.**

**HAZCHEM CODE: 2X
PACKAGING GROUP: III**



PACK SIZE: 1 L