

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

NUFARM
INTERVIX[®]

Herbicide for CLEARFIELD[®] crops

ACTIVE CONSTITUENT:
33 g/L IMAZAMOX present as the ammonium salt
15 g/L IMAZAPYR present as the ammonium salt

GROUP **B** HERBICIDE

For the early post-emergence control of certain annual grass and broadleaf weeds as part of the CLEARFIELD[®] Production System for canola, as specified in the DIRECTIONS FOR USE section of this label.

**IMPORTANT: READ THE ATTACHED LEAFLET
BEFORE OPENING OR USING THIS PRODUCT.**

Contents: 5 Litres

10 Litres

20 Litres

110 Litres

Nufarm Australia Limited
ACN 004 377 780
103-105 Pipe Road
Laverton North Victoria 3026
Tel: (03) 9282 1000
Fax: (03) 9282 1001



® Registered trademark of BASF

APVMA Approval No.: 60683/5L/0706
APVMA Approval No.: 60683/10L/0706
APVMA Approval No.: 60683/20L/0706
APVMA Approval No.: 60683/110L/0706

STORAGE AND DISPOSAL

(5L, 10L, 20L)

Store in the closed, original container in a dry, cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

Triple, or preferably pressure, rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

Envirodrum 110L Mini Bulk Returnable Container

Store the original sealed Envirodrum in a cool well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Nufarm Australia Limited.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the containers and preparing spray, wear cotton overalls (or equivalent clothing) buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

When tank mixing with other products consult also the safety directions for those products.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone: Australia 131126). If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, wash out immediately with water.

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS).

In case of emergency: Phone 1800 033 498

Ask for shift supervisor. Toll free 24 hours.

CONDITIONS OF SALE

"Nufarm Australia Limited (Nufarm) shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Nufarm's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Nufarm has any authority to add to or alter these conditions."

Batch No.:

Date of Manufacture:



(5L, 10L, 20L)



(110L only)

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

NUFARM
INTERVIX[®]

Herbicide for CLEARFIELD[®] crops

ACTIVE CONSTITUENT:
33 g/L IMAZAMOX present as the ammonium salt
15 g/L IMAZAPYR present as the ammonium salt

GROUP **B** HERBICIDE

For the early post-emergence control of certain annual grass and broadleaf weeds
as part of the CLEARFIELD[®] Production System for canola,
as specified in the DIRECTIONS FOR USE section of this label.

READ COMPLETE DIRECTIONS FOR USE
BEFORE USING THIS PRODUCT.

Nufarm Australia Limited
ACN 004 377 780
103-105 Pipe Road
Laverton North Victoria 3026
Tel: (03) 9282 1000
Fax: (03) 9282 1001



® Registered trademark of BASF

APVMA Approval No.: 60683/0706

DIRECTIONS FOR USE:

RESTRAINTS:

Apply ONLY to certified canola varieties with the CLEARFIELD Technology. DO NOT apply to triazine tolerant or conventional canola varieties.

DO NOT apply to crops that are stressed due to conditions such as waterlogging, too little moisture, frost, disease or nutritional disorders.

DO NOT apply by aircraft.

DO NOT apply if rain is expected within 2 hours of application.

DO NOT apply more than once per season to any one crop.

CROP	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Canola varieties with CLEARFIELD Technology only	Wild radish (<i>Raphanus raphanistrum</i>), Wild turnip (<i>Brassica tournefortii</i>)	300 to 500 mL/ha plus Hasten* or Kwickin* at 0.5L/100L spray volume	<p>Read Follow Crop comments and restrictions on the label prior to use.</p> <p>Apply to crop at the 2 to 6 leaf stage. Application to older crops is not harmful but weed control may be impaired because of poor contact and coverage of weeds.</p> <p>Apply to actively growing weeds in the 3-leaf to 2-tiller stage and broadleaf weeds in the 2 to 6 leaf stage.</p> <p>Use the higher rate when weed numbers are high or towards the upper end of the recommended growth stages, or when the crop is at the 5 to 6 leaf stage to ensure better contact and coverage.</p> <p>If other weeds require control, apply appropriate herbicides at least two weeks after INTERVIX Herbicide and only when signs of regrowth or renewed vigour appear, or the effects of INTERVIX Herbicide may affect their performance.</p>
	Capeweed (<i>Arctotheca calendula</i>), Field pea (<i>Pisum sativum</i>), Narrow leaf lupin (<i>Lupinus angustifolius</i>), Sub clover (<i>Trifolium subterraneum</i>), Wild radish (<i>Raphanus raphanistrum</i>), Wild turnip (<i>Brassica tournefortii</i>)	300 to 500 mL/ha plus Nufarm Archer® Herbicide at 150 to 300 mL/ha plus Hasten or Kwickin at 0.5L/100L spray volume	<p>Refer to critical comments for 300 to 500 mL/ha alone.</p> <p>Nufarm Archer Herbicide aids in the control of legume and composite weed species. Refer to the Archer label. Use rates above 150 mL/ha when these weeds are primary weeds in the paddock and when required by their growth stage.</p> <p>Archer above 150 mL/ha can slightly impair grass control. Refer to advice under three-way mixes.</p> <p>The addition of Archer does not affect the control of wild radish and wild turnip.</p> <p>(Refer to the Compatibility section of this label and the Archer label for further details of use)</p>
	Annual ryegrass ¹ (<i>Lolium rigidum</i>), Barley (<i>Hordeum vulgare</i>), Barley grass (<i>Hordeum leporinum</i>), Great brome (<i>Bromus diandrus</i>), Oat (<i>Avena sativa</i>), Rigid brome (<i>Bromus rigidus</i>), Wheat (<i>Triticum aestivum</i>), Wild oat (<i>Avena fatua</i>), Wild radish (<i>Raphanus raphanistrum</i>), Wild turnip (<i>Brassica tournefortii</i>)	300 to 500 mL/ha plus Aramo® Herbicide at 175 to 300 mL/ha plus Hasten or Kwickin at 0.5L/100L spray volume.	<p>Refer to critical comments for 300 to 500 mL/ha alone.</p> <p>DO NOT apply later than the 6-leaf stage of the crop as injury induced by Aramo may occur.</p> <p>Aramo Herbicide aids in the control of grasses. Refer to the Aramo label. INTERVIX Herbicide alone provides strong suppression of listed grasses other than annual ryegrass. The low rate of Aramo will normally be sufficient to achieve good control. Use rates above 175 mL/ha when grasses are primary weeds in the paddock, and when their growth stage requires it, to ensure highest levels of control.</p> <p>¹ Where Group A or Group B resistance in ryegrass is known, or thought to be developing in a paddock, an application of Nufarm TriflurX® Selective Herbicide should be made prior to sowing. Satisfactory control of populations with significant levels of either Group A and Group B resistance or both may not be achieved.</p> <p>The addition of Aramo does not affect the control of wild radish and wild turnip.</p> <p>(Refer to the Compatibility section of this label and the Aramo label for further details of use).</p>

CROP	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
<p>Canola varieties with CLEARFIELD Technology only (cont).</p>	<p>Annual ryegrass² (<i>Lolium rigidum</i>), Barley (<i>Hordeum vulgare</i>), Barley grass (<i>Hordeum leporinum</i>), Capeweed (<i>Arctotheca calendula</i>), Field pea (<i>Pisum sativum</i>), Great brome (<i>Bromus diandrus</i>), Narrow leaf lupin (<i>Lupinus angustifolius</i>), Oat (<i>Avena sativa</i>), Rigid brome (<i>Bromus rigidus</i>), Sub clover (<i>Trifolium subterraneum</i>), Wheat (<i>Triticum aestivum</i>), Wild oat (<i>Avena fatua</i>), Wild radish (<i>Raphanus raphanistrum</i>), Wild turnip (<i>Brassica tournefortii</i>)</p>	<p>300 to 500 mL/ha plus Nufarm Archer Herbicide at 150 to 300 mL/ha plus Aramo Herbicide at 175 to 300 mL/ha plus Hasten or Kwickin at 0.5L/100L spray volume.</p>	<p>Three-way tank mixes may be considered when legume and composite weeds and grasses are present together with wild radish or wild turnip.</p> <p>Refer to critical comments for 300 to 500 mL/ha alone, and in separate mixes with Nufarm Archer Herbicide and Aramo Herbicide.</p> <p>DO NOT apply later than the 6-leaf stage of the crop as injury induced by Aramo may occur.</p> <p>Archer and Aramo together aid in the control of legumes, composites and grasses. Refer to the Archer and Aramo labels. INTERVIX Herbicide alone provides varied degrees of suppression of all species listed. Use the low rates of Aramo and Archer for light infestations of target weeds.</p> <p>Use Aramo above 175 mL/ha when grasses are primary weeds and when their growth stage requires it. Use Archer above 150 mL/ha when legumes and composites are primary weeds and when their growth stage requires it.</p> <p>The addition of Aramo does not affect the control of broad leaf weeds. The addition of Archer above 150 mL/ha can impair grass control. DO NOT use above 150 mL/ha if a very high level of grass control is a primary objective.</p> <p>²Where Group A or Group B resistance in ryegrass is known, or thought to be developing in a paddock, an application of Nufarm TriflurX Selective Herbicide should be made prior to sowing. Satisfactory control of populations with significant levels of either Group A or Group B resistance or both may not be achieved.</p> <p>(Refer to the Compatibility section of this label and the Archer and Aramo labels for further details of use).</p>
	<p>Barley (<i>Hordeum vulgare</i>), Barley grass (<i>Hordeum leporinum</i>), Great brome (<i>Bromus diandrus</i>), Indian hedge mustard (<i>Sisymbrium orientale</i>), Oat (<i>Avena sativa</i>), Rigid brome (<i>Bromus rigidus</i>), Marshmallow (<i>Malva parviflora</i>), Sub clover (<i>Trifolium subterraneum</i>), Wheat (<i>Triticum aestivum</i>), Wild oat (<i>Avena fatua</i>), Wild radish (<i>Raphanus raphanistrum</i>), Wild turnip (<i>Brassica tournefortii</i>)</p> <p>Suppression only: Annual ryegrass³ (<i>Lolium rigidum</i>), Bedstraw spp. (<i>Galium tricornutum</i> and <i>G. aparine</i>), Doublegee (<i>Emex australis</i>), Silver grasses (<i>Vulpia bromoides</i> and <i>B. myuros</i>)</p>	<p>600 to 750 mL/ha plus Hasten or Kwickin at 0.5L/100L spray volume</p>	<p>Apply to crop at the 2 to 6 leaf stage. Application to older crops is not harmful but weed control may be impaired because of poor contact and coverage of weeds.</p> <p>Apply to actively growing grass weeds in the 3-leaf to 2-tiller stage and broadleaf weeds in the 2 to 6 leaf stage.</p> <p>Use the higher rate when weed numbers are high or towards the upper end of the recommended growth stages, or when the crop is at the 5 to 6 leaf stage, to ensure better contact and coverage.</p> <p>Weeds will either be killed in high numbers (weeds controlled) or in moderate numbers (weeds suppressed). In both cases, surviving plants will be stunted and will be uncompetitive with the crop, and seed set will be prevented or greatly reduced.</p> <p>If other weed species require control, apply appropriate herbicides at least two weeks after INTERVIX Herbicide and only when signs of regrowth or renewed vigour appear or the effects of INTERVIX Herbicide may affect their performance.</p> <p>³The control of annual ryegrass varies from excellent to poor depending on the status of Group B resistance in the population and environmental conditions. Where the population is expected to exceed 200 plants per sqm, or a high level of control is required, or the ryegrass is known to be resistant or thought to be developing resistance, an application of Nufarm TriflurX Selective Herbicide should be made prior to sowing. A tank mix with Aramo Herbicide may also be necessary.</p>

CROP	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
<p>Canola varieties with CLEARFIELD Technology only (cont).</p>	<p>Barley (<i>Hordeum vulgare</i>), Barley grass (<i>Hordeum leporinum</i>), Capeweed (<i>Arctotheca calendula</i>), Field pea (<i>Pisum sativum</i>), Great brome (<i>Bromus diandrus</i>), Indian hedge mustard (<i>Sisymbrium orientale</i>), Narrow leaf lupin (<i>Lupinus angustifolius</i>), Oat (<i>Avena sativa</i>), Rigid brome (<i>Bromus rigidus</i>), Marshmallow (<i>Malva parviflora</i>), Sub clover (<i>Trifolium subterraneum</i>), Wheat (<i>Triticum aestivum</i>), Wild oat (<i>Avena fatua</i>), Wild radish (<i>Raphanus raphanistrum</i>), Wild turnip (<i>Brassica tournefortii</i>)</p> <p>Suppression only: Annual ryegrass⁴ (<i>Lolium rigidum</i>), Bedstraw spp. (<i>Galium tricornutum</i> and <i>G. aparine</i>), Doublegee (<i>Emex australis</i>), Silver grasses (<i>Vulpia bromoides</i> and <i>B. myuros</i>)</p>	<p>600 to 750 mL/ha plus Nufarm Archer Herbicide at 150 to 300 mL/ha plus Hasten or Kwickin at 0.5L/100L spray volume</p>	<p>Refer to critical comments for 600 to 750 mL/ha alone.</p> <p>Nufarm Archer Herbicide aids in the control of legume and composite weed species. Refer to the Archer label. Use rates above 150 mL/ha when these weeds are primary weeds in the paddock and when required by their growth stage.</p> <p>Archer above 150 mL/ha can slightly impair grass control. When legumes or composites are primary weeds together with grasses specifically ryegrass, refer to advice under three-way mixes.</p> <p>The addition of Archer does not affect the control of other weeds controlled by INTERVIX Herbicide.</p> <p>⁴The control of annual ryegrass by INTERVIX Herbicide varies from excellent to poor depending on the status of Group B resistance in the population and environmental conditions. Where the population is expected to exceed 200 plants per sqm, a high level of control is required, and the ryegrass is known to be resistant, or thought to be developing resistance, an application of Nufarm TriflurX Selective Herbicide should be made prior to sowing. A tank mix with Aramo Herbicide should also be considered. DO NOT tank mix with Archer above 150 mL/ha without including Aramo in a three-way mix.</p> <p>(Refer to the Compatibility section of this label and the Archer label for further details of use).</p>
	<p>Annual ryegrass⁵ (<i>Lolium rigidum</i>), Barley (<i>Hordeum vulgare</i>), Barley grass (<i>Hordeum leporinum</i>), Great brome (<i>Bromus diandrus</i>), Indian hedge mustard (<i>Sisymbrium orientale</i>), Oat (<i>Avena sativa</i>), Rigid brome (<i>Bromus rigidus</i>), Marshmallow (<i>Malva parviflora</i>), Sub clover (<i>Trifolium subterraneum</i>), Wheat (<i>Triticum aestivum</i>), Wild oat (<i>Avena fatua</i>), Wild radish (<i>Raphanus raphanistrum</i>), Wild turnip (<i>Brassica tournefortii</i>)</p> <p>Suppression only: Bedstraw spp. (<i>Galium tricornutum</i> and <i>G. aparine</i>), Doublegee (<i>Emex australis</i>), Silver grasses (<i>Vulpia bromoides</i> and <i>B. myuros</i>)</p>	<p>600 to 750 mL/ha plus Aramo Herbicide at 175 to 300 mL/ha plus Hasten or Kwickin at 0.5L/100L spray volume.</p>	<p>Refer to critical comments for 600 to 750 mL/ha alone.</p> <p>DO NOT apply later than the 6-leaf stage of the crop as injury induced by Aramo may occur. Refer to the Aramo Herbicide label.</p> <p>INTERVIX Herbicide alone provides very good to excellent control of listed grasses other than annual ryegrass. The control of annual ryegrass varies from excellent to poor depending on the status of Group B resistance in the population and environmental conditions.</p> <p>This tank mix need only be considered to ensure the highest level of broad-spectrum grass control including ryegrass, or when three-way mixes with Nufarm Archer Herbicide at rates above 150 mL/ha are used to ensure against minor antagonism.</p> <p>The low rate of Aramo will normally be sufficient to achieve good control. Use rates above 175 mL/ha only when ryegrass is a primary weed, or when it is known to have a level of Group B resistance, or when the majority of plants are tillering.</p> <p>⁵Where Group A or Group B resistance in ryegrass is known, or thought to be developing in a paddock, an application of Nufarm TriflurX Selective Herbicide should be made prior to sowing. Satisfactory control of populations with significant levels of either Group A and Group B resistance or both may not be achieved.</p> <p>The addition of Aramo does not affect the control of wild radish and wild turnip.</p> <p>(Refer to the Compatibility section of this label and the Aramo label for further details of use).</p>

CROP	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
	<p>Annual ryegrass⁶ (<i>Lolium rigidum</i>), Barley (<i>Hordeum vulgare</i>), Barley grass (<i>Hordeum leporinum</i>), Capeweed (<i>Arctotheca calendula</i>), Field pea (<i>Pisum sativum</i>), Great brome (<i>Bromus diandrus</i>), Indian hedge mustard (<i>Sisymbrium orientale</i>), Lupin (<i>Lupinus</i> spp.), Oat (<i>Avena sativa</i>), Rigid brome (<i>Bromus rigidus</i>), Sub clover (<i>Trifolium subterraneum</i>), Wheat (<i>Triticum aestivum</i>), Wild oat (<i>Avena fatua</i>), Wild radish (<i>Raphanus raphanistrum</i>), Wild turnip (<i>Brassica tournefortii</i>)</p> <p>Suppression only: Bedstraw spp. (<i>Galium tricorutum</i> and <i>G. aparine</i>), Doublegee (<i>Emex australis</i>). Silver grasses (<i>Vulpia bromoides</i> and <i>B. myuros</i>)</p>	<p>600 to 750 mL/ha plus Nufarm Archer Herbicide at 150 to 300 mL/ha plus Aramo Herbicide at 175 to 300 mL/ha plus Hasten or Kwickin at 0.5L/100L spray volume.</p>	<p>Three-way tank mixes may be considered when legume and composite weeds and grasses are present together with wild radish or wild turnip or other weeds controlled by INTERVIX Herbicide.</p> <p>Refer to critical comments for 600 to 750 mL/ha alone, and in separate mixes with Nufarm Archer Herbicide and Aramo Herbicide.</p> <p>DO NOT apply later than the 6-leaf stage of the crop as injury induced by Aramo may occur.</p> <p>Refer to the labels of Nufarm Archer Herbicide and Aramo Herbicide. INTERVIX Herbicide alone controls or suppresses to various degrees all species listed. The low rates of Aramo and Archer should be adequate for lighter infestations of target weeds.</p> <p>Use Aramo above 175 mL/ha when ryegrass is a primary weed, or when it is known to have a level of Group B resistance, or when the majority of plants are tillering, or when Archer is used above 150 mL/ha.</p> <p>The addition of Aramo does not affect the control of broad leaf weeds. The addition of Archer above 150 mL/ha can slightly impair grass control but the inclusion of Aramo overcomes this problem.</p> <p>⁶Where Group A or Group B resistance in ryegrass is known, or thought to be developing in a paddock, an application of Nufarm TriflurX Selective Herbicide should be made prior to sowing. Satisfactory control of populations with significant levels of either Group A or Group B resistance or both may not be achieved.</p> <p>(Refer to the Compatibility section of this label and the Archer and Aramo labels for further details of use).</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

Canola with CLEARFIELD Technology: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 5 WEEKS AFTER APPLICATION.

Harvest For Grain: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

Nufarm INTERVIX Herbicide is for use in canola varieties with CLEARFIELD Technology only. Varieties with CLEARFIELD Technology are those that have been bred specifically to be tolerant to INTERVIX Herbicide.

INTERVIX Herbicide is absorbed through the leaves, green stems and roots of susceptible weeds and moved from the point of contact throughout the plant. Weeds will either die or will remain stunted and will not compete with the crop. Symptoms of kill may take one to two weeks to develop with death occurring up to one month from treatment. Symptoms first appear at the growth points where young foliage becomes discoloured and distorted before dying.

INTERVIX Herbicide is primarily a post-emergence product. Best results will be achieved when good contact and coverage of weeds occurs and weeds are actively growing. The product must be mixed with Hasten or Kwickin as per the Directions for Use.

INTERVIX Herbicide also has some residual soil activity under good soil moisture conditions. Residual effects on weeds can be reduced when dry soil conditions follow application before the herbicide has moved to the root zone. Best results will be achieved when application is made to moist soil or if approximately 10 mm rain follows within several days of application.

Vigorous crop growth will assist in suppressing weeds not completely killed and those germinating later.

MIXING

INTERVIX Herbicide is a water-soluble liquid (SL) formulation. Pour the required amount of the product into a spray tank containing almost the total amount of water required. Mix thoroughly. If INTERVIX Herbicide is added during filling, foaming may occur. If excessive foaming becomes a problem, add a silicone based antifoaming agent at the manufacturers recommended rate. DO NOT use a suction probe unless the antifoaming agent has already been added to the spray tank water. Consult your distributor for specific information on suitable antifoaming agents. When tank mixing this product with other recommended compatible products, first add the other product(s) to the tank and mix thoroughly before adding INTERVIX Herbicide.

INTERVIX Herbicide may be applied in hard or soft water. The product is corrosive to mild steel. Use ONLY stainless steel, fibreglass, plastic or plastic-lined containers for mixing, storage and application.

APPLICATION

DO NOT apply by aircraft.

Apply in minimum 70 L water per hectare. When the crop is older than 4 leaf or very leafy or when the total weed population exceeds 200 plants per m², apply in a minimum of 100 L water per hectare to improve contact and coverage. INTERVIX Herbicide should be applied a minimum of two hours before rainfall or irrigation. If tank-mixed with other products, follow recommendations for the mixing partner should these extend beyond two hours.

EQUIPMENT CLEAN-UP

Thoroughly flush all spray equipment with water following the use of INTERVIX Herbicide and before use with other products. If tank-mixed with other products, also follow clean-up procedures recommended for the mixing partner.

COMPATIBILITY

INTERVIX Herbicide is chemically compatible with FASTAC[®] Duo Insecticide, Sonic[®] 200EC Insecticide, Nufarm Dimethoate Systemic Insecticide, Le-Mat* 290SL Insecticide and Nufarm Endosulfan 350EC Insecticide. It is also chemically and biologically compatible with Nufarm Archer[®] Herbicide, Lontrel* 750 SG Herbicide and Aramo[®] Herbicide and chemically compatible with Select* Herbicide. Simultaneous mixes of Archer and Aramo with INTERVIX Herbicide (three-way mixes) chemically and biologically compatible. Other simultaneous mixes have not been tested.

All tank mixes are chemically stable over a 24-hour period. In the event of delayed spraying, store tank load out of direct sunlight and maintain agitation if possible. Mixes with Archer and Aramo are also biologically stable over a 24-hour delay. Biological stability of other mixes is unknown.

RE-ENTRY

DO NOT re-enter treated areas until spray has dried. If re-entry is necessary wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves.

RESISTANT WEEDS WARNING

GROUP B HERBICIDE

INTERVIX Herbicide is a member of the imidazolinone group of herbicides. The product has the inhibitors-of-acetolactate-synthase (ALS) mode of action. For weed resistance management, the product is a Group B herbicide. Some naturally-occurring weed biotypes resistant to the product and other Group B herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group B herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Australia Ltd accepts no liability for any losses that may result from failure of this product to control resistant weeds.

BEST MANAGEMENT PRACTICE (BMP) PROGRAM

A detailed program has been developed that outlines sound agronomic and integrated weed management practices, designed to optimise the performance of the CLEARFIELD Production System and minimise the potential for the development of herbicide resistance in weed populations. Consultation on BMP should be undertaken with a CLEARFIELD-accredited agronomist prior to use of INTERVIX Herbicide in the CLEARFIELD Production System for canola. Implementation of the BMP is an essential part of herbicide resistance management.

FOLLOW CROPS

This product, like all other imidazolinone herbicides, is broken down in the soil by microbes in wet, aerobic conditions. Under conditions that do not favour breakdown, carry-over soil residues can affect susceptible follow crops. Normally safe residue levels may still affect follow crops when soil nutrition is low or marginal, when cold and very wet soil conditions prevail, or when soil pathogens or nematodes are present. As environmental and agronomic factors make it impossible to eliminate all risks associated with this product, rotational crop injury is always possible.

Note: when the intention is to grow cereals on CLEARFIELD canola stubble (treated with INTERVIX) self-sown canola volunteers must be removed before they mature beyond 2-leaf, all macro and micro-nutrients must be maintained at levels necessary to grow the planned crops, and sulfonyleureas must not be used.

The following minimum re-cropping intervals (months after application) should be observed.

Months after Application	Follow Crops
0	Canola varieties with CLEARFIELD Technology, Wheat varieties with CLEARFIELD Technology
10	Chickpeas, Faba beans, Field peas, Lucerne, Lupins, Pasture legumes, vetch, Oats* , Triticale* , Barley* , Non-CLEARFIELD Wheat*
34	Triazine tolerant and conventional Canola, All other crops

*** Barley, Oats, Triticale, Non-CLEARFIELD Wheat:**

The following additional requirements apply if it is intended to sow these cereals during the next winter season:

- DO NOT apply INTERVIX Herbicide later than the end of August (no later than the end of July in WA).
- DO NOT use INTERVIX Herbicide in areas where rainfall from spraying of cereals is expected to be below **150mm (for 300-375mL/ha use), 200mm (for up to 500mL/ha use) and 250mm (for 600-750mL/ha use).**
- **DO NOT use above 375mL/ha in the Lower Great Southern region of Western Australia.**

For all situations, if expected rainfall is not received following use of INTERVIX Herbicide, consult your local Nufarm representative before planting non-CLEARFIELD cereals. In calculating rainfall actually received, place greater emphasis on rain received from application up to the end of Spring and lesser emphasis on break rains. If single **isolated** heavy summer and autumn falls are required to achieve rainfall targets consult your local Nufarm representative for advice.

Trial work is constantly in progress in order to refine these recommendations.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT spray within 50 m of wetlands or waterways.

CROP SAFETY

This product may, in some circumstances, lead to transient crop yellowing and temporary slowing of growth but plants soon recover and yield is unaffected. This effect may be more pronounced when the product is used under poor growth conditions. DO NOT use this product on any canola variety other than certified varieties with CLEARFIELD technology. Extreme crop damage will result to triazine tolerant and conventional canola varieties.

PROTECTION OF LIVESTOCK

INTERVIX Herbicide is of low hazard to bees.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

(5L, 10L, 20L)

Store in the closed, original container in a dry, cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

Triple, or preferably pressure, rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

Envirodrum 110L Mini Bulk Returnable Container

Store the original sealed Envirodrum in a cool well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Nufarm Australia Limited.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the containers and preparing spray, wear cotton overalls (or equivalent clothing) buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

When tank mixing with other products consult also the safety directions for those products.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, wash out immediately with water.

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS).

In case of an emergency: Phone 1800 033 498

Ask for shift supervisor. Toll free 24 hours.

CONDITIONS OF SALE

"Nufarm Australia Limited ('Nufarm') shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Nufarm's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Nufarm has any authority to add to or alter these conditions."

® Registered trademark of BASF
Archer, Aramo, Triflur X, Sonic
are registered trademarks of
Nufarm Australia Limited.

* Other registered trademarks.