

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

GENFARM IMIDACLOPRID 200SC INSECTICIDE

ACTIVE CONSTITUENT: 200 g/L IMIDACLOPRID

GROUP	4A	INSECTICIDE
--------------	-----------	--------------------

For the control of various Insect pests of cotton, Fruit, vegetables and
Ornamentals as specified in The DIRECTIONS FOR USE table

CONTENTS: 1L, 10L, 20L, 60L

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USING THIS PRODUCT

Genfarm Crop Protection Pty Ltd
Suite 2, Level 3, 64 Talavera Road,
Macquarie Park,
NSW, 2113
Tel: (02) 9889 5400

STORAGE AND DISPOSAL

Store in the closed original container in a cool, dry, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings or spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers to below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use. When preparing product for use, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia:13 11 26)

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which is available from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which the buyer uses it. Genfarm Crop Protection Pty Ltd accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

APVMA Approval No.: 62165/1L/0708, 62165/10L/0708, 62165/20L/0708, 62165/60L/0708

Batch No;

Date of Manufacture:

In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS AUSTRALIA WIDE
--	---

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

GENFARM IMIDACLOPRID 200SC INSECTICIDE

ACTIVE CONSTITUENT: 200 g/L IMIDACLOPRID

GROUP	4A	INSECTICIDE
--------------	-----------	--------------------

For the control of various Insect pests of cotton, Fruit, vegetables and
Ornamentals as specified in The DIRECTIONS FOR USE table

APVMA APPROVAL NO: 62165/0708

THIS BOOKLET IS PART OF THE LABEL

Genfarm Crop Protection Pty Ltd
Suite 2, Level 3, 64 Talavera Road,
Macquarie Park,
NSW, 2113
Tel: (02) 9889 5400

DIRECTIONS FOR USE

FOLIAR SPRAY APPLICATIONS

CROP	PEST	RATE	WHP	CRITICAL COMMENTS
Cotton	Aphids	250 mL/ha + Pulse Penetrant at 0.2% v/v (2 mL/L water) or equivalent organosilicone surfactant (eg Freeway Gold)	13 weeks	<p>The addition of Pulse Penetrant or equivalent is critical for the performance of Genfarm Imidacloprid 200SC Insecticide. Apply early in the establishment of an aphid infestation when numbers are low (ie no more than 1 or 2 leaves per plant with honeydew present). Applications made later than this may result in reduced control.</p> <p>Shorter residual control may be evident and a repeat application of a registered aphicide (follow the Cotton Insecticide Resistance Management Strategy for cotton aphid) may be required to achieve complete control:</p> <ul style="list-style-type: none"> ◆ if applications of Genfarm Imidacloprid 200SC Insecticide plus Pulse or equivalent are timed too late (see above); or ◆ if existing high density aphid colonies (hotspots) are present; or ◆ if aphids have established throughout the plant canopy (especially lower in the canopy); or ◆ if there is high reinfestation pressure; or ◆ if there is rapid crop growth; or ◆ if Genfarm Imidacloprid 200SC Insecticide plus Pulse or equivalent is used following a spray-failure (eg resistance to organophosphate or carbamate insecticides). Note: Where resistance to carbamates or organophosphates is suspected, Genfarm Imidacloprid 200SC Insecticide plus Pulse or equivalent should be used first so as not to delay control of the aphids present. <p>Aphids treated with Genfarm Imidacloprid 200SC Insecticide plus Pulse or equivalent may still be present on the plant but will not be feeding. Control of aphids should initially be assessed by a reduction in fresh honeydew and not on the presence of aphids on the plant. After ingesting Genfarm Imidacloprid 200SC Insecticide, aphids may take up to 5 days to die.</p>
	Mirids, Brown flea beetle			Apply when pest numbers reach treatment threshold levels as determined by field checks.
Stone fruit	Green peach aphid Black peach aphid	<p>Dilute spraying 25 mL/100L</p> <p>Concentrate spraying Refer to Mixing/Application section</p>	21 days	Apply at first sign of aphid infestation. Apply as a full cover spray, ensuring thorough coverage. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Do not use in equipment that requires rates greater than 125 mL/100 L of water (i.e. greater than 5 X concentrate).

CROP	PEST	RATE	WHP	CRITICAL COMMENTS
Cucurbits	Green peach aphid	25 mL/100 L or 300 mL/ha	1 day	Apply at first sign of aphid infestation.
Capsicum Egg plant Potato			7 days	
Tomato			3 days	
Brassicas	Grey cabbage aphid Turnip aphid		7 days	Apply at first sign of aphid infestation. Add a wetting agent.
Sweet potato	Silverleaf whitefly, including type B	25 mL/100 L or 250 mL/ha	7 days	Apply at first sign of whitefly or melon thrips infestation. Apply dilute sprays (25 mL/100 L) to run off. Ensure thorough coverage of underside of leaves. Use of droppers will improve coverage of underside of leaves.
Cucumber			1 day	
Egg plant	Melon thrips		7 days	
Roses	Aphids	25 mL/100 L	-	Apply as a thorough cover spray at first sign of insect infestation.
Ornamental plants	Aphids Azalea lace bug Bronze orange bug, Harlequin bug, Citrus mealybug Greenhouse thrips, Fullers rose weevil			
	Hibiscus flower beetle	50 mL/100 L		Spray buds and flowers as needed.
	Longtailed mealybug	50 mL/100 L + surfactant		Apply 3 sprays 2 weeks apart. Use a non-ionic surfactant at label rate.
	Psyllids	25 mL/100 L		Spray at first sign and then a week later.
	Soft scales	25 mL/100 L		Spray in late spring or when small scales are first seen. Apply 3 sprays 2 weeks apart. Add a wetting agent.
Duboisia	Green peach aphid	25 mL/100 L		Apply when aphid numbers reach spray threshold levels as determined by regular monitoring. Ensure thorough coverage of all leaves.
Pandanus trees	Flatid (<i>Jamella australiae</i>)	Spot spray 875 mL/100 L of water Stem injection 1.75 L/1 L of water		Spot spray: Spray 100 mL of mixture directly into the leafy throat of each head. Stem injection: Drill holes 0.5 to 1 cm in diameter and 10 cm deep at an angle of 30° 1 to 1.5 m above ground level. Drill one hold per limb (or trunk in single trunked trees). Apply 5 mL of mixture in each hole and seal the hole. Do not reapply in the same holes. Uptake of Genfarm Imidacloprid 200SC Insecticide, and therefore control of the pest in heavily infested heads already showing severe damage, will be slow and may be incomplete.

SOIL DRENCH APPLICATIONS

CROP	PEST	RATE	CRITICAL COMMENTS
Elm	Elm leaf beetle	7 mL/25 mm of tree diameter at breast height	Mix the required dose in sufficient water to adequately treat each tree. Use at least 50 L of mix per tree up to a tree diameter of 400-500 mm and then 100 L per tree for larger trees. Inject mix to a depth of 20-30 cm in a minimum of 4 injection sites per tree, 0.75 to 1.5 m apart, arranged in an evenly spaced grid to just beyond the drop line. Ensure root zone is adequately moist with active root growth. Keep treated area moist for 7-10 days after treatment. Treat at least 6-10 weeks prior to pest attack in late winter or early spring when roots are active. DO NOT treat if soil is waterlogged.
Azaleas in pots	Azalea lace bug	3.5 mL/250 mL water/pot	Use as a soil drench for pots up to 20 L capacity. Prior to application remove mulch and dead vegetation, and moisten the soil surface. Apply the Genfarm Imidacloprid 200SC Insecticide mixture, and then water it in well immediately after application.
Ornamentals in pots	Scarab beetle larvae	3.5 mL/5 L water	Use as a soil drench. 5 L of mixture will treat twenty 6 L pots. Prior to application remove mulch and dead vegetation, and moisten the soil surface. Apply the Genfarm Imidacloprid 200SC Insecticide mixture, and then water it in well immediately after application.
Roses	Aphids	3.5 mL/ ² L water/plant	Use as a soil drench by pouring mixture evenly around drop zone. Use this rate for plants up to 1 m high. For each additional metre of plant height, add 2 mL extra of Genfarm Imidacloprid 200SC Insecticide to the 2 L of water. Prior to application remove mulch and dead vegetation, and moisten the soil surface. Apply the Genfarm Imidacloprid 200SC Insecticide mixture, and then water it in well immediately after application.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

Cotton: DO NOT HARVEST FOR 13 WEEKS AFTER APPLICATION

DO NOT GRAZE OR CUT FOR STOCK FOOD

Cucurbits: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION

Tomatoes: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION

Brassicas, Capsicum, Eggplant, Potatoes, Sweet potatoes:

DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION

Stone fruit: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION

DO NOT graze any treated area, or cut for stock food.

DO NOT feed produce harvested from treated area to animals, including poultry.

GENERAL INSTRUCTIONS

GROUP	4A	INSECTICIDE
-------	----	-------------

Insecticide Resistance Warning

For insecticide resistance management, Genfarm Imidacloprid 200SC Insecticide is a Group 4A insecticide. Some naturally occurring insect biotypes resistant to Genfarm Imidacloprid 200SC Insecticide and other Group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Genfarm Imidacloprid 200SC Insecticide and other Group 4A insecticides are used repeatedly. The effectiveness of Genfarm Imidacloprid 200SC Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use. Genfarm Crop Protection Pty Ltd accepts no liability for any losses that may result from the failure of Genfarm Imidacloprid 200SC Insecticide to control resistant insects. Genfarm Imidacloprid 200SC Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, Genfarm representative or local agricultural department agronomist.

Resistance Management Strategy

Cotton aphid in cotton

Observe the cotton industry insecticide Resistance Management Strategy (IRMS).

Aphids, whitefly and melon thrips in various crops

Do not apply Genfarm Imidacloprid 200SC Insecticide (or other group 4A insecticides) in consecutive sprays within and between seasons. Rotate with registered insecticides from other mode of action groups.

Confined environments such as glasshouses:

Annuals – Do not apply more than one spray of Genfarm Imidacloprid 200SC Insecticide (or other group 4A insecticides) to any one crop.

Perennials – Rotate with registered insecticides from other groups. Use a maximum of three Genfarm Imidacloprid 200SC Insecticide (or other group 4A insecticide) sprays in any 12 month period.

Mixing/Application

Prior to pouring, shake container vigorously, then add the required quantity of Genfarm Imidacloprid 200SC Insecticide to water in the spray vat while stirring or with agitators in motion.

Special Instructions for Stone Fruit

Dilute Spraying

- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way:
EXAMPLE ONLY
 1. Dilute spray volume as determined above: For example 1500 L/ha
 2. Your chosen concentrate spray volume: For example 500 L/ha
 3. The concentration factor in this example is: 3 X (ie 1500 L ÷ 500 L = 3)
 4. As the dilute label rate is 25 mL/100 L for stone fruit, then the concentrate rate becomes 3 x 25, that is 75 mL/100 L of concentrate spray.
- The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.
- Do not use a concentrate rate higher than that specified in the Critical Comments.

- For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

Application (Cotton)

Thorough coverage of cotton plants is essential to achieve maximum performance from Genfarm Imidacloprid 200SC Insecticide plus Pulse. Equipment should be calibrated to achieve a **minimum of 60 droplets/cm² on the target foliage**. A droplet **Volume Median Diameter (VMD)** for optimum performance from Genfarm Imidacloprid 200SC Insecticide plus Pulse is dependent on equipment and is defined below. Do not apply when unfavourable environmental conditions may reduce the quality of spray coverage.

Ground Application (Cotton)

Application using ground equipment should be made using **hollow cone nozzles** with a **minimum spray volume of 100 L/ha**. Hollow cone nozzles are recommended but if flat fan nozzles are used, higher water volumes will be required and nozzles should be configured to ensure thorough coverage. A droplet **VMD of 150 – 180 microns** must be used. Where multiple nozzles per row are used, they should be of the same specification to ensure that each nozzle contributes an equal proportion of the required dose. Where multiple nozzles per row are used (particularly for banded applications) ensure the correct nozzle overlap pattern is achieved on the target foliage. **Banded applications less than 100% are not recommended beyond the 15 node crop stage.**

Aerial Application (Cotton)

Apply in a **minimum spray volume of 25 L/ha**. A droplet **VMD of 120 -150 microns** must be used. Do not exaggerate swath width or exceed a swath width of 20 to 22 m. Do not apply Genfarm Imidacloprid 200SC Insecticide plus Pulse using Ultra Low Volume (ULV) methods. The use of large droplet placement equipment is not recommended.

Compatibility

Genfarm Imidacloprid 200SC Insecticide 200 SC is compatible with Antracol, Baycor, Nitofol. Do not mix concentrates together but add each to the spray tank separately. As formulations of other manufacturers' products are beyond the control of Genfarm, all mixtures should be tested prior to mixing commercial quantities. As changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays, Genfarm cannot be responsible for the behaviour of such mixtures.

Export of Treated Produce

Growers should note that MRLs or import tolerances do not exist in all markets for edible produce treated with Genfarm Imidacloprid 200SC Insecticide. If you are growing edible produce for export, please check with Genfarm for the latest information on MRLs and import tolerances before using Genfarm Imidacloprid 200SC Insecticide.

Note on Ornamentals

Genfarm Imidacloprid 200SC Insecticide 200 SC Insecticide has been used on a wide range of ornamental plant species without damage. However, some species and varieties are particularly sensitive to chemical sprays and as this is often related to local conditions it is advisable to treat only a small number of plants first, in order to ascertain their reaction before treating the whole crop.

PROTECTION OF LIVESTOCK

Dangerous to bees. DO NOT spray any plants in flower while bees are foraging.
DO NOT graze any treated area, or cut for stock food.
DO NOT feed produce harvested from treated area to animals, including poultry.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. A spraydrift minimization strategy should be employed at all times when aerially applying sprays. The strategy envisaged is exemplified by the cotton industry's Best Management Practices Manual.

STORAGE AND DISPOSAL

Store in the closed original container in a cool, dry, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings or spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers to below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use. When preparing product for use, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia: 13 11 26)

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which is available from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of Genfarm Crop Protection Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which the buyer uses it. Genfarm Crop Protection Pty Ltd accepts no responsibility for any consequences resulting from the use of this product. Genfarm Crop Protection Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

Genfarm Crop Protection Pty Ltd

Suite 2, Level 3, 64 Talavera Road,
Macquarie Park,
NSW, 2113
Tel: (02) 9889 5400