

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Farmoz

Chief[®] 250 SC

SEED DRESSING FUNGICIDE

ACTIVE CONSTITUENT: 250 g/L IPRADIONE

GROUP **2** FUNGICIDE

For control of Brown Leaf Spot and the suppression of Hypocotyl Rot, in the first four weeks after emergence in Lupins and for the control of Hypocotyl Rot in Potatoes as specified in the Directions for Use table.

www.farmoz.com.au

DIRECTIONS FOR USE:

CROP	DISEASE	STATE	RATE	CRITICAL COMMENTS
Lupins	Brown Leaf Spot (<i>Pleiochaeta setosa</i>), Rhizoctonia Hypocotyl Rot (<i>Rhizoctonia solani</i>) (Suppression only)	WA only	Pre-sowing Treatment (Feb-May) 100-400 mL/100 kg seed Post-harvest Treatment (Dec-Jan) 400 mL/100 kg seed	Mix with water and apply in a total volume of 600 mL mixture/100 kg seed using suitable seed dressing application equipment. Ensure thorough coverage of seed. Use in conjunction with other disease management practices. See General Instructions for details. Use high rate in areas likely to be exposed to high disease pressure. (eg. Central and Eastern Wheat Belt areas with close rotations and a history of Brown Leaf Spot or where other disease management practices are not followed). Application of CHIEF 250 SC will provide suppression of Hypocotyl Rot and can reduce Pleiochaeta Root Rot in some situations.
	Brown Leaf Spot (<i>Pleiochaeta setosa</i>)	NSW, Vic, SA only	500 mL/100 kg seed in 500 mL water	Apply using suitable seed dressing application equipment. Ensure thorough coverage of seed. Application can reduce Pleiochaeta Root Rot in some situations.
Potatoes	Hypocotyl Rot (Black Scurf) (<i>Rhizoctonia solani</i>)	All States	800 mL/tonne seed material	CHIEF 250 SC will protect emerging shoots from Hypocotyl Rot, improving overall germination. CHIEF 250 SC may also reduce occurrence of Black Scurf on the harvested potatoes. Ensure good coverage of seed material and planting furrow. DO NOT plant into waterlogged soil.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD (Harvest and Grazing):

LUPINS, POTATOES: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

Lupins only: This product will reduce inoculum levels on inoculated seed and should only be used on seed to be sown in ground where existing soil inoculum levels will provide adequate nodulation. Spray inoculation is not recommended.

This product should be used as part of an integrated disease management strategy, particularly in areas with high disease pressure. Where possible, the following disease management practices should be used in conjunction with CHIEF 250 SC:

- Stubble Retention
- Direct Drilling
- Correct Seeding Depth
- Correct Rotations
- Clean Seed Source
- Correct Seeding Rate
- Avoid Late Sowing

MIXING

Shake the container well before using. Add the required volume of CHIEF 250 SC to the water whilst agitating.

APPLICATION

Lupins only: Slowly add the mixture to the seed and mix thoroughly ensuring thorough coverage of the seed.

Potatoes only: Ensure good coverage of seed material and planting furrow. This can be achieved by applying CHIEF 250 SC as a fine spray to the seed at the time of planting using spray equipment mounted on the planter, and nozzles located at three points on each planter row to ensure uniform coating of the seed. A minimum water volume of 80 L/tonne should be used.

Further information can be obtained from a FARMOZ representative.

FUNGICIDE RESISTANCE WARNING

FARMOZ CHIEF® 250 SC Seed Dressing Fungicide is a member of the Dicarboximide group of fungicides. For fungicide resistance management CHIEF 250 SC is a Group 2 fungicide.

GROUP **2** FUNGICIDE

Some naturally occurring individual fungi resistant to CHIEF 250 SC and other Group 2 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by CHIEF 250 SC or other Group 2 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, FARMOZ Pty. Ltd. accepts no liability for any losses that may result from the failure of CHIEF 250 SC to control resistant fungi.

PROTECTION OF LIVESTOCK

DO NOT feed treated seed to animals, including poultry. Store treated seed away from feed.

DO NOT allow treated seed to contaminate grain intended for animal consumption.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT feed treated seed or otherwise expose wildlife or domestic birds. DO NOT contaminate streams, rivers, or waterways with the chemical or bags which have held treated seed.

PROTECTION OF OTHERS

When treated seed is stored, it should be kept apart from other grains or seeds, and the containers or bags should be clearly marked to show that the contents have been treated with CHIEF 250 SC.

DO NOT use treated seed for human consumption.

DO NOT allow treated seed to contaminate food for human consumption.

Bags which have held treated seed should not be used for any other purpose. Any seed not used for sowing should be destroyed.

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN. Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling break, crush, or puncture, and bury empty containers in a local authority landfill. If no landfill is available, bury the empty containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. If product on skin, immediately wash area with soap and water. Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, DO NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for CHIEF 250 SC is available from FARMOZ on request. Call Customer Service on (02) 9431 7800.

CONDITIONS OF SALE: The use of FARMOZ CHIEF 250 SC Seed Dressing Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by FARMOZ Pty. Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and FARMOZ Pty Ltd accepts no responsibility for any consequence whatsoever from the use of this product.

® Registered trademark of a Makhteshim-Agan Company

NOT A DANGEROUS GOOD ACCORDING TO THE AUSTRALIAN DANGEROUS GOODS (ADG) CODE.

PACK SIZES: 5 L, 20 L