

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CORVETTE[®] FLOWABLE

Fungicide

ACTIVE CONSTITUENT: 500 g/L IPRADIONE

GROUP **B** FUNGICIDE

For control of certain fungal diseases in various crops and situations as specified in the Directions for Use table

IMPORTANT : READ THE ATTACHED LEAFLET BEFORE USE

CONTENTS : 5 Litres



Crop Care Australasia Pty Ltd
Unit 15/16 Metroplex Avenue
Murarrie Qld 4172

STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. Wear suitable protective clothing, gloves and goggles. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from your supplier or from the Crop Care website: www.cropcare.com.au

Conditions of sale

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Care Australasia Pty Ltd within thirty days of delivery.

In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 498 ALL HOURS - AUSTRALIA WIDE
---	--

® Corvette is a Registered Trademark.



Crop Care Australasia Pty Ltd
ACN 061 362 347
Unit 15/16 Metroplex Avenue
MURARRIE Qld 4172
Tel : (07) 3909 2000

Insert Barcode,
drumMUSTER
logo and Label
code as required

APVMA Approval No. : 59702/5L/0608

BN

DOM

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CORVETTE[®] FLOWABLE

Fungicide

ACTIVE CONSTITUENT: 500 g/L IPRADIONE

GROUP	B	FUNGICIDE
-------	----------	-----------

For control of certain fungal diseases in various crops and situations as specified in the Directions for Use table

IMPORTANT : READ THIS LEAFLET BEFORE USING THIS PRODUCT

Crop Care Australasia Pty Ltd, Unit 15/16 Metroplex Avenue, Murarrie Qld 4172

DIRECTIONS FOR USE

Tree Crops/Vines:

RATE					CRITICAL COMMENTS
In the following table, all rates are given for dilute spraying. For concentrate spraying, refer to the Special Instructions for Tree Crops/Vines section.					For all uses in this table: Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.
CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Almonds	Blossom blight, Brown rot (<i>Monilinia</i> spp., <i>Sclerotinia</i> spp.)	All States	50 mL/100 L water	Nil	Apply first at full bloom and if conditions are favourable for disease development, up to two subsequent applications can be made; at petal fall and up to four weeks after petal fall.
Boysenberries	Grey mould (<i>Botrytis cinerea</i>)		100 mL/100 L water	1 day	Spray at 10% blossom and full bloom. For fruit protection, apply at 2 to 3 weeks pre-harvest.
Grapes				7 days	Good crop hygiene will aid in the control of disease. This use is subject to a CropLife fungicide resistance management strategy: 1. If three or fewer bunch rot sprays are applied in a season, use only one spray per season containing Corvette Flowable (or other Group B fungicides). 2. If four or more bunch rot sprays are applied in a season, use no more than two sprays of Corvette Flowable (or other Group B fungicides). 3. DO NOT apply more than two consecutive sprays of Corvette Flowable (or other Group B fungicides), including from the end of one season to the start of the following season. 4. Late season fungicide treatments should be applied before botrytis infection reaches unacceptably high levels in the vineyard.
Kiwifruit	Botrytis blight (<i>Botrytis</i> spp.)	NSW, Vic & WA only			Apply the spray to vines every 10 to 14 days ensuring that all fruit is thoroughly wet. Apply 3 applications at 10 to 14 day intervals from 10% bloom to petal fall for protection of flowers and young fruit. Apply a further 2 applications of Corvette Flowable to control late season Botrytis.
Macadamias	Botrytis blight (<i>Botrytis</i> spp.)	All States	50 mL/100 L water	Nil	Apply as a thorough cover spray to flower racemes when they open. A follow up spray may be needed one week later if wet conditions persist during flowering. Remove nuts under trees prior to spraying.
Mandarins (non-bearing)	Alternaria leaf spot (brown spot) (<i>Alternaria alternata</i>)	Qld, WA & NT only	100 mL/100 L water		Apply to non-bearing trees of Murcott variety monthly from first flush in spring until flushing ceases in the autumn. Reduce intervals to fortnightly during periods of wet weather.
Passionfruit	Alternata spot (brown spot) (<i>Alternaria</i> spp., <i>Alternaria passiflorae</i>)	Qld, NSW, WA & NT only		7 days	This use is subject to a CropLife fungicide resistance management strategy: 1. Maintain a protective cover with protectant fungicide such as mancozeb. 2. Limit the use of Corvette Flowable to strategic periods, i.e. before, during and after extended wet periods. 3. Always tank mix Corvette Flowable with a protectant such as mancozeb. 4. DO NOT apply more than four Corvette Flowable (or other Group B fungicide) sprays in a season.

Raspberries	Grey mould (<i>Botrytis cinerea</i>)	All States	100 mL/100 L water	1 day	Spray at 10% blossom and full bloom. For fruit protection, apply at 2 to 3 weeks pre-harvest.
Stone fruit: Apricots Cherries Nectarines Peaches Plums	Orchard Spraying Blossom blight (<i>Monilinia fructicola</i> , <i>Monilinia laxa</i>) Brown rot (<i>Monilinia fructicola</i> , <i>Monilinia laxa</i>)		50 to 75 mL/100 L water	Nil	For control of blossom blight, spray at 10% blossom, full bloom and petal/shuck fall. For control of subsequent brown rot in fruit, spray at 3 weeks and 1 week pre-harvest. Use the higher rate under severe conditions of challenge, or for single applications of Corvette Flowable in the spray program. This use is subject to a CropLife fungicide resistance management strategy: 1. DO NOT apply more than 2 consecutive sprays of Corvette Flowable (or other Group B fungicides). 2. A post-harvest treatment should also be counted as an application. 3. The last blossom blight spray and the first pre-harvest brown rot spray should be regarded as consecutive applications. 4. The spray program should be considered and the strategy applied on a whole-orchard basis.
Youngberries	Grey mould (<i>Botrytis cinerea</i>)		100 mL/100 L water	1 day	Spray at 10% blossom and full bloom. For fruit protection, apply at 2 to 3 weeks pre-harvest.

Fruit – Post-harvest dipping:

CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Pome fruit: Apples Pears	Post-harvest dipping Storage rots (<i>Penicillium</i> spp.) (<i>Botrytis</i> spp.) (<i>Gloeosporium</i> spp.)	All States	100 mL/100 L water	Nil	To minimise the development of post-harvest rots handle fruit carefully to avoid fruit injury and dip promptly after harvest. Remove any infected fruit from the packing house immediately and destroy. When dipping, allow sufficient time to thoroughly wet the fruit. Top up dip with 100 mL Corvette Flowable in 100 L of water. This use is subject to a CropLife fungicide resistance management strategy: 1. For the last pre-harvest spray, use a fungicide with a different mode of action to the fungicide planned for use as a post-harvest treatment. 2. Where alternative fungicide groups are available, rotate to use as many different modes of action as possible.
Stone fruit: Apricots Cherries Nectarines Peaches Plums	Post-harvest dipping Brown rot (<i>Monilinia fructicola</i> , <i>Monilinia laxa</i>) *Transit rot (<i>Rhizopus</i> spp.)				To minimise the development of post-harvest rots handle fruit carefully to avoid fruit injury and dip promptly after harvest. Remove any infected fruit from the packing house immediately and destroy. When dipping, allow sufficient time to thoroughly wet the fruit. Top up dip with 200 mL Corvette Flowable in 100 L of water. A non-ionic wetting agent should be added. *Transit rot is suppressed at this rate. This use is subject to a CropLife fungicide resistance management strategy: 1. For the last pre-harvest spray, use a fungicide with a different mode of action to the fungicide planned for use as a post-harvest treatment. 2. Where alternative fungicide groups are available, rotate to use as many different modes of action as possible.

Berries:

(See Tree Crops/Vines for boysenberries, raspberries and youngberries)

CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Strawberries	Grey mould (<i>Botrytis cinerea</i>)	All States	1.0 L/ha where spray volume is less than 1000 L/ha OR 100 mL/100 L water where spray volume equals or exceeds 1000 L/ha	1 day	This use is subject to a CropLife fungicide resistance management strategy: 1. Apply a program of protectant fungicides during flowering. If conditions favour disease development during this period, use Corvette Flowable. 2. DO NOT apply more than two successive sprays of Corvette Flowable (or other Group B fungicides).

Vegetables:

CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Celery	Sclerotinia rot (pink rot) (<i>Sclerotinia sclerotiorum</i>)	All States	1.0 L/ha where spray volume is less than 1000 L/ha OR	1 day	Commence spraying 1 to 2 weeks post-transplanting and then every 2 to 3 weeks. Use only five sprays.
Lettuces	Sclerotinia rot (drop) (<i>Sclerotinia sclerotiorum</i> , <i>Sclerotinia minor</i>)	Tas, WA & NT only	100 mL/100 L water where spray volume equals or exceeds 1000 L/ha	7 days	Spray should be directed to the stems at ground level and to the underside of lower leaves. This use is subject to a CropLife fungicide resistance management strategy: 1. Apply Corvette Flowable as a seedling drench soon after emergence. 2. Apply a protectant fungicide as a high volume foliar spray before planting out, then Corvette Flowable immediately after planting. 3. Maintain a cover with protectant fungicide sprays at 7-10 day intervals. 4. If weather conditions favour Botrytis infection, tank mix the protectant with Corvette Flowable. 5. DO NOT apply Corvette Flowable (or other Group B fungicides) more than four times per season, irrespective of the target disease.
	Grey mould (<i>Botrytis</i> spp.)	Tas, WA & NT only			

Potatoes	Sclerotinia rot (<i>Sclerotinia sclerotiorum</i>)	All States	500 mL to 1.0 L/ha where spray volume is less than 1000 L/ha	Nil	Apply 2 sprays, once immediately before and once immediately after hilling-up. For most effective treatment, concentrate the spray at the base of the stems and surrounding soil surface, where the fungus is active. Use the higher rate where disease is severe.
	Target spot (early blight) (<i>Alternaria solani</i>)		OR 50 to 100 mL/100 L water where spray volume equals or exceeds 1000 L/ha		Ensure thorough coverage to the whole plant. Treatment is generally not required until after flowering. Use the higher rate where disease is severe. This use is subject to a CropLife fungicide resistance management strategy: 1. Limit the use of Corvette Flowable to periods when conditions favour disease development. 2. DO NOT apply more than six Corvette Flowable (or other Group B fungicide) sprays in one season. 3. Apply no more than two consecutive sprays of Corvette Flowable (or other Group B fungicides).
	Hypocotyl rot (black scurf) (<i>Rhizoctonia solani</i>)		400 mL/ tonne seed material		Corvette Flowable will protect emerging shoots from hypocotyl rot, improving overall germination. Corvette Flowable may also reduce occurrence of black scurf on the harvested potatoes. Ensure good coverage of seed material and planting furrow. This can be achieved by applying Corvette Flowable as a fine spray to the seed at the time of planting using spray equipment mounted on the planter, and nozzles located at three points on each planter row to ensure uniform coating of the seed. DO NOT plant into waterlogged soil. A minimum water volume of 80 L/tonne seed should be used.
Tomatoes	Sclerotinia rot (<i>Sclerotinia sclerotiorum</i>)	All States	1.0 L/ha where spray volume is less than 1000 L/ha	7 days	Spray at 14-day intervals from transplanting and throughout the period of disease pressure.
	Grey mould (<i>Botrytis cinerea</i>)		OR 100 mL/100 L water where spray volume equals or exceeds 1000 L/ha		Commence spraying 3 to 4 weeks after transplanting or at the onset of disease. Repeat treatment at 14-day intervals or when conditions favour spread of the disease, i.e. at trimming or deleafing. This use is subject to a CropLife fungicide resistance management strategy: 1. Alternate or tank mix Corvette Flowable with a protectant such as chlorothalonil (eg. Barrack®). Avoid applying two Corvette Flowable (or other Group B fungicide) sprays in succession, unless tank mixed with a protectant. 2. DO NOT apply more than four Corvette Flowable (or other Group B fungicide) sprays in a season.
	Target spot (early blight) (<i>Alternaria solani</i>)	Qld, Tas, WA & NT only			Commence spraying 1 week post-transplanting. Use adequate water to give thorough coverage of the plants. Use high volume spray equipment. This use is subject to a CropLife fungicide resistance management strategy: 1. Limit the use of Corvette Flowable to periods when conditions favour disease development. 2. DO NOT apply more than four Corvette Flowable (or other Group B fungicide) sprays in one season. 3. Apply no more than two consecutive sprays of Corvette Flowable (or other Group B fungicides).

Field Crops:

CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Lucerne	Lucerne leaf spot (<i>Stemphylium botryosum</i>)	Qld & WA only	250 mL to 500 mL/ha where spray volume is less than 1000 L/ha OR 25 to 50 mL per 100 L water where spray volume equals or exceeds 1000 L/ha	7 days	Spray every 10 to 14 days when cool, damp weather favours the disease. Use the higher rate under conditions of high disease pressure.
	Leptosphaerulina leaf spot (<i>Leptosphaerulina trifolii</i>)				Apply in at least 300 L water/ha every 10 to 14 days when cool, damp weather favours the disease. Use the higher rate under conditions of high disease pressure.
Peanuts	Sclerotinia rot (<i>Sclerotinia sclerotiorum</i> , <i>Sclerotinia minor</i>)		1.0 L/ha OR 220 mL/100 L water (spot application)	12 days	Apply when disease first appears. Repeat if necessary. Use a high water volume to ensure good coverage of foliage and stem at ground level. Do not mix Corvette Flowable with a foliar fungicide due to the different target positions on the plant.
Soybeans	Black leaf blight (<i>Arkoala nigra</i>)	NSW & WA only	1.0 L/200 to 400 L water/ ha	7 weeks	If disease is present on leaves apply an initial spray at early pod set (pods approximately 5 mm long). An additional spray 14 days later may be required if wet seasonal conditions prevail.

Ornamentals and Turf:

CROP	DISEASE	STATE	RATE	WHP	CRITICAL COMMENTS
Ornamentals	Botrytis blight (<i>Botrytis cinerea</i>)	All States	100 mL/100 L water	Nil	Spray at 14 day intervals commencing when the disease first becomes apparent and continuing until conditions no longer favour the disease. Spraying saintpaulia and poinsettia flowers may result in some petal scorch. Tepid water should be used and wet plants protected from direct sunlight. This use is subject to a CropLife fungicide resistance management strategy: 1. DO NOT apply more than two consecutive sprays of Corvette Flowable (or other Group B fungicides).

Recreational Turf	Brown patch (<i>Rhizoctonia solani</i>) Dollar spot (<i>Sclerotinia homoeocarpa</i>)	All States	60 to 90 mL/10 to 150 L water/100m ²	-	<u>Curative Program</u> : Repeat treatment after 14 days. Use the higher rate under conditions of severe disease pressure. <u>For preventative treatment</u> : Apply as a monthly spray program. This use is subject to a CropLife fungicide resistance management strategy: 1. DO NOT apply more than two consecutive sprays of Corvette Flowable (or other Group B fungicides), unless mixed with a protectant fungicide from Group X or Y .
	Brown patch (<i>Curvularia</i> spp.) (<i>Dreschlera</i> spp.) (<i>Fusarium culmorum</i>) (<i>Rhizoctonia solani</i>)	Vic only			
	Fusarium patch (<i>Fusarium nivale</i>)	All States	90 mL/10 to 150 L water/100m ²		<u>For Curative Treatment</u> : Repeat treatment in 14 days. <u>For Preventative Treatment</u> : Apply as a monthly spray program. This use is subject to a CropLife fungicide resistance management strategy: 1. DO NOT apply more than two consecutive sprays of Corvette Flowable (or other Group B fungicides), unless mixed with a protectant fungicide from Group X or Y .
	Leaf spot (<i>Dreschlera</i> spp.)		60 mL/10 to 150 L water/100m ²		
	Spring dead spot (<i>Leptosphaeria</i> spp.)*	NSW, SA & ACT only	65 mL/10 to 150 L water/100m ²		<u>For Preventative Treatment only</u> : Apply as a monthly soil drench throughout the year. Water in thoroughly after application. *When used in a preventative program this product will limit the development of <i>Leptosphaeria</i> and improve the rate of turf recovery. This use is subject to a CropLife fungicide resistance management strategy: 1. DO NOT apply more than two consecutive sprays of Corvette Flowable (or other Group B fungicides), unless mixed with a protectant fungicide from Group X or Y .
	Spring dead spot (<i>Helminthosporium</i> spp.) <i>Curvularia</i> spp.)	Vic, WA & NT only			
	Helminthosporium disease affecting leaf, crown and roots: (Black and white Helminthosporium spot: <i>Bipolaris</i> spp. <i>Drechslera</i> spp. <i>Exserohilum</i> spp.)	All States	45 mL/10 to 150 L water/100m ²		<u>For Curative Treatment</u> : Apply as a foliar spray and repeat application after 7 to 14 days. For white spot symptoms on couch grass (<i>Cynodon dactylon</i>) only, use higher water volumes of 100 to 150L of water per 100m ² . <u>For Preventative Treatment</u> : Apply as a high volume foliar spray. Repeat applications at monthly intervals from February to May. This use is subject to a CropLife fungicide resistance management strategy: 1. DO NOT apply more than two consecutive sprays of Corvette Flowable Fungicide (or other Group B fungicides), unless mixed with a protectant fungicide from Group X or Y .

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS :

Almonds, macadamias, mandarins, potatoes, stone fruit: NOT REQUIRED WHEN USED AS DIRECTED

Boysenberries, celery, raspberries, strawberries, youngberries: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION

Grapes, kiwifruit, lettuce, tomatoes and passionfruit: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION

Peanuts: DO NOT HARVEST FOR 12 DAYS AFTER APPLICATION

Soybeans: DO NOT HARVEST FOR 7 WEEKS AFTER APPLICATION

Lucerne: DO NOT GRAZE OR CUT FOR STOCK FOOD WITHIN 7 DAYS OF TREATMENT

Export of Treated Produce

Growers should note that MRLs or import tolerances may not exist in all markets for produce treated with Corvette Flowable. If you are growing produce for export, please check with Crop Care Australasia Pty Ltd for the latest information on MRLs and import tolerances BEFORE using Corvette Flowable.

GENERAL INSTRUCTIONS

Fungicide Resistance Warning

GROUP	B	FUNGICIDE
--------------	----------	------------------

Corvette Flowable Fungicide is a member of the dicarboximide group of fungicides. For fungicide resistance management the product is a Group B fungicide.

Some naturally occurring individual fungi resistant to the product and other Group B fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group B fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Crop Care Australasia Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

Resistance Management

Resistant strains of fungi can develop to this product and other fungicides. To reduce the possibility of this occurrence, and where alternatives are available, rotate to use products with as many different modes of action as possible. Where specific resistance management strategies are established these are detailed in the Critical Comments.

Mixing

Note: Corvette Flowable may be unstable in conditions where the pH is 7 or higher. It is therefore essential to check the pH of the spray solution before adding Corvette Flowable. A suitable registered buffering agent (eg. Primabuff® Bio-Degradable Adjuvant) may have to be added to bring the pH down below 7.

Add the required amount of Corvette Flowable to the spray tank containing half the required volume of water. Mix thoroughly and make up to the required volume with water.

Application

Good disease control requires even, thorough coverage of the target area. Application should be made using appropriate spray equipment and sufficient water to provide adequate penetration and coverage. Equipment settings and water volume may need to vary, depending on the growth stage of the crop.

Special Instructions for Tree Crops/Vines

Dilute Spraying

- ◆ Use a sprayer designed to apply high spray volumes, up to the point of run-off and matched to the crop being sprayed.
- ◆ Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient spray solution to cover the crop to the point of run-off. Avoid excessive run-off.
- ◆ The required spray volume to achieve point of run off may be determined by applying different test volumes, using different settings on the sprayer, or from industry guidelines or other expert advice.
- ◆ Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of run-off.

- ◆ The required dilute spray volume to achieve point of run off will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying

- ◆ Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies spray volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- ◆ Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen spray volume.
- ◆ Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- ◆ The mixing rate for concentrate spraying can then be calculated in the following way:

EXAMPLE ONLY

1. Dilute spray volume as determined above: For example 1500 L/ha
 2. Your chosen concentrate spray volume: For example 500 L/ha
 3. The concentration factor in this example is: 3 X (i.e. $1500 \text{ L} \div 500 \text{ L} = 3$)
 4. If the dilute label rate is 100 mL/100 L, then the concentrate rate becomes 3 x 100, that is 300 mL of product per 100 L water for concentrate spraying.
- ◆ The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.
 - ◆ For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

Turf Nutrition

This product acts very rapidly and where the turf is of a low nutritional status, a slight yellowing may be seen. This will be corrected by nitrogenous fertilizer application. Where possible, it is recommended that the fertilizer program should precede the fungicide application.

Compatibility

This product may be combined with:

*Aliette[®] WG (see NOTE below), Bugmaster[®] Flowable, calcium chloride, chlorpyrifos 500 g/L EC, copper oxychloride, dimethoate (e.g. Saboteur[®]), Dithane[®] DF, Dithane M-45[®], DPA, endosulfan (e.g. Endosan[®] 350EC), fenitrothion, Kelthane[®] EC, maldison, metalaxyl, methomyl (e.g. Lannate[®], Nudrin[®]) or parathion-methyl.

When tank mixing products the order of mixing is determined by formulation type. As a guide the following mixing sequence is recommended:

- | | |
|---|------------------------------|
| 1. Wettable powders | 6. Solutions |
| 2. Suspension concentrates (e.g. Corvette Flowable) | 7. Emulsifiable concentrates |
| 3. Water Dispersible Granules | 8. Soluble concentrates |
| 4. Suspo-emulsions (e.g. Corvette Liquid) | 9. Wetting agents and oils |
| 5. Soluble powders | |

With any mixture, thoroughly agitate immediately before applying. It is not recommended to mix this product with more than one of the above chemicals in the tank. DO NOT tank mix this product with fertilisers. Mixtures with some fertilisers, e.g. urea, may cause foliar damage.

NOTE: *Mixing Corvette Flowable with Aliette[®] WG may result in some settling out.

As formulations of other manufacturers' products are beyond the control of Crop Care Australasia Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF LIVESTOCK

Do not graze treated areas or feed turf clippings from treated areas to animals including poultry.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Toxic to aquatic organisms. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. Wear suitable protective clothing, gloves and goggles. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from your supplier or from the Crop Care website: www.cropcare.com.au

Conditions of sale

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Care Australasia Pty Ltd within thirty days of delivery.

In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 498 ALL HOURS - AUSTRALIA WIDE
---	--

Corvette[®] is a Registered Trademark.

Aliette[®], Barrack[®], Bugmaster[®], Dithane M45[®], Endosan[®], Kelthane[®], Lannate[®], Nudrin[®], Primabuff[®] and Saboteur[®] are Registered Trademarks.



Crop Care Australasia Pty Ltd
ACN 061 362 347
Unit 15/16 Metroplex Avenue
MURARRIE Qld 4172
Tel : (07) 3909 2000

APVMA Approval No. : 59702/0608