

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FYFANON[®] ULV

INSECTICIDE

ACTIVE CONSTITUENT: 1169 g/L MALDISON
(an anticholinesterase compound)

GROUP	1B	INSECTICIDE
-------	-----------	-------------

An ultra low volume formulation for the control of a wide range of insect pests in certain crops and eucalypts as shown under Directions for Use

CONTENTS (20 litres, 200 litres)

CHEMINOVA AUSTRALIA PTY LTD
A.B.N. 59 472 616 773
A.C.N. 110 199 169
12 JULIUS AVE
NORTH RYDE NSW 2113
PH: 1800 624 597
www.cheminova.com.au

DIRECTIONS FOR USE RESTRAINT

DO NOT apply to melons or cucumbers when wet.

CROP/ SITUATION	PEST	STATE	RATE/ha	WHP days	CRITICAL COMMENTS
Cereal crops, Pastures, Pasture seed crops	Black Field Cricket	NSW, Vic, SA, WA only	450 mL	1	Apply in the early morning, when crickets are active or swarming.
	Small Plague Grasshopper	All States			Apply to infested areas.
Cereals, Pastures, Non crop areas	Black Field Cricket	Vic, SA, WA, Tas, NT only	Bait 110 mL/10 kg whole or kibbled grain/ha (Use clean wheat, barley or oats, not dusty grain)		Immature crickets: Mix in a drum or cement mixer. As treated grain absorbs Fyfanon rapidly, it is unnecessary to leave standing and grain can be used immediately.
			Bait 110-220 mL/10-20 kg whole or kibbled grain/ha (Use clean wheat, barley or oats, not dusty grain)		Mature crickets: Mix in a drum or cement mixer. As treated grain absorbs Fyfanon rapidly, it is unnecessary to leave standing and grain can be used immediately. Excess grain will kill moderate numbers of re-invading crickets as treated grain remains active for 4-6 weeks. Use higher rate for heavy pest pressure, where plentiful alternative feed exists, or for certainty of control. Spread in late afternoon and evening early in the season, and morning late in the season. Baiting may be ineffective if large amounts of pasture seed are present.
Cereal crops, Lucerne, Pastures, Pasture seed crops, Grain legume crops	Redlegged earth mite, Lucerne Flea	NSW only	225 mL		Apply after hatching, usually 3-4 weeks after opening rains.
		Vic, SA, WA only	275 mL		
Cereal crops, Maize, Pastures, Pasture seed crops, Rice, Sorghum	Common Armyworm	NSW, Vic, Qld only	700 mL		Spray entire crop or pasture when infestation is widespread. Spray along the front when pests are moving as an army.
Crucifers, Cucurbits, Canola (Rapeseed), Safflower, Sunflower	Rutherglen Bug	Vic, Qld, SA, WA, Tas only	450-900 mL	3	Use higher rate when plant growth is dense. Sunflowers: spray at bud stage
Linseed	Common Armyworm	Vic, Qld only	700 mL	1	Apply when Armyworms are actively feeding, in the early morning or late afternoon
Lucerne, Pastures, Pasture seed crops	Bluegreen Aphid, Spotted Aphid	All States	450 mL	1	Apply at first sign of pest. Repeat as necessary.
	Wingless Grasshopper				Apply to infested areas. To prevent re-invasion, spray a wide protective barrier around crop or pasture.
Maize, Pasture, Pasture seed crops, Sorghum	Australian Plague Locust		700 mL		Apply in hopper bands directly onto locusts.
Peaches	Oriental Fruit Moth	NSW, Vic, SA only	900 mL	3	Apply before trees are in full leaf at first sign of pest and repeat at signs of further moth flights.
Peas	Pea Weevil	NSW, Vic, SA, WA only	550 mL		Apply when first flowers begin to wither.

CROP/SITUATION	PEST	STATE	RATE/ha	WHP days	CRITICAL COMMENTS
Pome fruit, Stone fruit	Wingless Grasshopper	NSW, Vic, Qld, SA, Tas only	450 mL		Apply to infested areas.
		WA only	560 mL		
Sorghum	Sorghum Midge	NSW, Vic, Qld, SA only	450 mL	1	Apply at the flowering stage when adult pest numbers indicate likely damage. Repeat as necessary.
	Rutherglen Bug	All States	450-900 mL		Use higher rate when plant growth is dense.
Stone Fruit				3	Use higher rate when tree growth is dense.
Eucalyptus forests	Phasmatid Nymphs	NSW, Vic, SA only	450 mL	-	Apply at first sign of pest. Repeat as necessary.
Mosquito breeding areas	Adult Mosquitoes	All States	280 mL		Apply when adults emerge. Repeat as necessary. When fogging dissolve product in a suitable diluent. Dilution rate is dependent on output speed over ground and swath width of fogger. (Refer to General Instructions).

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

Cereals, lucerne, pastures, forage crops: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 1 DAY AFTER APPLICATION

Fruit, vegetables: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION

GENERAL INSTRUCTIONS

FYFANON is a contact insecticide and only partial control will be achieved if insects are protected from spray by dense foliage or if spray coverage is inadequate.

All vehicles should be removed from areas to be sprayed as paintwork may be damaged.

MIXING

FYFANON should be used undiluted except when fogging. When fogging for mosquito control, dilute with a kerosene/diesel distillate mixture of 1 L : 8 L, respectively.

APPLICATION

Apply by aircraft or mist spray using properly calibrated equipment designed for ultra low volume application.

INSECTICIDE RESISTANCE WARNING

GROUP	1B	INSECTICIDE
--------------	-----------	--------------------

For insecticide resistance management Fyfanon ULV Insecticide is a Group 1B insecticide. Some naturally occurring insect biotypes resistant to Fyfanon ULV Insecticide and other Group 1B insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Fyfanon ULV Insecticide or other Group 1B insecticides are used repeatedly. The effectiveness of Fyfanon ULV Insecticide on resistant individuals could be significantly reduced. Since the occurrence of resistant individuals is difficult to detect prior to use, Cheminova Australia Pty Ltd accepts no liability for any losses that may result from the failure of Fyfanon ULV Insecticide to control resistant insects. Fyfanon ULV Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, Cheminova Australia Pty Ltd representative or local agricultural department agronomist.

Read Directions for Use, Precautionary Statements, Safety Directions and First Aid instructions prior to opening. Use only at the recommended growth stages. Use only on recommended crops.

PRECAUTIONS

Avoid contact with food, food utensils, or places where food is prepared or stored.

RE-ENTRY PERIOD

DO NOT allow entry into treated areas until the spray has dried. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist and a washable hat and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF LIVESTOCK

Dangerous to bees. DO NOT spray on any plants in flower while bees are foraging. DO NOT spray over or upwind of

beehives. Remove bees from the area before blanket spraying.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto adjacent areas, particularly wetlands, waterbodies or watercourses. Dangerous to fish and aquatic invertebrates. DO NOT contaminate streams, rivers or waterways with the chemical or used containers

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsing to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length pvc gloves. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If swallowed, splashed on skin or in eyes, or inhaled, contact a Poisons Information Centre Phone Australia 131126 or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised. Give atropine if instructed.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet available from www.cheminova.com.au.

WARNING

This product must be used, handled and stored strictly as directed in accordance with this label, packaging and other reference material ("Directions"). To the extent permitted by law Cheminova Australia Pty Ltd and its related companies will have no liability for any injury, loss or damage arising from a failure to follow the Directions.

B/N

DOM Bar Code

FOR 24 HOUR SPECIALIST ADVICE

IN EMERGENCY ONLY

PHONE 1800 033 111

APVMA Approval No 49539/100187

® Registered Trademark of Cheminova A/S, DENMARK