

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SIPCAM
MCPA 500
SELECTIVE HERBICIDE

**ACTIVE CONSTITUENT: 500g/L MCPA
PRESENT AS THE DIMETHYLAMINE SALT**

GROUP	I	HERBICIDE
-------	----------	-----------

**FOR SELECTIVE CONTROL OF BROADLEAF WEEDS IN CEREALS,
LINSEED, PASTURES, SUGAR CANE AND TURF
AS SPECIFIED IN THE DIRECTIONS FOR USE TABLE**

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

Contents: 5, 20 200 Litres

**SIPCAM PACIFIC AUSTRALIA PTY LTD
Suite 11, 23-31 Gheringhap Street
GEELONG VIC 3220
ACN 073 176 888
TEL: 03 5223 3746**

NRA Approval No.

GENERAL INSTRUCTIONS:

Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavourable. The best conditions are when the soil is moist, weather fine and rain unlikely within 6 hours of application.

DO NOT spray crops or weeds outside the stages indicated in the Table as damage, loss of yield or inadequate weed control may result.

Resistant Weeds Warning:

Sipcam MCPA 500 Selective Herbicide is a member of the Phenoxy group of herbicides. Sipcam MCPA 500 Selective Herbicide has the disruptors of plant cell growth mode of action. For weed resistance management Sipcam MCPA 500 Selective Herbicide is a Group I herbicide. Some naturally-occurring weed biotypes resistant to Sipcam MCPA 500 Selective Herbicide and other disruptors of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Sipcam MCPA 500 Selective Herbicide or other disruptors of plant cell growth herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Sipcam Pacific Australia Pty Ltd accepts no liability for any losses that may result from the failure of Sipcam MCPA 500 Selective Herbicide to control resistant weeds.

MIXING AND APPLICATION: This product mixes readily with water. Half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly before adding the remaining water.

BOOM SPRAYING: Use 30L-120L water per hectare. Use a minimum of 170L water per hectare for Linseed.

AERIAL SPRAYING: Use 10L-90L water per hectare.

Equipment that has been used for this chemical should be well cleaned with a 2% solution of cloudy ammonia or equivalent followed by several clear water rinses before use for the application of other materials to sensitive plants.

LEGUME TOLERANCE: The use of MCPA is preferable to 2,4-D amine or ester in situations where legumes are undersown in the crop. The tolerance of pasture legumes to applied product can vary with species, variety, growing conditions, stage of growth and companion crop cover. DO NOT spray cereal crops undersown with lucerne as damage may result.

TANK MIXES: When tank mixing add the required amount of compatible product (and surfactant if recommended) when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly before spraying.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS: DO NOT apply under weather conditions, or from spraying equipment, which may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT: DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL: Store in the closed original container, in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or (preferably) pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. Break, crush or puncture and bury empty containers in a local authority landfill. If not available bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS: Harmful if swallowed. Attacks eyes. Will irritate the skin. When preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre (TEL: 131126). If swallowed, and if more than 15 minutes from a hospital, induce vomiting, preferably using Ipecac Syrup APF.

NOTICE TO BUYER

This product is not to be used for any purpose or in any way contrary to label instructions. The buyer by use or application of this product acknowledges so doing upon the terms and conditions agreed between Sipcam Pacific Australia Pty Limited and the buyer but not limited to the following.

To the extent permitted by law all conditions and other warranties or other rights of action which buyer or any other user may have against Sipcam Pacific Australia Pty Limited (Sipcam) or Seller are hereby excluded. Sipcam hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Sipcam or on its behalf unless it can be established that such information or advice was provided negligently and that the product has been used strictly as directed. Notwithstanding the generality of this notice Sipcam will not be held liable for any loss, injury or damage indirect or consequential arising from the sale, supply, use or application of this product. Sipcam's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefor.

BATCH No.

DOM:

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SIPCAM
MCPA 500
SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 500g/L MCPA
PRESENT AS THE DIMETHYLAMINE SALT

**FOR SELECTIVE CONTROL OF BROADLEAF WEEDS IN CEREALS,
LINSEED, PASTURES, SUGAR CANE AND TURF
AS SPECIFIED IN THE DIRECTIONS FOR USE TABLE**

SIPCAM PACIFIC AUSTRALIA PTY LTD
Suite 11, 23-31 Gheringhap Street
GEE LONG VIC 3220
ACN 073 176 888
TEL: 03 5223 3746

NRA Approval No.

DIRECTIONS FOR USE:

RESTRAINTS: DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

DO NOT apply if rain is likely within 6 hours.

1. FIELD CROPS

Situation & Crop	Weeds Controlled	State	Rate/Ha	Critical Comments
Wheat, Barley, Oats, Triticale	Refer to Weed Table	SA, Tas. Only	700mL-2.1L	Apply through the tillering stage after the crop plants have 5 leaves and before they commence jointing.
		Qld. only		Apply from early tillering (4 to 5 fully emerged main stem leaves plus one or more tillers) until before the boot stage (visible swelling of the head at the top of the main stem).
		NSW only		Apply 700mL rate after crop has reached 5 leaf stage and weeds marked E in weed table have 3-6 leaves. Higher rates after the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller.
		Vic. only	700mL-1.5L	Apply 700mL rate after crop has reached 2-3 leaf stage and weeds marked E in weed table have 2-4 leaves. Higher rate after the crop plants have 3 leaves to fully tillered.
		WA only	900mL-2L	Apply to crop from 5 leaf to Flag leaf emergence. (Zadoks 15-37)
Wheat, Barley, Triticale, Cereal Rye	Capeweed, Double Gee, Erodium, Mustard, Rapistrum, Wild Turnip, Wild Radish	WA only	400mL plus 350mL Flowable Diuron	CROP: 3-4 leaf stage. (Zadoks 13-14)
			500mL plus 400mL Flowable Diuron	CROP: 4-5 leaf stage (Zadoks 14-15) Apply by aircraft or boom with 20-100L/ha water, when most weeds have germinated and are in 2-5 leaf stage. Crop and weeds should be dry at time of application. Some temporary yellowing of crop may occur after application. Undersown sub-clovers may be slightly retarded. DO NOT apply to undersown medic.
	Amsinckia (Yellow Burr), Capeweed, Charlock, Deadnettle Double Gee (Spiny Emex, Three Corner Jack), Erodium sp., Mustard, Turnip Weed, White Ironweed (Corn Gromwell, Sheep Weed), Wild Radish, Wild Turnip	NSW, Vic only	500mL plus 500mL Flowable Diuron	Apply by boom spray or aircraft when the crop is at the 3-5 leaf stage (Vic only). Weeds must be in the 2-5 leaf stage Do not apply to undersown medic or lucerne. Do not use on Kamilaroi, Kite, Olympic or Shortim Wheat.
	Capeweed, Mustard, Three Corner Jack, Turnip Weed, Wild Radish, Wild Turnip, Ward's Weed, Common Ice Plant, Pimpernel	SA only	350mL plus 500mL Flowable Diuron	Do not use on very light sandy soils.
Wheat, Barley, Oats, Triticale	Soldier Thistle	SA only	700mL plus 200mL Lontrel L	Apply to crop at tillering stage after the crop has 3 leaves and before the crop commences jointing.
	Skeleton Weed		1L plus 500mL Lontrel L	Apply to rosettes from 5-15cm diameter when the crop is between the 5 leaf and late tillering growth stages. It is essential that all the Skeleton weed is emerged and minimum of 5cm diameter which is usually not before August This rate will provide control of skeleton weed until harvest and ;sill substantially reduce seed numbers the following season.
Linseed	Refer to Weed Table	Qld only	700mL-2.1L	Apply when crop is 10-15cm tall with at least 170L/ha of water. Do not spray after buds appear. Some wilting effect may be seen following application but crop usually recovers quickly.
		NSW, Vic, SA, Tas only	700mL-1.1L	Do not apply by aircraft

Sugar Cane	Blue Top, Chinese Burr, Flannel Weed, Gambia Pea, Bell Vine, Streaked Rattle Pod, Bindweed, Pink Convolvulus, Cupids Flower, Merremia Vine, Morning Glory	Qld only	1.4L	POST EMERGENT: Apply as a directed inter-row spray. Do not exceed rates with phenoxy sensitive varieties.
	As above plus Fat Hen, Noogoora Burr		2.1L	
	All of the above		10L	
Flax	Refer to Weed Table	Vic, Tas only	490mL	Apply to crop when 8-20cm high.

2. PASTURES, SEED CROPS & TURF

Situation & Crop	Weeds Controlled	State	Rate/Ha	Critical Comments
Grass Pastures	Refer to Weed Table	All States	700mL-4L	Apply to established grass based pastures. Damage may occur to legumes if present. See "Legume tolerance" below. NOTE: Capeweed, Paterson's Curse and Variegated Thistle may be poisonous to stock after spraying.
	Salvation Jane	SA only	1L plus 400mL Flowable Diuron	Use this mixture only if spraying is delayed until early winter. Rosettes should not exceed 20cm diameter.
	Soldier Thistle	SA only	700mL-2.1L	Use low rate on seedlings with no more than 2 true leaves. Use 100 litres of water for best results. If clover present do not exceed 600mL.
	Seedling and established Horehound	NSW only	3L	Apply in autumn when horehound plants are at the seedling stage. If sub clover or medic plants are present they will be damaged by this application.
Grass Seed Crops	Refer to Weed Table	Qld., SA Tas., Vic., WA only	700mL-4L	Apply to established grass seed crops.
Sub-Clover Seed Crops	Refer to Weed Table	Vic., SA only	700mL-1.5L	Do not exceed rate specified. Spray in autumn & early winter.
Pastures Spray/Graze Technique	Caltrop, Capeweed, Charlock, Mustards, Paterson's Curse, Shepherds Purse, Saffron, Slender, Spear or Variegated Thistle, Turnip Weed, Wild Radish, Wild Turnip	NSW, Vic only	350mL-1.4L	Spray actively growing 6-8 weeks old weeds. Introduce stock 7-10 days after spraying, preferably sheep as cattle are less effective. Use stocking rate at least 5 times heavier than normal until weeds have been reduced but before survival of desirable pasture species is threatened. Lucerne and medics may be damaged and should be grazed short before spraying Refer "Legume Tolerance". Precaution: .An increased quantity of poisonous plants may be eaten by stock using Spray/Graze, eg Caltrop, Capeweed, Paterson's Curse and Variegated Thistle and deaths could result from causes such as nitrate poisoning.
		Tas only	700mL	
	Refer to Weed Table.	WA only	925mL-1.85L	
	Cape Weed, Dandelion, Seedling Dock, Three Corner Jack, Geranium, Erodium, Mustard, Annual Thistles, Turnip Weed, Wild Radish, Wild Turnip.	SA only	500mL	
	Soldier Thistle		600mL	
				With Paterson's Curse preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock. Do not graze horses or pigs on Paterson's Curse. Observe Withholding Period of at least 7 days after spraying.

Turf	Refer to Weed Table	All States	500mL or 5mL per 100 sq. metres	Newly laid - Delay until well established or at least two months old. Apply in high volume to actively growing weeds. See "Legume Tolerance". Buffalo grass is more easily damaged than other grasses.
			1.4-2.8L or 14-28mL per 100 sq. metres	Established - Apply in high volume to actively growing weeds. Do not mow for 2 days before application. Some transitory damage may occur to fine turf grasses and strawberry clover. See "Legume Tolerance". Buffalo grass is more easily damaged than other grasses.

3. SPOT SPRAYING

Situation & Crop	Weeds Controlled	State	Mixing Rates /Comments
High Volume Spraying	Refer to Weed Table	All States	Add 1/10th of rate on Weed Table to 150 litres of water. Each 150 litres of mix will cover 1000m ² (1/10th Ha) eg. If rate in Weed Table is 1.4L use 140mL/150L water.
Knapsack Application	Refer to Weed Table	All States	Add 1/100th of rate on Weed Table to 10 litres of water. Each 10 litres of mix will cover 100m ² (1/100Ha.) eg. if rate in Weed Table is 1.4L use 14mL/10L Water.

4. NON CROP AREAS, FALLOW LAND, INDUSTRIAL & COMMERCIAL AREAS, RIGHTS-OF-WAY

Situation & Crop	Weeds Controlled	State	Rate/Ha	Critical Comments
Non Crop Areas, Fallow Land, Industrial and Commercial Areas, Rights of Way	Refer to Weed Table	All States	Refer to Weed Table	Apply at rates shown in Weed Table when weeds only are present and no crop or pasture is involved.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD:

CEREALS, PASTURE - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS:

Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavourable. The best conditions are when the soil is moist, weather fine and rain unlikely within 6 hours of application.

DO NOT spray crops or weeds outside the stages indicated in the Table as damage, loss of yield or inadequate weed control may result.

Resistant Weeds Warning:

Sipcam MCPA 500 Selective Herbicide is a member of the Phenoxy group of herbicides. Sipcam MCPA 500 Selective Herbicide has the disruptors of plant cell growth mode of action. For weed resistance management Sipcam MCPA 500 Selective Herbicide is a Group I herbicide. Some naturally-occurring weed biotypes resistant to Sipcam MCPA 500 Selective Herbicide and other disruptors of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Sipcam MCPA 500 Selective Herbicide or other disruptors of plant cell growth herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Sipcam Pacific Australia Pty Ltd accepts no liability for any losses that may result from the failure of Sipcam MCPA 500 Selective Herbicide to control resistant weeds.

MIXING AND APPLICATION: This product mixes readily with water. Half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly before adding the remaining water.

BOOM SPRAYING: Use 30L-120L water per hectare. Use a minimum of 170L water per hectare for Linseed.

AERIAL SPRAYING: Use 10L-90L water per hectare.

Equipment that has been used for this chemical should be well cleaned with a 2% solution of cloudy ammonia or equivalent followed by several clear water rinses before use for the application of other materials to sensitive plants.

LEGUME TOLERANCE: The use of MCPA is preferable to 2,4-D amine or ester in situations where legumes are undersown in the crop. The tolerance of pasture legumes to applied product can vary with species, variety, growing conditions, stage of growth and companion crop cover. DO NOT spray cereal crops undersown with lucerne as damage may result.

TANK MIXES: When tank mixing add the required amount of compatible product (and surfactant if recommended) when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly before spraying.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS: DO NOT apply under weather conditions, or from spraying equipment, which may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT: DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL: Store in the closed original container, in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or (preferably) pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. Break, crush or puncture and bury empty containers in a local authority landfill. If not available bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS: Harmful if swallowed. Attacks eyes. Will irritate the skin. When preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre (TEL: 131126). If swallowed, and if more than 15 minutes from a hospital, induce vomiting, preferably using Ipecac Syrup APF.

NOTICE TO BUYER

This product is not to be used for any purpose or in any way contrary to label instructions. The buyer by use or application of this product acknowledges so doing upon the terms and conditions agreed between Sipcam Pacific Australia Pty Limited and the buyer but not limited to the following.

To the extent permitted by law all conditions and other warranties or other rights of action which buyer or any other user may have against Sipcam Pacific Australia Pty Limited (Sipcam) or Seller are hereby excluded.

Sipcam hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Sipcam or on its behalf unless it can be established

that such information or advice was provided negligently and that the product has been used strictly as directed.

Notwithstanding the generality of this notice Sipcam will not be held liable for any loss, injury or damage indirect or consequential arising from the sale, supply, use or application of this product. Sipcam's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefor.

WEED TABLE:

Where weeds are to be sprayed in a crop or pasture, use only the rate given for the particular crop or situation indicated under the Directions for Use table.

In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed.

The rates listed in the Weed Table below are for use where weeds only are present and no crop or pasture is involved. Use at these rates in a crop or pasture other than for spot spraying may cause damage.

This product should only be used in those States where a rate or range of rates indicated for the particular weeds listed.

Application Rate: Per Hectare							
Weeds Controlled	Vic	NSW	SA	Tas	Qld	WA	Critical Comments
Ball Mustard	-	-	1.7L	-	-	-	Apply at rosette stage
Bathurst Burr	2L	1-2L	2L	2L	2L	2L	Spray young seedlings only.
Black Bindweed	-	1.4-2L	-	-	1.4-2L	-	Apply at young stage prior to flowering.
Californian Burr	-	1-2L	-	2.2L	-	-	Spray young seedlings only.
Cape Tulip	-	-	-	-	-	2L	Apply just prior to flowering. Gives mild suppression only
Capeweed	-	2.1L	2.1L	2.1L	2.1L	-	Apply at seedling stage.
Carrot Weed	2L	-	-	-	2L	-	Apply at seedling stage.
Charlock	700mL E	1L E	700mL	1-2.2L	700mL	-	Spray up to rosette stage.
Cutleaf Mignonette	-	-	2.1L	-	-	-	Apply at young rosette stage.
Dandelion	2L	2L	2L	2L	2L	2L	Apply at young rosette stage
Deadnettle	-	2.1L	-	-	2.1L	-	Moderately susceptible. Apply at seedling stage.
Docks	1.5L	-	2.1L	2.1L	-	-	Apply at rosette stage and before flower heads emerge. Only the top growth of old established plants will be affected.
Fat Hen	2L	1-2L	-	2L	2L	-	Spray up to rosette stage.
Fennel	2.7L	2.7L	2.7L	2.7L	2.7L	-	Apply at young stage.
Field Bindweed	2L	-	2L	2L	2L	2L	Apply at bud stage, repeating over several seasons.
Field Cress	-	-	-	2.2L	-	-	Apply at young stage.
Flatweed or Catsear	-	-	-	2.2L	-	1.4L	Apply when weeds are 10-15 cm high.
Fumitory (red)	1.4L	1.4L	1.4L	-	1.4L	-	Apply at young stage prior to flowering.
Hoary Cress	4.5L	-	2.1L	-	4.5L	-	Apply before flowering.
Horehound	-	3L	2.1L	-	-	-	Apply at rosette stage.
Hedge Mustard	1.1L E	-	700mL	700mL	700mL	1.25L	Apply at rosette stage.
Hexham Scent or Mellilotus	1.5L	1.2L	1.4L	-	1.4L	-	Apply at young seedling stage.
Lesser Swinecress	-	-	-	2.2L	-	-	Apply at young stage.
Lincoln Weed	-	-	700mL	-	-	-	Apply at rosette stage..
London Rocket	-	-	-	-	-	1.25L	
Lupins	-	700mL-1.5L	-	-	-	1.25L	Spray up to 10cm high.

Mintweed	2L	2L	-	-	2L	-	Spray young seedlings only.
Mustards	1.1L E	1L E	1.1L	1.1L	-	1.25L	Spray up to rosette stage.
Noogoora Burr	2L	1-2L	2L	-	2L	2L	Spray young seedlings only.
Opium Poppy	-	-	-	2.2L	-	-	Apply at young stage.
Paterson's Curse	-	1-1.5L	-	-	4L	-	Apply early rosette stage.
Pimpernel	-	-	700mL	-	-	-	Apply at rosette stage.
Plantains	-	2L	-	2L	2L	-	Apply at young stage.
Rapeseed	-	1-1.5L	-	-	-	-	Spray up to rosette stage.
Rough Poppy	-	700mL- 1.5L	-	-	-	-	Spray up to rosette stage.
Safflower	-	700mL-1L	-	-	-	-	Spray up to rosette stage.
Shepherds Purse	-	-	-	2.2L	-	-	Apply at rosette stage.
Skeleton Weed	-	1.5-2L	-	-	-	-	Spray rosettes before aerial growth commences.
Sorrel	-	-	2.1L	-	-	-	Apply to rosette stage.
Stinkwort	2L	1-1.5L	2L	-	-	2L	Apply when weeds are 10-15cm high.
Sunflower (seedlings)	-	1-1.5L	-	-	-	-	Spray up to rosette stage.
Thistles:							
-Artichoke	1.6L	-	2.1L	-	2L	700mL	Apply at rosette stage.
-Nodding	-	2.5L	-	-	-	-	Spray early rosettes. Repeat spray required in following Spring or Autumn.
-Saffron	1.5-2.5L	1-2L	1.4-2.5L	1.4-2L	1.4-2.5L	1.5- 2.5L	Spray up to rosette stage. (September in WA)
Scotch or Cotton	2.2L	-	700mL- 1.4L	2.2L	-	-	Apply at seedling stage
Slender	3L	1-2L	1L	3L	-	3L	Spray young seedlings only.
Soldier	1.1L	-	1.1L	-	-	-	Spray young seedlings. Use higher rate for larger weeds.
Spear	-	11.5-2L.5- 2L	-	2.2L	-	-	Apply at rosette stage.
Star	-	-	1-1.4L	-	-	-	Apply at seedling stage. Use higher rate for larger weeds.
Variegated	1.5L	1-2L	700mL- 1.4L	1.5L	1.5L	1.5L	Spray young seedlings only. Use 2L/ha (SA) for well developed weeds.
Turnip Weed	1.1L E	1L E	1L	-	-	-	Spray up to rosette stage.
Wild Artichoke	-	-	2.1L	-	-	-	Apply to rosette stage.
Wild Radish	1.4L	1L E	1.4L	2L	700mL- 1.1L	1.2L	Spray up to rosette stage.
Wild Turnip	1.1L E	1L E	700mL	2L	700mL- 1.1L	900mL	Spray up to rosette stage.