

Main panel

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SIPCAM
LVE MCPA* 500
HERBICIDE

ACTIVE CONSTITUENT: 500g/L MCPA
PRESENT AS THE 2-ETHYL HEXYL ESTER

GROUP	I	HERBICIDE
-------	----------	-----------

FOR SELECTIVE CONTROL OF BROADLEAF WEEDS IN
CEREALS AND PASTURES
AS SPECIFIED IN THE DIRECTIONS FOR USE TABLE

CONTENTS: 20, 200 Litres

SIPCAM PACIFIC AUSTRALIA PTY LTD
Suite 11, 23 - 31 Gheringhap Street
GEE LONG VIC 3220
ACN 073 176 888

NRA Approval No.

Ancillary panel 1

DIRECTIONS FOR USE:

SITUATION	WEEDS CONTROLLED	STATE	RATE/ha	CRITICAL COMMENTS
Wheat	Refer Weed Table	All States	350mL-2.1L	Apply from 5 true leaf through to the emergence of the flag leaf. Use lower rates when crop is 2-3 leaf stage. Use high rates after 5 leaf stage.
Oats, Barley, Triticale and Cereal Rye			350mL-1.6L	Apply from 5 true leaf through to the emergence of the flag leaf. Use lower rates when crop is 2-3 leaf stage. Use high rates after 5 leaf stage. Only apply after 5 leaf stage on barley.
Pastures		Vic, NSW, Qld SA, Tas only	350mL-2.1L	Observe legume tolerance rates under General Instructions.

WEED TABLE: CONTROL RECOMMENDATIONS FOR SPECIFIC WEEDS

Weeds Controlled	Rate/ha	Critical Comments
Bathurst Burr	1.6L	Spray young seedlings only. Only moderately susceptible
Capeweed	1-1.7L	Spray young rosettes. Becomes more tolerant with age.
Charlock	350mL-1.6L	Spray in rosette stage. Young rosettes very susceptible.
Dandelion	1.6L	Spray in early rosette stage.
Fat Hen	1.6L	Spray young rosettes. Becomes more tolerant with age.
Fumitory (NOT TAS)	1.1L	Spray up to first true leaf stage. Only moderately susceptible.
Hedge Mustards	560mL-1L	Apply at rosette stage.
London Rocket	1.1L	Spray in early rosette stage.
Noogoora Burr	1.6L	Spray young seedlings. Becomes more tolerant with age.
Skeleton Weed	1.1-1.6L (NSW only)	Spray in rosette stage only. Only moderately susceptible.
Stinkwort	1.6L	Apply when weeds are 10-15cm high.
Thistles		
-Saffron	1.1-1.9L	Spray young rosettes before leaves become spiny.
-Scotch	700mL-1.7L	Spray in early rosette stage.
-Slender	2.1L	Spray in rosette stage only. Only moderately susceptible.
-Spear	1.5-2.1L	Spray in early rosette stage. Only moderately susceptible.
-Variegated	840mL-1.1L	Spray at pre-cabbage stage.
Wild Radish	840mL-1.6L	Spray in rosette stage. Low rates for younger weeds.
Wild Sage	1.1L	Spray young seedlings.
Wild Turnip	350mL-1.5L	Spray young rosettes. Low rates for younger weeds.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD:

CEREALS, PASTURE - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS:

Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavourable. The best conditions are when the soil is moist, weather fine and rain unlikely within 6 hours of application.

DO NOT spray crops or weeds outside the stages indicated in the Table as damage, loss of yield or inadequate weed control may result. DO NOT spray in high winds.

LEGUME TOLERANCE:

The relative tolerance of legumes is:

Lucerne	not tolerant
Medics	low tolerance
Sub-clover	moderate tolerance
White clover	moderate tolerance
Strawberry clover	high tolerance

DO NOT spray cereal crops undersown with lucerne as damage may result. The tolerance of pasture legumes to applied product can vary with species, variety, growing conditions, stage of growth and companion crop cover.

Resistant Weeds Warning:

SIPCAM LVE MCPA 500 Herbicide is a member of the Phenoxy group of herbicides. SIPCAM LVE MCPA 500 Herbicide has the disruptors of plant cell growth mode of action. For weed resistance management SIPCAM LVE MCPA 500 Herbicide is a Group I herbicide. Some naturally-occurring weed biotypes resistant to SIPCAM LVE MCPA 500 Herbicide and other disruptors of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by SIPCAM LVE MCPA 500 Herbicide or other disruptors of plant cell growth herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Sipcam Pacific Australia Pty Ltd accepts no liability for any losses that may result from the failure of SIPCAM LVE MCPA 500 Herbicide to control resistant weeds.

MIXING AND APPLICATION: This product mixes readily with water. Half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly before adding the remaining water.

BOOM SPRAYING: Use 30L-120L water per hectare.

AERIAL SPRAYING: Use 20L-30L water per hectare.

HIGH VOLUME SPRAYING: Use 200mL per 100L water for easy to kill weeds, eg Mustards, Wild Radish, and Wild Turnip. Use 400mL per 100L water for difficult to control weeds eg Slender Thistle.

CEREALS: Apply 50-200L (WA 50-100L) water per hectare. Use higher volumes for heavy weed infestation or dense crop cover.

Ancillary panel 1 continued

Equipment that has been used for this chemical should be well cleaned with a 2% solution of cloudy ammonia or equivalent followed by several clear water rinses before use for the application of other materials to sensitive plants.

TANK MIXES: When tank mixing add the required amount of compatible product (and surfactant if recommended) when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly before spraying.

COMPATIBILITY: SIPCAM LVE MCPA 500 Herbicide is compatible with bromoxynil, chlorsulfuron, dicamba, diclofop methyl, glyphosate and metsulfuron methyl. Test all mixtures before using commercial quantities.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS: DO NOT apply under weather conditions, or from spraying equipment, which may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers, or waterways with the chemical or used containers.

STORAGE AND DISPOSAL: Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS: Harmful if absorbed by skin contact or inhaled or swallowed. Will irritate the eyes and skin. Do not inhale vapour or spray mist. Repeated exposure may cause allergic disorders. When opening container and preparing spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and face shield or goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist, a washable hat, and elbow-length PVC gloves. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 1126. If swallowed, and if more than 15 minutes from a hospital, induce vomiting, preferably using Ipecac Syrup APF. Additional information is listed in the Material Safety Data Sheet.

NOTICE TO BUYER

This product is not to be used for any purpose or in any way contrary to label instructions. The buyer by use or application of this product acknowledges so doing upon the terms and conditions agreed between Sipcaml Pacific Australia Pty Limited and the buyer but not limited to the following.

To the extent permitted by law all conditions and other warranties or other rights of action which buyer or any other user may have against Sipcaml Pacific Australia Pty Limited (Sipcaml) or Seller are hereby excluded. Sipcaml hereby gives notice to

buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Sipcam or on its behalf unless it can be established that such information or advice was provided negligently and that the product has been used strictly as directed. Notwithstanding the generality of this notice Sipcam will not be held liable for any loss, injury or damage indirect or consequential arising from the sale, supply, use or application of this product. Sipcam's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefor.