

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Mesurol[®]

SNAIL AND SLUG BAIT

Active Constituent: 20 g/kg METHIOCARB (an anticholinesterase compound)

CONTAINS BITREX[™]

For the control of snails and slugs in berry crops, cereals, gardens, nurseries, oilseed crops, orchards, pastures and vegetable crops and for false wireworm beetle on sunflowers

GENERAL INSTRUCTIONS

Export of treated produce

Growers should note that MRLs or import tolerances do not exist in all markets for edible produce treated with Mesurol. If you are growing edible produce for export, please check with Bayer CropScience Pty Ltd for the latest information on MRLs and import tolerances BEFORE using Mesurol.

PRECAUTION

Avoid application of pellets to foliage of edible crops.

Store in original container. Lock in a safe place preventing access of children, animals, poultry or ducks. Keep away from domestic pets. Keep away from dogs. Dogs find this bait attractive and it may kill them. If pets are poisoned, contact a veterinary surgeon.

PROTECTION OF LIVESTOCK

After filling applicator, clean up spilled pellets so that they are not eaten by animals and birds. Dogs find this bait attractive and if ingested it may kill them. Do not heap as heaping is unnecessary and such heaps could be eaten by dogs. If a dog eats pellets take it to a veterinarian immediately. Do not give dogs Ipecac Syrup. Do not treat areas on which poultry graze.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used container.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Lock in a safe place preventing access of children, animals, poultry or ducks. Keep away from domestic pets. Ensure plastic bag is completely empty before disposal. Puncture and bury empty bag in a local authority landfill. If no landfill is available, bury the bag below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty bags and product should not be burnt.

SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with eyes and skin. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. Obtain an emergency supply of atropine tablets 0.6 mg.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (telephone 13 11 26). If swallowed, give one atropine tablet every 5 minutes until dryness of the mouth occurs. If poisoned by skin absorption or through lungs, remove any contaminated clothing, wash skin thoroughly and give atropine tablets as above. Get to a doctor or hospital quickly.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from www.bayercropscience.com.au.

EXCLUSION OF LIABILITY

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

APVMA Approval No.: 33274/0304

Mesurol[®] is a registered trademark of Bayer

FOR 24 HOUR SPECIALIST ADVICE
IN EMERGENCY ONLY
PHONE 1800 033 111

DIRECTIONS FOR USE

CROP/SITUATION	PEST	RATE	CRITICAL COMMENTS
Berry crops (including grapevines), Cereals, Gardens, Nurseries, Oilseed crops, Orchards, Pastures, Vegetable crops	Common garden snail, slugs	5.5 kg/ha or 11-22 kg/ha	For most infestations apply low rate. For heavy infestations or where pasture is tall or dense apply higher rate. Scatter bait evenly onto ground where snails or slugs occur. DO NOT HEAP PELLETS. When applying in vegetable crops, ensure pellets do not become lodged in plant foliage. Equipment such as fertiliser spinners, combines or sod seeders are satisfactory for spreading the pellets and can easily be calibrated to apply 5.5 kg/ha. Gloves should be worn when pellets are spread by hand.
	White Italian snail, white snail (Not Qld)		
Sunflowers (Qld, SA only)	False wireworm beetle	2.5 kg/ha (10 pellets/m ²)	Apply Mesurol 1-3 days after sowing. Scatter bait evenly onto ground where false wireworm beetles occur. DO NOT HEAP PELLETS.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

Edible crops: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION

Treated areas: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION