

DIRECTIONS FOR USE

CROP/SITUATION	PEST	RATE	CRITICAL COMMENTS
Berry crops (including grapevines), Cereals, Gardens, Nurseries, Oilseed crops, Orchards, Pastures, Vegetable crops	Common garden snail, slugs	5.5 kg/ha or	For most infestations apply low rate. For heavy infestations or where pasture is tall or dense apply higher rate. Scatter bait evenly onto ground where snails or slugs occur. DO NOT HEAP PELLETS. When applying in vegetable crops, ensure pellets do not become lodged in plant foliage. Equipment such as fertiliser spinners, combines or sod seeders are satisfactory for spreading the pellets and can easily be calibrated to apply 5.5 kg/ha. Gloves should be worn when pellets are spread by hand.
	White Italian snail, white snail (Not Qld)	11-22 kg/ha	
Sunflowers (Qld, SA only)	False wireworm beetle	2.5 kg/ha (10 pellets/m ²)	Apply MesuroI 1-3 days after sowing. Scatter bait evenly onto ground where false wireworm beetles occur. DO NOT HEAP PELLETS.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

Edible crops: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION

Treated areas: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION