

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



**SNAIL AND SLUG BAIT**

**Active Constituent: 20 g/kg METHIOCARB (an anticholinesterase compound)**

**CONTAINS BITREX™**

**For the control of snails and slugs in berry crops, cereals, gardens, nurseries, oilseed crops, orchards, pastures and vegetable crops and for false wireworm beetle on sunflowers**

**GENERAL INSTRUCTIONS**

**Export of treated produce**

Growers should note that MRLs or import tolerances do not exist in all markets for edible produce treated with Mesuro!. If you are growing edible produce for export, please check with Bayer CropScience Pty Ltd for the latest information on MRLs and import tolerances BEFORE using Mesuro!.

**PRECAUTION**

Avoid application of pellets to foliage of edible crops.

Store in original container. Lock in a safe place preventing access of children, animals, poultry or ducks. Keep away from domestic pets. Keep away from dogs. Dogs find this bait attractive and it may kill them. If pets are poisoned, contact a veterinary surgeon.

**PROTECTION OF LIVESTOCK**

After filling applicator, clean up spilled pellets so that they are not eaten by animals and birds. Dogs find this bait attractive and if ingested it may kill them. Do not heap as heaping is unnecessary and such heaps could be eaten by dogs. If a dog eats pellets take it to a veterinarian immediately. Do not give dogs Ipecac Syrup. Do not treat areas on which poultry graze.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used container.

**STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Lock in a safe place preventing access of children, animals, poultry or ducks. Keep away from domestic pets.

Ensure plastic bag is completely empty before disposal. Puncture and bury empty bag in a local authority landfill. If no landfill is available, bury the bag below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty bags and product should not be burnt.

**SAFETY DIRECTIONS**

Product is poisonous if swallowed. May irritate the eyes. Avoid contact with eyes and skin. When using the product, wear cotton overalls buttoned to the neck and wrist and a washable hat. If applying by hand, wear elbow-length PVC gloves. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

**FIRST AID**

If swallowed, splashed on skin or in eyes, or inhaled, contact a Poisons Information Centre (Phone Australia 131126) or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised. Give atropine if instructed.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet, which can be obtained from [www.bayercropscience.com.au](http://www.bayercropscience.com.au).

**EXCLUSION OF LIABILITY**

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

APVMA Approval No.: 33274/1209

Mesuro!® is a registered trademark of Bayer

FOR 24 HOUR SPECIALIST ADVICE  
IN EMERGENCY ONLY  
PHONE 1800 033 111

**DIRECTIONS FOR USE**

<b>CROP/SITUATION</b>	<b>PEST</b>	<b>RATE</b>	<b>CRITICAL COMMENTS</b>
Berry crops (including grapevines), Cereals, Gardens, Nurseries, Oilseed crops, Orchards, Pastures, Vegetable crops	Common garden snail, slugs	5.5 kg/ha or	For most infestations apply low rate. For heavy infestations or where pasture is tall or dense apply higher rate. Scatter bait evenly onto ground where snails or slugs occur. <b>DO NOT HEAP PELLETS.</b> When applying in vegetable crops, ensure pellets do not become lodged in plant foliage.  Equipment such as fertiliser spinners, combines or sod seeders are satisfactory for spreading the pellets and can easily be calibrated to apply 5.5 kg/ha. Gloves should be worn when pellets are spread by hand.
	White Italian snail, white snail (Not Qld)	11-22 kg/ha	
Sunflowers (Qld, SA only)	False wireworm beetle	2.5 kg/ha (10 pellets/m <sup>2</sup> )	Apply Mesuroi 1-3 days after sowing. Scatter bait evenly onto ground where false wireworm beetles occur. <b>DO NOT HEAP PELLETS.</b>

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIODS:**

**Edible crops: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION**

**Treated areas: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION**