

**CAUTION**

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Bouncer[®]

Herbicide

ACTIVE CONSTITUENT: 720 g/L METOLACHLORGROUP **K** HERBICIDE

Controls certain annual grasses and broadleaf weeds in certain crops as specified in the directions for use

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

© Bouncer is a registered trademark of Nufarm Australia Limited

nufarm.com.au

Nufarm Australia Limited
ACN 004 377 780
103-105 Pipe Road
Laverton North Victoria 3026
Tel: (03) 9282 1000
Fax: (03) 9282 1001

Contents: 1 Litres
Contents: 5 Litres
Contents: 10 Litres
Contents: 20 Litres
Contents: 110 Litres
Contents: 200 Litres
Contents: 1000 Litres

APVMA Approval No.: 52024/59176

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product

For refillable containers:

Store in the closed, original container, in a cool, well ventilated area away from children, animals, food, feedstuffs, seed and fertilizers. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Envirodrum[®] 110L Mini Bulk Returnable Container:

Store the original sealed Envirodrum in a cool well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Nufarm Australia Limited.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Tel: Australia 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier or the Nufarm website - www.nufarm.com.au

In case of emergency: Phone 1800 033 498 Ask for shift supervisor. Toll free 24 hours.



CONDITIONS OF SALE

"Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute." BN / DOM / Barcode / drumMUSTER logo / Envirodrum /



ENVIRONMENTALLY HAZARDOUS SUBSTANCES, LIQUID, N.O.S. (CONTAINS METOLACHLOR)
UN No: 3082
PACKING GROUP: III
HAZCHEM: 2 Z
IN AN EMERGENCY DIAL: 000 POLICE OR FIRE BRIGADE
IN AN EMERGENCY, SPECIALIST ADVICE NUFARM AUSTRALIA LIMITED 1800 033 498

"NOTE: MANUFACTURED IN AUSTRALIA – DG CLASS 9 MARKING ONLY REQUIRED FOR SHIPMENT TO TASMANIA – THIS IS DONE BY NUFARM LOGISTICS"



CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Bouncer®

Herbicide

ACTIVE CONSTITUENT: 720 g/L METOLACHLOR

GROUP **K** HERBICIDE

Controls certain annual grasses and broadleaf weeds in certain crops as specified in the directions for use

READ COMPLETE DIRECTIONS FOR USE BEFORE USING THIS PRODUCT.

® Bouncer is a registered trademark of Nufarm Australia Limited.

nufarm.com.au

APVMA Approval No.: 52024/59176

DIRECTIONS FOR USE

CROP	WEEDS CONTROLLED	STATE	RATE	CRITICAL COMMENTS
Barley, Oats	Suppression of Annual ryegrass	NSW, ACT, Vic, Tas, SA WA only	750mL-1L/ha	Apply to moist soils immediately after planting and before crop and weeds emerge. Use the lower rate on light soils. Application should be made to a moist seed bed and sufficient rain to thoroughly wet the top 3-4cm of soil should occur within 10 days after spraying. Damage may occur if crop is sown too shallow (less than 4cm). Damage can occur on light sandy soil types after heavy rain. See General instructions for use in zero-tillage systems.
Barley, Oats, Triticale, Wheat	Toad rush	NSW, ACT, Vic, Tas, SA WA only	300-500mL/ha	Apply at or immediately after planting and before crop and weeds emerge. Application should be made to a moist seed bed and sufficient rain to thoroughly wet the top 3-4cm of soil should occur within 10 days after spraying. Use rates towards the higher end of the range in wet areas or when longer residual control is required or in situations where crop competition is minimal. When mixed with Nugran® (Triasulfuron 750) apply pre-sowing and incorporate by sowing operation (for Wheat only). See General instructions for use in zero-tillage systems.
Broccoli, Brussels sprouts, Cabbages, Cauliflowers	Barnyard grass, Blackberry nightshade, Chickweed (suppression), Deadnettle, Fat hen, Lovegrass, Pigeon grass, Pigweed (suppression), Potato weed, Shepherd's purse, Stinging nettle, Summer grass, Wireweed (suppression)	NSW, ACT, Vic, Qld, Tas, NT only	3-4L/ha	Apply immediately after transplanting. Sufficient irrigation to wet the top 3-4cm of soil should be applied within 24 hours. Use rates towards the higher end of the range where Blackberry nightshade, Chickweed, Fat hen, Lovegrass, Pigeon grass, Pigweed, Potato weed and Wireweed are the major problem. See warning – Brassica crops
	Crowsfoot grass, Fierce thornapple (suppression), Liverseed grass, Winter grass	NSW, ACT only		Apply immediately after transplanting. Sufficient irrigation to wet the top 3-4cm of soil should be applied within 24 hours. Use rates towards the higher end of the range where Apple of Peru, Common sowthistle, Fierce thornapple, Liverseed grass, Redroot amaranth, Twiggly turnip and Nettleleaf goosefoot are the major problems.
	Apple of Peru, Awnless barnyard grass, Fierce thornapple (suppression) Liverseed grass, Yellow weed, Redroot amaranth (suppression)	Qld only		WARNING: Qld, only stunting and reduced yield may occur in broccoli in red soils and light alluvial soils.
	Annual ryegrass, Common cotula, Common sowthistle (suppression), Crowsfoot grass, Fiddle dock, Italian ryegrass, Liverseed grass, Nettleleaf goosefoot (suppression), Twiggly turnip (suppression), Winter grass.	Vic, Tas only		See Warning – Brassica crops

CROP	WEEDS CONTROLLED	STATE	RATE	CRITICAL COMMENTS
Broccoli, Brussel sprouts, Cabbages, Cauliflower and Chinese Cabbage	Barnyard grass, Blackberry nightshade, Chickweed (suppression), Deadnettle, Fat hen, Lovegrass, Pigeon grass, Pigweed (suppression), Potato weed, Shepherd's purse, Stinging nettle, Crab grass, Wireweed (suppression), Winter grass, Fiddle dock, Goosefoot (suppression)	WA only	3-4L/ha	On soils of extremely low organic matter use the lower rate. Apply immediately after transplanting. Sufficient irrigation to wet the top 3-4cm of soil (but not greater than 25mm irrigation) should be applied within 24 hours. Soil type permitting, use rates towards the higher end of the range where Blackberry nightshade, Chickweed, Fat hen, Lovegrass, Pigeon grass, Pigweed, and Potato weed are the major problem. See warning – Brassica crops
Canola	Toad rush	NSW, ACT, SA, Vic, Tas, Qld,	300-500mL/ha	Apply at or immediately after planting and before crops and weeds emerge. Apply to moist soil, use rates towards the higher end of the range where longer residual control is required.
		WA Only		Apply before crops and weeds emerge. Apply to moist soil. To avoid crop injury use rates towards the lower end of the range on light sandy soils.
Cotton (Including Roundup Ready Flex cotton and Liberty Link cotton)	Barnyard grass, Crowsfoot grass, Liverseed grass, Lovegrass, Pigeon grass, Summer grass, Wandering Jew	NSW, WA, Qld only	2L/ha	Pre-emergent. Apply before, at, or immediately after planting and before crops and weeds have germinated using a low volume boom sprayer applying 60-120L of mixture per hectare. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3-4cm of soil. If rain does not occur or irrigation is unavailable incorporate chemical to a depth of 3-4cm. Use rates towards the higher end of the range on heavy soils or where a heavy grass population is expected. DO NOT throw untreated soil onto treated areas as this will reduce weed control. DO NOT use on soils where the sand content exceeds 70%. Use lower rate where the sand content of the soil is within the range 50%-70%. See Warning – Cotton
	Awnless barnyard grass (<i>Echinochloa colona</i>) Barnyard grass (<i>Echinochloa crus-galli</i>) Dinebra (<i>Dinebra retroflexa</i>), Crowsfoot grass, Liverseed grass, Lovegrass, Pigeon grass, Summer grass, Wandering Jew		2.0 L/ha	Over the top or directed spray applications: Apply as a standalone over-the-top or directed spray application to a weed-free soil surface, using a boom sprayer applying 85-120L of mixture per hectare. Apply only one application of Bouncer Herbicide per crop. Apply from the 4 node to 18 node crop growth stage. For maximum efficacy ensure even distribution of spray across the furrow. Use a directed spray if shading by the crop is likely to effect herbicide placement. Some minor transient leaf spotting or leaf scorching may occur after application. To avoid any leaf spotting or leaf scorching use a directed or shielded spray. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3-4cm of soil. If rain does not occur or irrigation is unavailable to incorporate reduced residual control may occur.
Roundup Ready Flex cotton			2.0 L/ha + 1.0-1.5 kg/L Roundup Ready Herbicide with Plantshield	Apply as an over-the-top or directed spray application when weeds are present. Apply only one application of Bouncer Herbicide per crop.

CROP	WEEDS CONTROLLED	STATE	RATE	CRITICAL COMMENTS
Green beans Navy beans	Blackberry nightshade, Common sowthistle, Fat hen, Potato weed, Bamyard grass, Crowsfoot grass, Liverseed grass, Lovegrass, Pigeon grass, Summer grass and suppression of Pigweed.	NSW, ACT, Qld, Vic, SA only	3-4L/ha	Pre-emergent. Apply before, at or immediately after planting and before crops and weeds have germinated. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3-4cm of soil. If rain does not occur or irrigation is unavailable incorporate chemical to a depth of 3-4cm. DO NOT throw untreated soil onto treated areas as this will reduce weed control.
	Powell's amaranth	Tas only	4L/ha	Pre-emergent. Apply before, at or immediately after planting and before crops and weeds have germinated. For sufficient Amaranth control the BOUNCER application should be followed by 1L/ha Basagran® (2 leaf Amaranth) plus 2 L/ha Basagran® (4 leaf Amaranth). Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3-4cm of soil. If rain does not occur or irrigation is unavailable incorporate chemical to a depth of 3-4cm. DO NOT throw untreated soil onto treated areas as this will reduce weed control.
Maize (not Waxy maize) Sweet com	Bamyard grass, Crowsfoot grass, Liverseed grass, Lovegrass, Pigeon grass, Summer grass, Wandering Jew.	NSW, ACT, Vic, Qld, SA, NT, Tas only	2-4L/ha	Pre-emergent. Apply before, at, or immediately after planting and before crops and weeds have germinated using a low volume boom sprayer applying 60-120L of mixture per hectare. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3-4cm of soil. If rain does not occur or irrigation is unavailable incorporate chemical to a depth of 3-4cm. Use rates towards the higher end of the range on heavy soils or where a heavy grass population is expected. DO NOT throw untreated soil onto treated areas as this will reduce weed control. DO NOT use on soils where the sand content exceeds 70%. Use lower rate where the sand content of the soil is within the range 50%-70%.
	Blackberry nightshade, Caltrop, Common thornapple, Mintweed, Needle burr, Noogoora burr, Pigweed, Redroot amaranth, Slim amaranth		2-4L/ha plus 2.5L/ha of Flowable Nu-trazine Liquid	
Pastures – Trikkala Subclover, Halfa white clover, Maral Persian clover, Paradana balansa clover, Redquin red clover, Phalaris and Cocksfoot. Frontier balansa clover	Toad rush	Vic, ACT, NSW, NT, SA only WA only	400-500mL/ha	Apply at or immediately after planting and before pasture species and weeds emerge. Application should be made to a moist seed bed. Sufficient rain to thoroughly wet the top 3-4cm of soil should occur within 10 days after spraying. Use rates towards the higher end of the range in wet areas or when longer residual control is required. DO NOT use in soils where prolonged waterlogging is anticipated otherwise unacceptable crop retardation will occur. Avoid damage to pasture species by avoiding overlaps and double spraying in headlands where White, Persian and Balansa clovers are sown and if the 500mL rate is being used.
Peanuts Soybeans Sunflowers	Bamyard grass, Crowsfoot grass, Liverseed grass, Lovegrass, Pigeon grass, Summer grass, Wandering Jew.	Qld, NSW, NT only NSW, ACT, Vic, Qld, SA, NT only	2-4L/ha	Apply before pre-irrigation where soil crusting is likely. Apply before, at, or immediately after planting and before weeds germinate. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3-4cm of soil. If rain does not occur or irrigation is unavailable incorporate chemical to a depth of 3-4cm. DO NOT throw untreated soil onto treated areas as this will reduce weed control. Use rates towards the higher end of the range on heavy soils or where a heavy grass population is expected.
Sorghum treated with Concep* II Sorghum Seed Safener	Bamyard grass, Crowsfoot grass, Liverseed grass, Lovegrass, Pigeon grass, Summer grass, Wandering Jew	NSW, ACT, Qld, NT, only	2-4L/ha	Pre-emergent. Apply before, at or immediately after planting and before crops and weeds have germinated using a low volume boom sprayer applying 60-120L of mixture per hectare. DO NOT apply to sorghum seed not previously treated with Concep II Sorghum Seed Safener. Rain or irrigation is necessary within 10 days of spraying to thoroughly wet the top 3-4cm of soil. If rain does not occur or irrigation is unavailable incorporate chemical to a depth of 3-4cm. Use rates towards the higher end of the range on heavy soils or where a heavy grass population is expected. DO NOT throw untreated soil onto treated areas as this will reduce weed control. Use the lower rate on sandy soils.
	Blackberry nightshade, Caltrop, Common thornapple, Mintweed, Needle burr, Noogoora burr, Pigweed, Redroot amaranth, Slim amaranth		2-4L/ha plus 2.5L of Flowable Nu-trazine Liquid	

CROP	WEEDS CONTROLLED	STATE	RATE	CRITICAL COMMENTS
Sugar cane (plant and ratoon)	Awnless barnyard grass, Barnyard grass, Bellvine, Billygoat weed (Blue top), Blackberry nighshade, Caltrop, Common sowthistle (Milk thistle), Common thornapple, Crowsfoot grass, Green Summer grass, Guinea grass, Mexican clover, Mintweed, Needle burr, Passionfruit vines, Pigeon grass, Pigweed, Redroot amaranth, Square weed, Star of Bethlehem, Summer grass, Wild rose	Southern Qld only (Bundaberg South)	2.2-2.9L/ha plus 2.7-3.6L/ha Flowable Nu-trazine Liquid or 1.5-2kg of Nu-trazine 900DF	Apply as a pre- or post-emergent application to crop. Apply once per year only. Application should be made to a moist soil before weeds and grasses have germinated. If conditions remain dry for a period of 10 days after spraying, irrigation or a shallow cultivation (2.5 cm) may assist results. DO NOT throw untreated soil on to treated areas as this will reduce weed control. In Southern Queensland, use rates toward the higher end of the range on heavy soils where a high grass population is expected. In Northern Queensland, application must be made to moist soil and rainfall or higher end of the range where high Green Summer grass population is expected. Where broadleaf weeds and grasses have emerged and are in the 2-4 true leaf stage, knockdown herbicides such as Primato*Z (2 L/ha), Nufarm Flowable Diuron (500g/L) at 1L/ha or Nuquat® 250 (1.2-1.6L/ha), applied as a post directed spray, can be added. Where broadleaf weeds only have emerged and are at the 2-4 true leaf stage then Amicide® 625 at 1.6L/ha should be added. In all cases add a non-ionic surfactant. Follow all instructions and restrictions on the Nu-trazine labels
		Northern Qld only (Mackay North)	2.9-3.6L/ha plus 3.6-4.5L/ha Flowable Nu-trazine Liquid or 2-2.5kg of Nu-trazine 900DF	
Sweet potatoes	Black pigweed, Redshank, Potato weed, Barnyard grass, Crowsfoot grass, Liverseed grass, Lovegrasses, Pigeon rass, Summer grass, Suppression of Evening primrose	NSW, ACT, Qld, Vic, NT, SA only	3L/ha	Pre-emergent. Apply immediately after transplanting before weeds have germinated. Sufficient irrigation to wet the soil through the weed zone should be applied within 24 hours. Further weed germination may occur following reilling due to exposure of untreated soil.
Tobacco (transplanted in field)	Barnyard grass, Lovegrass, Pigeon grass, Summer grass, and suppression of Common thornapple (<i>Datura stramonium</i>) and Nighthshades (<i>Solanum sarrachoides</i> and <i>S.gigrum</i>)	Vic, NSW only	3L/ha	Apply by low volume boom spray prior to transplanting and before weeds and grasses germinate. The seed bed should be fine and free of large clods. Thoroughly incorporate within 7 days of spraying using a rotary hoe or similar equipment set to work to a depth of 7.5cm. DO NOT use disc cultivating implements. An even coverage of the herbicide is essential. The herbicide needs to be activated by moisture through to the top 3-4cm of soil at the time of application or within 4 days of incorporation. This can be achieved by applying to moist soil, irrigating soon after incorporation or irrigating immediately after transplanting. Avoid boom overlaps as double treatment may cause growth suppression. Temporary crop retardation may be observed when soil conditions remain dry after application or where black root rot may occur.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

COTTON:

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED

GRAZING: DO NOT ALLOW LIVESTOCK TO GRAZE CROPS, COTTON STUBBLE OR BE FED GIN TRASH

CANOLA:

DO NOT GRAZE TREATED FIELDS FOR 10 WEEKS AFTER APPLICATION.

GREEN BEANS:

DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 10 WEEKS AFTER APPLICATION.

PASTURES:

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 13 WEEKS AFTER APPLICATION.

SWEET POTATOES:

DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 23 WEEKS AFTER APPLICATION.

WHEAT, BARLEY, OATS, TRITICALE:

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION.

FORAGE ALL OTHER CROPS EXCEPT SWEET POTATOES AND COTTON:

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 13 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

RESISTANCE WEEDS WARNING

GROUP	K	HERBICIDE
-------	----------	-----------

Bouncer Herbicide is a member of the amide group of herbicides with diverse sites of action. For weed resistance management this is Group K herbicide. Some naturally-occurring weed biotypes resistant to Bouncer Herbicide and other Group K herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Bouncer Herbicide or Group K herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm accepts no liability for any losses that may result from the failure of Bouncer Herbicide to control the resistant weeds. Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or a Nufarm representative.

USES

The product is a short residual pre-emergent herbicide for the control of many important annual grasses and broadleaf weeds in the crops nominated. The product does not control the important broadleaf weeds in maize, sweet corn and Concep II treated sorghum, but when mixed with Flowable Nu-trazine control of many important annual weeds and grasses is obtained without the problem of carryover of herbicides affecting succeeding crops. The product acts mainly through shoot (coleoptile) absorption. Its effectiveness depends on the occurrence of rainfall or irrigation to move it down into the top 3-4cm of soil, where it must be present at the time of weed germination.

MIXING

Add the required amount directly to the spray tank, fill with water and mix well.

COMPATIBILITY

Bouncer® is compatible with Convoy®, Convoy® DF, Nutron 900 DF, Flowable Nu-trazine Liquid, Nu-trazine 900 DF, Nugran®, Nuquat®, Roundup® CT, Roundup PowerMAX®.

WARNING – Cotton

Heavy rain following an application of Bouncer, together with below average soil temperatures, can result in crop injury, due to the inability of the germinating plant being able to metabolise the herbicide. Bouncer should not be used in situations where these conditions are likely to occur.

WARNING – Brassica crops

Crop retardation may occur where Bouncer is used on soils that have a low organic matter content and which contain more than 60% of fine sand and silt and less than 5% clay in the top 30cm. On these soil types, use rates of Bouncer towards the lower end of the rate range and apply onto moist soil. Irrigation after application should be limited to no more than 25mm.

Zero-tillage (Canola, Barley, Oats, Wheat, Triticale):

When sowing using knife points and zero tillage and leaving a seeding slot, DO NOT apply post seeding unless harrows (or similar) have filled in the seeding slot before application. Alternatively, apply Bouncer immediately before sowing.

DO NOT USE ON SOILS CONTAINING LESS THAN 5% CLAY IN THE TOP 30cm .

DO NOT USE MORE THAN ONCE PER YEAR.

PROTECTION OF CROPS, NATIVE AND OTHER NON TARGET PLANTS

DO NOT plant crops other than maize, sweet corn, tobacco, cotton, soybeans, sunflowers, peanuts, broccoli, brussel sprouts, cabbages, cauliflowers or Concep II treated sorghum within six months of application of Bouncer.

DRIFT WARNING

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pasture.

PROTECTION OF LIVESTOCK

DO NOT graze or cut sweet potato leaves for stock food.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

RE-ENTRY PERIOD

DO NOT enter treated area until spray has dried.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

For refillable containers:

Store in the closed, original container, in a cool, well ventilated area away from children, animals, food, feedstuffs, seed and fertilizers. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Envirodrum 110L Mini Bulk Returnable Container:

Store the original sealed Envirodrum in a cool well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Nufarm Australia Limited.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. Sensitive workers should use protective clothing. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Tel: Australia 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier or from Nufarm website - www.nufarm.com.au

In case of emergency: Phone 1800 033 498 Ask for shift supervisor. Toll free 24 hours.

CONDITIONS OF SALE

Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute.

Nufarm Australia Limited

ACN 004 377 780

103-105 Pipe Road

Laverton North Victoria 3026

Tel: (03) 9282 1000

Fax: (03) 9282 1001

© Bouncer, Envirodrum, Convoy, Nugran, Nuquat and Amicide are registered trademarks of Nufarm Australia Limited

© Roundup and Roundup PowerMAX are registered trademarks of Monsanto Technology LLC used under licence by Nufarm Australia Limited

* Other registered trademarks